

EARLY PALEOZOIC EVOLUTION OF THE PERI-GONDWANA PLATES

Wczesnopaleozoiczna ewolucja perygondwańskich płyt litosfery

Jan GOLONKA¹, Michał KROBICKI¹,
Paweł POPRAWA², Zbigniew PAUL³ & Andriej KHUDDOLEY⁴

¹AGH University of Science and Technology,
Faculty of Geology, Geophysics and Environmental Protection,
Department of General Geology, Environmental Protection and Geotourism;
al. Mickiewicza 30, 30-059 Krakow, Poland;
e-mail: jan_golonka@yahoo.com, krobicki@geol.agh.edu.pl

²Polish Geological Institute;
ul. Rakowiecka 4, 00-975 Warsaw, Poland;
e-mail: pawel.poprawa@pgi.gov.pl

³Polish Geological Institute, Carpathian Branch;
ul. Skrzatów 1, 31-560 Krakow, Poland;
e-mail: zbigniew.paul@pgi.gov.pl

⁴St. Petersburg State University;
University Nab. 7/9, St. Petersburg 199034, Russia;
e-mail: khudoley@AH3549.spb.edu

Treść: Przedstawiono główne etapy wczesnopaleozoicznej ewolucji perygondwańskich płyt litosferycznych w nawiązaniu do historii superkontynentu Rodinia, a zwłaszcza efektu neoproterozoicznego ryftowania, które doprowadziło do izolacji Gondwany, Laurencji, Syberii i Bałtyki. Wskazano z kolei na wczesnopaleozoiczne (późny kambry – wczesny ordowik) ryftowanie terranów awalońskich, które objęło m.in. północno-zachodnią i południową Polskę. Na tym tle zobrażowano również wczesnopaleozoiczne wydarzenia geotektoniczne w dalekowschodniej Azji.

Słowa kluczowe: Perygondwana, Rodinia, Avalonia, wczesny paleozoik

Key words: Peri-Gondwana, Rodinia, Avalonia, Early Paleozoic

PANNOTIA AND PERI-GONDWANA TERRANES

The Proterozoic history of the supercontinent is defined by two major orogenies. The Grenvillian orogeny around 1100 Ma is related to the formation of supercontinent Rodinia (Dalziel 1991, Hoffman 1991). The Cadomian/Pan-African orogeny is related to the assembly of the supercontinent Pannotia (Dalziel *et al.* 1994, Dalziel 1997, Golonka 2000, Golonka *et al.* 2006a, b)

around the Precambrian-Cambrian boundary. More than 500 hundred million years between these two events allows assuming two full Wilson orogenic cycles during this time. It allows also many different, speculative paleogeographic approaches, causing lively discussion.

The Cadomian orogeny caused the deformation and magmatic events of terranes from Iberia through Armorica. The Baltica (Eastern Europe) might have collided with the Cadomian part of Gondwana during the Vendian time causing deformation in the Timan area and proto-Uralian area. The Pechora-Timan belt (Roberts & Siedlecka 2002) and fragments of Ural, Novaya Zemlya and Taimyr are related to the Cadomian belt (Golonka 2000, 2002). At the same time rifting occurred along the other Baltica border, related perhaps to the opening of the Iapetus Ocean (Poprawa 2006). Laurentia rifted away from Pannotia along future Iapetus Ocean during Vendian time and along the Ouachita Ocean during Cambrian time (Golonka *et al.* 2006c). Pannotia supercontinent is not so badly constrained, however, it was short lived. Its history resembled somewhat the history of Pangea, which assembled finally during Early Jurassic and broke-up during Middle Jurassic time. One of the possible, speculative reconstructions is linking Central Asian Orogenic Belt terranes (Tuva?) with the Cadomian orogeny in Europe (Golonka *et al.* 2006b).

Both Gondwana and Baltica were included in the Pannotia supercontinent (Golonka *et al.* 2006a, b). The continents forming the core of Gondwana include South America, Africa, Madagascar, India, Antarctica and Australia. The location of numerous smaller continental blocks that bordered Gondwana is less certain. These smaller blocks were assembled during the Early Paleozoic time to Gondwana, forming the so-called Peri-Gondwanian zone. The following were adjacent to Gondwana at some time during the Paleozoic: Yucatan, Florida, Avalonia, central European (Cadomian) terranes between the Armorica and Bohemian Massif, Moesia, Iberia, Apulia and the smaller, southern European terranes, central Asian terranes (Karakum and others), China (several separate blocks), and the Cimmerian terranes of Turkey, Iran, Afghanistan, Tibet and Southeast Asia.

THE RIFTING OF AVALONIAN TERRANES

The south-dipping subduction developed along the central margin of Gondwana in Late Cambrian – Early Ordovician time. It caused the onset of rifting of the Avalonian terranes. Traditionally, the continent of Avalonia consists of northwestern and possibly southern Poland, some accreted terranes in the basement of East Carpathians and their foredeep, terranes in northern Germany, the Ardennes of Belgium and northern France, England, Wales, southeastern Ireland, the Avalon Peninsula of eastern Newfoundland, much of Nova Scotia, southern New Brunswick, and some coastal parts of New England. Brunovistulicum and Małopolska massives, parts of the Scythian platform, Dzurilla massif of Caucasus, terranes in Tian-Shan and adjacent parts of Kazakhstan and Southern Mongolia terrane could have constitute of the eastern extension of the Avalonia (Paul *et al.* 2003, Golonka *et al.* 2006b). The Avalonian plates probably started to rift from Gondwana and move towards Baltica in the Late Tremadocian and were in a drift stage by the Llanvirnian. This subduction caused also the formation of the vast Rheic Ocean. The Turkmen and Solonker oceans in Asia could constitute the eastern parts of this Rheic Ocean. The plates drifted northward toward the nucleus of Asia (Fig. 1).




Fig. 1. Plate tectonic map of the Middle Ordovician (plates position as of 472 Ma): 1 – oceanic spreading center and transform faults, 2 – subduction zone, 3 – thrust fault, 4 – normal fault, 5 – transform fault

Fig. 1. Mapa tektoniki płyt środkowego ordowiku (pozycja płyt 472 milionów lat temu): 1 – centrum spredingu oceanicznego i uskoki transformujące, 2 – strefa subdukcji, 3 – nasunięcie, 4 – uskoki normalny, 5 – uskoki przesuwczy

THE PALEOZOIC OROGENIES IN CENTRAL ASIA

The sedimentary sequences in the Gobi desert area in Mongolia as well as in the Chinese Inner Mongolia record the plate tectonic development of Central Asia. Collision between microcontinents (Salafrican orogeny) during Late Cambrian – Early Ordovician time in the Mongolia-Tuva area marked the onset of the formation of the Amuria (Mongolia) microcontinent. Relationship of eastern peri-Gondwana terranes and Avalonia plates remain unknown and speculative. It is not impossible that South China and Indochina plates also were rifted from Gondwana in the Ordovician. The uplift and volcanism support such a possibility. According to Shouxin & Yongyi (1991), the Ordovician conformably overlies the Cambrian over most of the South China plate. The northern part of the plate (Yangzi Platform) was covered with carbonates and mixed carbonate/clastic facies. The southern part of the plate is partially uplifted and partially covered by deep water synorogenic clastic deposits – more than 4000 m of weakly metamorphosed flysch, sandstones and graptolitic shales. Similar rocks formed on the margins of Indochina plate. They are known as Pa Ham formation (Ordovician-Silurian).

Late Silurian was the time of the major development of the Caledonian orogeny and final closure of the Iapetus (Fig. 2). The collision between Baltica and Greenland continued, marked by nappes in Norway and Greenland. After the complete closure of the Iapetus Ocean, the continents of Baltic, Avalonia, and Laurentia formed the continent of Laurussia (Ziegler 1989). This Caledonian orogeny and transpressional collision of Gondwana and Laurentia

was related to the formation of the enigmatic large supercontinent Oldredia, which existed during Early Devonian times, and included all major plates (Golonka 2000, Golonka *et al.* 2006b).




Fig. 2. Plate tectonic map of the Late Silurian (plates position as of 425 Ma): 1 – oceanic spreading center and transform faults, 2 – subduction zone, 3 – thrust fault, 4 – normal fault, 5 – transform fault

Fig. 2. Mapa tektoniki płyt późnego syluru (pozycja płyt 425 milionów lat temu): 1 – centrum spredin-gu oceanicznego i uskoc transformujący, 2 – strefa subdukcji, 3 – nasunięcie, 4 – uskoc normalny, 5 – uskoc przesuwczy

The above mentioned events indicate possible connection of Asiatic plates with Oldredia. During Middle Devonian times, rifting of continental margins of Oldredia led to disassembly of the supercontinent (Golonka 2000, 2006b). The collision of the North China plate and closure of Solonker ocean (Şengör & Natalin 1996) in Permian time concluded the orogenic process.

This research has been in part financially supported by the AGH University of Science and Technology in Kraków, grant No. 11.11.140.447.

REFERENCES

- Dalziel I.W.D., 1991. Pacific margins of Laurentia and east Antarctica-Australia as a conjugate rift pair. Evidence and implications. *Geology*, 19, 598–601.
- Dalziel I.W.D., 1997. Neoproterozoic-Paleozoic geography and tectonics; review, hypothesis, environmental speculation. *Geological Society of America Bulletin*, 109, 16–42.
- Dalziel I.W.D., Dalla Salda L.H & Gahagan L.M., 1994. Paleozoic Laurentia-Gondwana interaction and the origin of the Appalachian-Andean mountain system. *Geological Society of America Bulletin*, 106, 2, 243–252.
- Golonka J., 2000. *Cambrian-Neogene plate tectonic maps*. Wydawnictwa Uniwersytetu Jagiellońskiego, Kraków, 1–125.

- Golonka J., 2002. Plate-tectonic maps of the Phanerozoic. In: Kiessling W., Flügel E. & Golonka J. (eds), Phanerozoic reef patterns. *SEPM Special Publication*, 72, 21–75.
- Golonka J., Gahagan L., Krobicki M., Marko F., Oszczytko N. & Ślącza A., 2006a. Plate Tectonic Evolution and Paleogeography of the Circum-Carpathian Region. In: Golonka J. & Picha F. (eds), The Carpathians and their foreland: Geology and hydrocarbon resources. *American Association of Petroleum Geologists, Memoir*, 84, 11–46.
- Golonka J., Krobicki M., Pająk J., Nguyen Van Giang & Zuchiewicz W., 2006b. *Global plate tectonics and paleogeography of Southeast Asia*. Faculty of Geology, Geophysics and Environmental Protection, AGH University of Science and Technology; Arkadia, 1–128.
- Golonka J., Ślącza A. & Picha F., 2006c. The Western Carpathians and Ouachitas: A comparative study of geodynamic evolution. In: Golonka J. & Picha F. (red.), The Carpathians and their foreland: Geology and hydrocarbon resources, *American Association of Petroleum Geologists, Memoir*, 84, 561–584.
- Hoffman P.E., 1991. Did the breakout of Laurentia turned Gondwanaland inside-out? *Science*, 252, 1409–1412.
- Paul Z., Golonka J., Wójcik A. & Khudoley A., 2003. The Iapetus Ocean, Rheic Ocean and Avalonian Terranes in Central Asia. *Geolines*, 16, 79–80.
- Poprawa P., 2006. Neoproterozoic break-up of the supercontinent Rodinia/Pannotia recorded by development of sedimentary basins at the western slopes of Baltica. In: Matyja H. & Poprawa P. (red.), Facies, Tectonics and Thermal evolution of the Pomeranian sector of Trans-European Suture Zone and adjacent areas, *Prace Państwowego Instytutu Geologicznego*, 136, 165–188 (In Polish, English summary).
- Roberts D. & Siedlecka A., 2002. Timanian orogenic deformation along the northeastern margin of Baltic, northwest Russia and northeast Norway, and Avalonian-Cadomian connections. *Tectonophysics*, 352, 169–184.
- Şengör A.M.C. & Natalin B.A., 1996. Paleotectonics of Asia: fragment of a synthesis. In: Yin An & Harrison T.M. (red.), *The Tectonic Evolution of Asia*, Cambridge University Press, 486–640.
- Ziegler P.A., 1989. *Evolution of Laurussia*. Kluwer Academic Publishers, Dordrecht, Netherlands, 1–102.