

## POLISH SCIENTISTS PROFILE

Zbigniew W. ENGEL\*

### PROFESSOR WOJCIECH RUBINOWICZ'S CONTRIBUTION TO THE DEVELOPMENT OF ACTIVE METHODS OF VIBRATION CONTROL

#### SUMMARY

World known Polish scientist, physicist Professor Wojciech Rubinowicz (1889–1974) contributed immensely to the development of the theory of active methods of noise reduction. Prof. W. Rubinowicz was dealing mainly with optical phenomena, however it should be emphasised, that light waves and acoustic phenomena can be described by similar theoretical equations. Already in 1917 W. Rubinowicz developed the theory of the edge wave, which later on was named the Rubinowicz Edge Wave. He proved the equivalence of Fresnel's and Young's ideas concerning diffraction phenomena. He developed the mathematical description of the Kirchhoff's diffraction theory. Phenomena of mutual interference of light waves incident on screen edge and the effect in the form of wave compensation in certain zones were described. An incident wave on the screen surface (edge) can compensate itself with the wave of the same structure but phase shifted by  $\pi$  angle, which can cause the total extinction of light. Works of the Polish scientist, which constituted the new approach to the wave phenomena can be considered the starting point in the development of the active method theory.

**Keywords:** edge wave, interference, active methods of noise reduction



#### WKŁAD PROFESORA WOJCIECHA RUBINOWICZA W ROZWÓJ METOD AKTYWNYCH

Światowej sławy polski naukowiec, fizyk, Profesor Wojciech Rubinowicz (1889–1974) przyczynił się ogromnie do opracowania teorii aktywnych metod redukcji hałasu. Prof. W. Rubinowicz zajmował się głównie zjawiskami optycznymi, jednakże należy podkreślić, że fale światła i zjawiska akustyczne mogą być opisywane przez zbliżone zależności teoretyczne. Już w roku 1917 W. Rubinowicz opracował teorię fali brzegowej, która otrzymała później nazwę teorii fali brzegowej Rubinowicza. Dowiódł równoważności idei Fresnela i Younga dotyczących zjawiska dyfrakcji. Opracował opis matematyczny teorii dyfrakcji Kirchhoffa. Opisał zjawisko wzajemnej interferencji fal światła zachodzącej na krawędzi ekranu oraz efekt kompensacji fal w pewnych obszarach. Fala wypadkowa na płaszczyźnie ekranu (krawędzi) kompensuje się z falą o tej samej strukturze, ale w fazie przesuniętej o kąt  $\pi$ , co powoduje jej całkowite wygaśnięcie. Praca polskiego naukowca, która stanowi nowe podejście do zjawisk falowych, może być rozważana jako punkt wyjściowy w rozwoju teorii metod aktywnych.

**Słowa kluczowe:** fala brzegowa, interferencja, metody aktywne redukcji hałasu

#### 1. INTRODUCTION

The recent twenty years have witnessed a rapid development of active methods of vibration and noise control. Research centers world-wide focus on active methods and the results have left the laboratory stage and new practical applications are being found. Underlying the active vibration control methods is the principle that sounds and vibrations to be reduced are compensated by noise and vibrations from additional sources. Of major importance is the fact that they incorporate an external source of energy. Controlled systems might supply or absorb vibro-acoustic energy in a predetermined manner, from any point of the system (device). The first concepts of active vibration reduction methods appeared in the late 19<sup>th</sup> century, stemming from the interference theory propounded in 1690 by Christiaan Huygens (1629–1695).

A well-known English researcher Thomas Young (1773–1829) in his work *Course of Lecture on Natural Philosophy*

(1807) remarked that „Two identical systems of waves leaving the nearby points will cancel at certain points or reinforce their actions at others” [15]. The basic concepts of noise reduction through the interference of the primary and secondary waves were given by lord Rayleigh (1842–1919). It is worthwhile to mention that the problem utilizing the phenomenon of interference does involve pure interference as in this case interference is achieved for waves propagating in opposite directions, while compensation (reduction) applies occurs in waves moving in the same direction (and hence the terms: “negative interference” and “destructive interference”).

The concept suggested by Rayleigh was utilized by a French scholar – Henry Coala in 1932 and by a German Paul Lueg (1933) while developing new patents in the area of active noise reduction.

\* AGH University of Science and Technology, Krakow, Poland

Underlying the active methods of sound compensation are theoretical backgrounds provided by M. Jessel. In his works Jessel recalls the effect of concavity, stemming from the well-known Rubinowicz's paradox.

## 2. PROFESSOR WOJCIECH RUBINOWICZ'S BIOGRAPHY

The outstanding Polish physicist was born in 1889 in Sadagora, near Czernowitz, in the family of a pharmacist Damian and Małgorzata, nee Brodowska. From 1900 to 1908 he attended the German high school in Czernowitz, then moved on to study at the University of Czernowitz, in 1914 he received a Ph.D. degree in mathematical physics. In 1916 he left for Munich, where he worked as Prof. Arnold Sommerfeld's assistant. On returning to Czernowitz he became a *Privatdozent*, having presented the dissertation *Die Beugungswelle in der Kirchhoffschen Theorie der Beugungsercheinung*. Invited by Niels Bohr, he went to Copenhagen in 1920, then took an appointment as a professor of theoretical physics in Ljubljana. In 1922 he became a full professor of theoretical physics at the Polytechnic Institute of Lvov.

In the academic year 1935/26 he was the Dean of this Faculty. In the period 1935–37 he worked as the head of the Department of General Mechanics at the Faculty of Engineering of the Polytechnic Institute of Lvov. From 1937 he was a professor of theoretical physics at the John Casimir University of Lvov.

In 1946 he was appointed the head of the Department of Theoretical Mechanics at the Warsaw University. From 1953 he held the position of the head of the Department of Optics and Mechanics, until his retirement. At the same time he also worked in the Institute of Physics of the Polish Academy of Sciences. In 1931 he was elected a member of the Polish Academy of Learning. From 1952 he was a member of the Polish Academy of Sciences.

In the years 1949–1952 and 1961–1974 he was the President of the Polish Society of Physics. Professor Wojciech Rubinowicz was a theoretical physicist, his major research interests included quantum theory of radiation, theory of diffraction and mathematical physics.

Professor Wojciech Rubinowicz was a theoretical physicist, his research work focused on quantum theory of radiation, the theory of diffraction and mathematical physics. His outstanding achievements, which made him world famous, included the discovery of rules for quantum transitions in atoms, the theory of electromagnetic quadruple radiation and works on light diffraction. He was the author of a great number of books, including monographs and textbooks.

Wojciech Rubinowicz died in Warsaw in 1974.

## 3. ANALYSIS OF WAVE MOTION IN THE WORKS OF W. RUBINOWICZ

Professor Rubinowicz's major research interest was the theory of diffraction. In 1917 the work *Die Beugungswelle in der Kirchhoffschen Theorie der Beugungsercheinung* appeared, where he transformed the Kirchhoff diffraction inte-

gral into what has been known as Rubinowicz representation. The Kirchhoff diffraction theory might be interpreted in two ways. The first interpretation utilises a concept of an edge wave, first proposed by Thomas Young, further developed by Prof. Rubinowicz, who also provided the theoretical backgrounds. The other interpretation involves the Fresnel's diffraction model (Augustin Fresnel 1788–1827), utilising the concept of zones. For a long time that approach was considered to be the only one.

Thomas Young claimed that diffraction occurs as a result of superimposition of a wave propagating in a free space with that appearing in the presence of a screen and hence can be treated as a wave reflected from the diffracting edge. After a while, however, Young abandoned his diffraction model in favour of the Fresnel's theory of zones. Nearly one hundred years later, Wojciech Rubinowicz showed that the Kirchhoff diffraction field can be divided. Rubinowicz decomposed the Kirchhoff wave into a geometrical –optical wave and a diffraction wave. In his work of 1924 he showed that in the first approximation the diffraction wave might be treated as a wave reflected from the diffraction edge. The starting point for Rubinowicz's work was the Helmholtz equation

$$\nabla^2 u + k^2 u = 0 \quad (1)$$

Wave motion was studied in the infinite space containing opaque bodies, referred to as screens. On their surface  $F$  the wave function  $u(P)$  should satisfy one of the boundary conditions:  $u = 0$  or  $\delta u / \delta n = 0$ . The physical interpretation of boundary conditions is associated with the physical meaning of the wave function. If the function  $u(P)$  is a velocity potential, then  $\mathbf{v} = \text{grad } u(P)$  denotes velocity and  $\mathbf{n} \cdot \text{grad}_P u(P) = \delta u / \delta n = 0$  indicates that the normal velocity component disappears and hence the screen surface should be immobile. In acoustics, such screens are referred to as "hard". In the case of the boundary condition, when  $u(P) = 0$  the screen is called "soft". On the surface of such screens the acoustic pressure tends to disappear. Let us consider an incident wave radiated by an isotropic and point light source  $L$  (Fig. 1).




Fig. 1. A rough sketch for Rubinowicz's analysis [12]

For this specific case we get

$$u(P) = \frac{e^{ikr_{LP}}}{r_{LP}} \quad (2)$$



portance was that part of the screen that would be shaded on the lit side because of wave-like shape of the diffracting screen. The screen can be deformed in any way whilst diffraction phenomena will not change. Such screen is called a “wavy” screen or Kirchhoff wavy screen. Rubinowicz observed step-like motion of a wave that on a wavy screen. The proposed interpretation is that such screens absorb incident waves. In the case of a screen formed by an arbitrary surface, it may happen that light rays, i.e. half-lines originating from the source should pass through the screen twice or even more times. Thus the behaviour of incident light on the Kirchhoff screen seems paradoxical. This phenomenon is known as Rubinowicz’s paradox. M. Jessel, one of the researchers who provided theoretical backgrounds of active methods of noise reduction, states in his works [6, 7] that the concavity effect originates from Rubinowicz’s paradox. While analysing the area in which an acoustic field occurs, one part of this field supports the Huygen’s source conformity, while the other supports the active source reduction. Rubinowicz predicted this effect on a Kirchhoff’s screen.

## 5. SUMMING UP

Kirchhoff’s theory of diffraction was studied theoretically by Wojciech Rubinowicz. The founders of active methods of noise reduction utilised the achievements of this outstanding Polish scholar who focused chiefly on optical phenomena, not on acoustics. Both optical and acoustic effects are wave-like phenomena governed by similar equations, that is why the works by this Polish scholar who started the new approach to the studies of waves might be regarded as the starting point, as theoretical backgrounds underlying the active methods. Unfortunately, major achievements of Polish scholars and their contribution to research achievements world-wide tend to be forgotten. Many authors specialising in active methods do not know the works of Prof. Wojciech Rubinowicz, and his works are rarely quoted. The main purpose of this paper is to recall the research achievements of Prof. Rubi-

nowicz and his contribution to the development of active methods.

*This study is a part of a research program no 4 T07C 00830 supported by the Ministry of Education and Science in Poland.*

## REFERENCES

- [1] Angevine O.L., Jessel M.J.M.: *Active Systems for Global Attenuation of Noise*. 1977(maszynopis ofiarowany przez Autorów)
- [2] Engel Z.: *Aktywna redukcja drgań i hałasu*. Mater. XI Sympozjum „Drgania w układach fizycznych”, Poznań 1984, pp. 124–125
- [3] Engel Z.: *Rozwój aktywnych metod zwalczania hałasu*. Mechanika, Kwartalnik AGH, t. 12, z. 1, 1993, pp. 9–28
- [4] Engel Z., Makarewicz G., Morzyński L., Zawieska M.: *Metody aktywne redukcji hałasu*. Warszawa, Wyd. CIOP 2001
- [5] Jessel M.J.M.: *Contribution aux theories du Principe de Huygens et de la Diffraction*. Université de Paris1962 (D.Sc. Thesis)
- [6] Jessel M.J.M.: *Noise Control by Means of Active Absorbers, part I: Theory*. Proc. Noise Control Conference, Warszawa 1976, pp. 221–224
- [7] Jessel M.J.M.: *Active Control in the Epistemology of Science and Technology*. Proc. Intern. Symp. ACSV, Tokyo 1991, pp. 235–242
- [8] Mangiante G.: *Noise Control by Means of Active Absorbers, part II: Active Sound Absorbers in a long Duct*. Proc. Noise Control Conference, Warszawa 1976, pp. 279–282
- [9] Rubinowicz W.: *Die Beugungswelle in der Kirchhoffschen Theorie der Beugungserscheinungen*. Ann. Physik, (4) 53, 1917, pp. 257–278
- [10] Rubinowicz W.: *Zur Kirchhoffschen Beugungstheorie*. Ann. Physik, (4),73, 1924, pp. 339–364
- [11] Rubinowicz W.: *Die Beugungswelle in der Kirchhoffschen Theorie der Beugung*. Warszawa, PWN 1957
- [12] Rubinowicz W.: *Teoria dyfrakcji Kirchhoffa i jej interpretacja na podstawie poglądów Younga*. Wrocław, Wyd. Ossol. 1972
- [13] Tokhi M.O., Leitch R.R.: *Active Noise Control*. Oxford, Clarendon Press 1992
- [14] Skudrzyk E.: *Die Grundlagen der Akustik*. Wien, Springer Verlag 1954
- [15] Wróblewski A.K.: *Historia fizyki*. Warszawa, PWN 2006
- [16] Petykiewicz J.: *Optyka falowa*. Warszawa, PWN 1986
- [17] Laue M.: *Interferenz und Beugung elektromagnetischer Wellen (mit Ausnahme der Roentgenstrahlen)*. Handb. Exp. Phys., 18, 211, Leipzig, Akad. Verl. 1928