

## ON A PROBLEM OF GEVORKYAN FOR THE FRANKLIN SYSTEM

Zygmunt Wronicz

*Communicated by P.A. Cojuhari*

**Abstract.** In 1870 G. Cantor proved that if  $\lim_{N \rightarrow \infty} \sum_{n=-N}^N c_n e^{inx} = 0$  for every real  $x$ , where  $\bar{c}_n = c_n$  ( $n \in \mathbb{Z}$ ), then all coefficients  $c_n$  are equal to zero. Later, in 1950 V. Ya. Kozlov proved that there exists a trigonometric series for which a subsequence of its partial sums converges to zero, where not all coefficients of the series are zero. In 2004 G. Gevorkyan raised the issue that if Cantor's result extends to the Franklin system. The conjecture remains open until now. In the present paper we show however that Kozlov's version remains true for Franklin's system.

**Keywords:** Franklin system, orthonormal spline system, trigonometric system, uniqueness of series.

**Mathematics Subject Classification:** 42C10, 42C25, 41A15.

### 1. INTRODUCTION

In 1870 G. Cantor proved in [5] the following theorem.

**Theorem 1.1.** *If  $\lim_{N \rightarrow \infty} \sum_{n=-N}^N c_n e^{inx} = 0$  for every real number  $x$ , where  $\bar{c}_n = c_n$ , then  $c_n = 0$  for  $n \in \mathbb{Z}$ .*

In 1950 V. Ya. Kozlov [13] proved that there exists a trigonometric series  $\sum_{n=0}^{\infty} a_n \cos nx + b_n \sin nx$  such that a subsequence of its partial sums is convergent to zero for  $x \in \mathbb{R}$  and not all the coefficients are equal to zero. By the Gram-Schmidt process to the Schauder basis Ph. Franklin constructed an orthonormal system of continuous piecewise linear functions with dyadic knots. It is an orthonormal Schauder basis in the space  $C[0, 1]$ , and also in the space  $L^2[0, 1]$ . In 1963 Z. Ciesielski [6] proved exponential type estimates for Franklin functions. Since then, it has been studied by many authors from different points of view. The Franklin system is an unconditional basis in  $L^p[0, 1]$ ,  $1 < p < \infty$  (S.V. Bochkarev [4]) and  $H^1[0, 1]$  (P. Wojtaszczyk [16]).

It has been used to prove the existence of a basis in the space  $C^1([0, 1] \times [0, 1])$  (independently by Z. Ciesielski [7] and by S. Schonefeld [15]).

In the case of Haar and Walsh systems the uniqueness problem was solved by F.G. Aratunyan and A.A. Talalyan in [2]. In 2004 G. Gevorkyan [12] raised the issue if Cantor’s result extends to the Franklin system. This conjecture remains open until now, and also for the periodic Franklin system which is obtained similarly to the nonperiodic case. The periodic Franklin system has been used to construct a basis in a disc algebra (S.V. Bochkarev [4]).

In 1938 J. Marcinkiewicz [14] obtained the following result.

**Theorem 1.2.** *For any complete in  $L^2[0, 1]$  an orthonormal system  $\{\varphi_n\}_{n=1}^\infty$  there exists a nonzero series  $\sum_{n=1}^\infty c_n \varphi_n(x)$  with a subsequence of its partial sums converging to zero almost everywhere in the interval  $[0, 1]$ .*

The purpose of the paper is to prove the ensuing result.

**Theorem 1.3.** *There is a nonzero Franklin series  $\sum_{n=0}^\infty a_n f_n(x)$  such that a subsequence  $\{s_{n_k}(x)\}_{k=1}^\infty$  of its partial sums is convergent to zero in the interval  $[0, 1]$ .*

## 2. PROOF OF THEOREM 1.3

It suffices to prove the theorem for the Franklin system for the interval  $I = [-1, 1]$ . We use odd functions in the proof. Because of it and the simplicity of calculations, we shall consider the Franklin system for this interval. Consider the following sequence  $\{\Delta_n\}_{n=1}^\infty$  of dyadic partitions of the interval  $I$ :  $\Delta_n = \{s_{n,i}\}_{i=0}^n$ ,  $s_{1,0} = -1$ ,  $s_{1,1} = 1$ ,

$$s_{n,i} = \begin{cases} \frac{i}{2^\mu} - 1 & \text{for } i = 0, 1, \dots, 2\nu, \\ \frac{i-\nu}{2^{\mu-1}} - 1 & \text{for } i = 2\nu + 1, \dots, n \end{cases} \tag{2.1}$$

for  $n = 2^\mu + \nu$ ,  $\mu = 0, 1, \dots$ ,  $\nu = 1, 2, \dots, 2^\mu$ .

We can obtain the Franklin system by means of cubic splines. We put

$$f_0 = \frac{1}{\sqrt{2}}, \quad f_1 = \sqrt{\frac{3}{2}}x.$$

Let  $g_n$  be a cubic spline with respect to the partition  $\Delta_n$ , i.e.  $g_n \in C^2(I)$  and it is a polynomial of degree at most 3 in each interval  $[s_{n,i-1}, s_{n,i}]$ . We assume that  $g_n(s_{n-1,j}) = 0$  for  $j = 0, 1, \dots, n - 1$  and  $g_n(s_{n,k}) = 1$  for  $s_{n,k} \in \Delta_n \setminus \Delta_{n-1}$  with  $g'_n(\pm 1) = 0$ . The spline  $g_n$  is unique. For the proof we refer to [1]. It is similar to that of the uniqueness of a spline  $S_n$  below. Because of its crucial role in the proof of the main result, we shall give it in detail. Integrating by parts, we check that the system  $\{f_n\}_{n=0}^\infty$ , where

$$f_n = \frac{g''_n}{\|g''_n\|}, \quad \|g''_n\|^2 = \int_{-1}^1 [g''_n(x)]^2 dx, \quad n = 2, 3, \dots,$$

is orthonormal in the interval  $I$  (see [1, 17, 18]).

Let

$$F(x) = \begin{cases} -1 & \text{for } x \in [-1, 0), \\ 0 & \text{for } x = 0, \\ 1 & \text{for } x \in (0, 1]. \end{cases} \tag{2.2}$$

We define the following sequence of functions  $\{S_n(x)\}_{n=0}^\infty$ .  $S_n$  is a cubic spline interpolating the function  $F$  on  $\Delta_n$ , i.e.  $S_n(s_{n,i}) = F(s_{n,i})$ ,  $S'_n(\pm 1) = 0$ . Let

$$\begin{aligned} M_i &= S''_n(s_{n,i}), \quad h_i = s_{n,i} - s_{n,i-1}, \quad y_i = F(s_{n,i}), \quad i = 0, 1, \dots, n, \\ y'_0 &= F'(-1), \quad y'_n = F'(1), \quad d_0 = \frac{6}{h_1} \left( \frac{y_1 - y_0}{h_1} - y'_0 \right), \quad d_n = \frac{6}{h_n} \left( y'_n - \frac{y_n - y_{n-1}}{h_n} \right), \\ d_j &= \frac{6}{h_j + h_{j+1}} \left( \frac{y_{j+1} - y_j}{h_{j+1}} - \frac{y_j - y_{j-1}}{h_j} \right), \quad \lambda_j = \frac{h_{j+1}}{h_j + h_{j+1}}, \\ \mu_j &= 1 - \lambda_j, \quad j = 1, \dots, n - 1, \quad \text{and } \lambda_0 = \mu_n = 1, \quad \lambda_n = \mu_0 = 0. \end{aligned}$$

$S_n$  is piecewise linear. Hence for  $x \in [s_{n,i-1}, s_{n,i}]$ ,  $i = 1, \dots, n$ , we get

$$\begin{aligned} S''_n(x) &= M_{i-1} \frac{s_{n,i} - x}{h_i} + M_i \frac{x - s_{n,i-1}}{h_i}, \\ S'_n(x) &= -M_{i-1} \frac{(s_{n,i} - x)^2}{2h_i} + M_i \frac{(x - s_{n,i-1})^2}{2h_i} + C_i, \\ S_n(x) &= M_{i-1} \frac{(s_{n,i} - x)^3}{6h_i} + M_i \frac{(x - s_{n,i-1})^3}{6h_i} + C_i(x - s_{n,i-1}) + D_i, \end{aligned}$$

where  $C_i$  and  $D_i$  are some constants. Using the interpolation conditions, we obtain

$$S'_n(x) = -M_{i-1} \frac{(s_{n,i} - x)^2}{2h_i} + M_i \frac{(x - s_{n,i-1})^2}{2h_i} + \frac{y_i - y_{i-1}}{h_i} - (M_i - M_{i-1}) \frac{h_i}{6}.$$

The function  $S'_n$  is continuous on the interval  $[-1, 1]$ . Hence  $S'_n(s_{n,i-}) = S'_n(s_{n,i+})$ ,  $i = 1, \dots, n - 1$ , and we obtain

$$\mu_j M_{j-1} + 2M_j + \lambda_j M_{j+1} = d_j, \quad j = 0, 1, \dots, n, \quad \text{where } M_{-1} := M_{n+1} := 0. \tag{2.3}$$

The matrix of this system has dominated the main diagonal. Hence, by the Gerschgorin theorem, the spline  $S_n$  is unique ([1]) and  $S''_n(x) = \sum_{k=0}^n a_k f_k(x)$  with not all  $a_n = 0$ .

Let  $n = 2^{k+1} = 2m$ . We have  $\mu_j = \lambda_j = \frac{1}{2}$ ,  $d_j = 0$  for  $j = 0, 1, \dots, m - 2, m, m + 2, \dots, 2m$  and  $d_{m-1} = 3m^2 = -d_{m+1}$ .

Further we need the following result.

**Theorem 2.1** (S. Demko [10]). *Let  $A = [a_{ij}]$  be an  $m \times m$  band matrix with bandwidth  $k$ , i.e.  $a_{ij} = 0$  for  $|i - j| \geq k$  and  $\|A\|_p$  denote matrix  $l_p$ -norm. Suppose there exist  $1 \leq p \leq \infty$  and  $M$  such that  $\|A\|_p \leq 1$  and  $\|A^{-1}\|_p \leq M$ . Then there exist constants  $C = C_{k,M}$  and  $q = q_{k,M}$ ,  $0 < q < 1$ , such that for  $A^{-1} = [b_{ij}]$*

$$|b_{ij}| \leq C q^{|i-j|}, \quad i, j = 1, \dots, m.$$

Let  $A = [a_{ij}]$  be the matrix of the first  $m$  equations of the system (2.3) and  $B = [b_{ij}] = A^{-1}$ . By means of Theorem 2.1 we prove that there exist constants  $C$  and  $0 < q < 1$  such that

$$|M_{m-k}| \leq Cm^2q^k, \quad k = 1, 2, \dots \tag{2.4}$$

Then for  $k \geq \sqrt{m}$

$$|M_{m-k}| \rightarrow 0 \quad \text{as } m \rightarrow \infty.$$

Let  $0 < \alpha < 1$ . We choose  $m$  such that  $\alpha > \frac{1}{\sqrt{m}}$ . Hence for each  $\varepsilon > 0$  there exists  $\delta > 0$  such that  $|M_{m-k}| < \varepsilon$  for all integers  $m > \delta$  and  $k \in [\sqrt{m}, m]$ , and

$$M_{m-k} = S''_{2m}(s_{n,m-k}) = S''_{2m}\left(-\frac{k}{m}\right), \quad \frac{k}{m} \geq \frac{1}{\sqrt{m}}.$$

Since  $S''_{2m}$  is piecewise linear, then  $|S''_{2m}(x)| < \varepsilon$  for  $x \in [-1, \alpha] \cup [\alpha, 1]$ , and we obtain

$$\lim_{m \rightarrow \infty} S''_{2m}(x) = \lim_{m \rightarrow \infty} s_{2m}(x) = \lim_{m \rightarrow \infty} \sum_{k=0}^{2m} a_k f_k(x) = 0.$$

The convergence is uniform on the intervals  $[-1, -\alpha]$  and  $[\alpha, 1]$  ( $0 < \alpha < 1$ ).

Now let  $n = 2^\mu + \nu$  and  $\nu = 2^{\mu-1} = m$ . In this case the quantities  $M_j$  satisfy the following system of equations

$$\begin{cases} 2M_0 + M_1 = 0, \\ M_{j-1} + 4M_j + M_{j+1} = d_j, \quad j = 1, \dots, 3m - 1, \\ M_{3m-1} + 2M_{3m} = 0, \end{cases} \tag{2.5}$$

where  $d_j = 0$  for  $j = 1, \dots, 2m - 2, 2m + 2, \dots, 3m - 1$ , and

$$d_{2m-1} = 24m^2, \quad d_{2m} = -8m^2, \quad d_{2m+1} = -6m^2.$$

To obtain  $M_{2m}$ , we use the Cramer formulas for the system (2.5). After elementary calculations, we write the nominator and the denominator in a form of block triangular matrices and we prove that  $|M_{2m}| \rightarrow \infty$  as  $m \rightarrow \infty$ .

**Remark 2.2.**

1) Theorem 1.1 remains true for the periodic Franklin system, and its proof is analogous to that above.

2) Let

$$g_0(x) = 1, \quad g_1(x) = x, \quad g_n(x) = \int_0^x f_{n-1}(t)dt,$$

where  $f_n$  is the  $n$ -th Franklin function,  $n = 2, 3, \dots$ . Applying the Gram-Schmidt process to this system, we obtain the Ciesielski orthonormal system of splines of degree 2. Proceeding in the same way, we get an orthonormal system of splines of degree 3. Repeating this process, we may obtain an orthonormal system of splines

of degree  $k$ , where  $k = 1, 2, \dots$ . These systems were introduced by Z. Ciesielski [8]. We may prove Theorem 1.1 for them (cf. [8, 18]). In this case we interpolate the function  $F$  from Theorem 1.1 by splines of odd degree and the function  $G(x) = 1$  for  $x \in [-1, 1] \setminus \{0\}$  and  $G(0) = 0$  by splines of even degree.

### 3. THE GEVORKYAN PROBLEM FOR THE SPACE $V$ OF FUNCTIONS

$f \in C[0, 1]$  WITH  $f(0) = 0$

Let  $\{\Delta_n\}_{n=1}^\infty$ ,  $\Delta_n = \{0 = x_0 < x_1 < \dots < x_n = 1\}$ , be a sequence of partitions of the interval  $[0, 1]$  with  $\Delta_{n-1} \subset \Delta_n$  ( $n = 1, 2, \dots$ ) and

$$\lim_{n \rightarrow \infty} \max_{1 \leq i \leq n} (x_i - x_{i-1}) = 0.$$

We define the Franklin system for the space  $V$  with the sequence  $\{\Delta_n\}_{n=1}^\infty$  as follows:  $f_1 = \sqrt{3}x$ ,  $g_{n+1}$  is a cubic spline such that  $g_{n+1}(x_j) = 0$  for  $x_j \in \Delta_n$  and  $g_{n+1}(x_k) = 1$  for  $x_k \in \Delta_{n+1} \setminus \Delta_n$ ,  $g''_n(0) = g'_n(1) = 0$ . Then we put

$$f_{n+1}(x) = \frac{g''_{n+1}}{\|g''_{n+1}\|}, \quad n = 1, 2, \dots$$

The system  $\{f_n\}_{n=1}^\infty$  is orthonormal in the space  $L^2[0, 1]$  and it forms a basis in the space  $V$  (see [1, 6, 9, 10]). Now we define the sequence  $\{S_n(x)\}_{n=1}^\infty$  of cubic splines such that  $S_n(x_k) = 1$  for  $x_k \in \Delta_n$ ,  $k = 1, 2, \dots, n$ ,  $S_n(0) = S''_n(0) = S'_n(1) = 0$ . As in Theorem 1.1, we may prove that  $S_n$  is unique.  $S''_n(x) = \sum_{k=1}^n a_k f_k(x)$ ,  $a_1 = -\sqrt{3}$ .

Let  $M_j = S''_n(x_j)$ ,  $j = 1, \dots, n$ . Using the Demko theorem, as in the proof of Theorem 1.1, we prove that for any  $\alpha \in (0, 1)$  the sequence  $\{S''_n(x)\}_{n=1}^\infty$  is uniformly convergent to zero in the interval  $[\alpha, 1]$ . Hence

$$\lim_{m \rightarrow \infty} S''_m(x) = \sum_{n=1}^\infty a_n f_n(x) = 0 \text{ on } [0, 1] \text{ and } a_1 \neq 0,$$

and thus we have proved the following result.

**Theorem 3.1.** *There is a nonzero Franklin series  $\sum_{n=1}^\infty a_n f_n$  in the space  $V$  with the sequence of partial sums converging to zero in the interval  $[0, 1]$ .*

### 4. PROBLEMS

Let  $\{\Delta_n\}_{n=1}^\infty$  be a given sequence of partitions of the interval  $I = [-1, 1]$ ,  $\Delta_n = \{-1 = t_{n,0} < t_{n,1} < \dots < t_{n,n} = 1\}$  with  $\Delta_n \subset \Delta_{n+1}$ , i.e. each point of  $\Delta_n$  is a point of  $\Delta_{n+1}$ . We assume that

$$\lim_{n \rightarrow \infty} \max_{1 \leq i \leq n} (t_{n,i} - t_{n,i-1}) = 0.$$

We define an orthonormal system of piecewise functions with respect to the sequence of partitions  $\{\Delta_n\}_{n=1}^{\infty}$  analogous to the Franklin system. We call it the general Franklin system (see [12]). We have the following problems: to prove a counterpart of Theorem 1.3 for

- a) any general Franklin system (see [17]),
- b) any complete orthonormal system in the space  $C[0, 1]$ .

## REFERENCES

- [1] J.H. Ahlberg, E.N. Nilson, J.L. Walsh, *The theory of splines and their applications*, Academic Press, New York-London, 1967.
- [2] F.G. Arutyunyan, A.A. Talalyan, *Uniqueness of Series in Haar and Walsh systems*, Izv. Akad. Nauk SSSR, Ser. Matem. **28** (1964), 1391–1408 [in Russian].
- [3] N.K. Bari, *Trigonometric series*, Moscow 1961 [in Russian].
- [4] S.V. Bochkarev, *Existence of a basis in the space of functions analytic in a disc and some properties of the Franklin system*, Mat. Sb. **95** (1974) 137, 3–18 [in Russian].
- [5] G. Cantor, *Über einen die Trigonometrischen Reihen betreffenden Lehrsatz*, Crelles J. für Math. **72** (1870), 130–138.
- [6] Z. Ciesielski, *Properties of the orthonormal Franklin system*, Studia Math. **23** (1963), 141–157.
- [7] Z. Ciesielski, *A construction of basis in  $C^{(1)}(I^2)$* , Studia Math. **33** (1969), 289–323.
- [8] Z. Ciesielski, *Constructive function theory and spline systems*, Studia Math. **53** (1975), 278–302.
- [9] Z. Ciesielski, J. Domsta, *Construction of an orthonormal basis in  $C^m(I^d)$  and  $W_p^m(I^d)$* , Studia Math. **41** (1972), 211–224.
- [10] S. Demko, *Inverses of band matrices and local convergence of splines projections*, SIAM J. Numer. Anal. **14** (1977) 4, 616–619.
- [11] Ph. Franklin, *A set of continuous orthogonal functions*, Math. Ann. **100** (1928), 522–529.
- [12] G.G. Gevorkyan, *Ciesielski and Franklin systems*, [in:] Approximation and Probability, Banach Center Publ. 72, Warszawa 2006, 85–92.
- [13] V.Ya. Kozlov, *On complete systems of orthogonal functions*, Mat. Sbornik N.S. **26** (1950) 68, 351–364 [in Russian].
- [14] J. Marcinkiewicz, *Quelques theorems sur les series orthogonales*, Ann. Soc. Polon. Math. **16** (1937), 84–96.
- [15] S. Schonefeld, *Schauder bases in spaces of differential functions*, Bull. Amer. Math. Soc. **75** (1969), 586–590.
- [16] P. Wojtaszczyk, *The Franklin system is an unconditional basis in  $H^1$* , Ark. Mat. **20** (1982), 293–300.

- [17] Z. Wronicz, *On the application of the orthonormal Franklin system to the approximation of analytic functions*, [in:] *Approximation Theory*, Banach Center Publ. 4, PWN, Warszawa 1979, 305–316.
- [18] Z. Wronicz *Approximation by complex splines*, *Zeszyty Nauk. Uniw. Jagiellon. Prace Mat.* **20** (1979), 67–88.

Zygmunt Wronicz  
wronicz@uci.agh.edu.pl

AGH University of Science and Technology  
Faculty of Applied Mathematics  
al. A. Mickiewicza 30, 30-059 Krakow, Poland

*Received: December 19, 2014.*

*Revised: January 9, 2016.*

*Accepted: May 15, 2016.*