

ON FIBONACCI NUMBERS IN EDGE COLOURED TREES

Urszula Bednarz, Dorota Bród, Anetta Szynal-Liana,
Iwona Włoch, and Małgorzata Wołowiec-Musiał

Communicated by Gyula O.H. Katona

Abstract. In this paper we show the applications of the Fibonacci numbers in edge coloured trees. We determine the second smallest number of all $(A, 2B)$ -edge colourings in trees. We characterize the minimum tree achieving this second smallest value.

Keywords: edge colouring, tree, tripod, Fibonacci numbers.

Mathematics Subject Classification: 11B37, 11C20, 15B36, 05C69.

1. INTRODUCTION AND PRELIMINARY RESULTS

The n -th Fibonacci number F_n is defined recursively by the second order linear recurrence relation of the form $F_n = F_{n-1} + F_{n-2}$ for $n \geq 2$ with the initial conditions $F_0 = F_1 = 1$. Research on graph interpretations of the Fibonacci numbers were initiated in 1982 by H. Prodinger and R.F. Tichy in [4]. They showed that the number of all independent sets in n -vertex path is equal to F_n . This simple observation triggered counting problems in graphs related to the Fibonacci numbers. In this paper we consider a special parameter in edge coloured graph which allows to get another graph interpretation of the Fibonacci numbers.

For concepts not defined here see [2, 3]. Let G be a finite, undirected, simple graph with the vertex set $V(G)$ and the edge set $E(G)$. The order (number of vertices) and size (number of edges) of G are denoted by n and m , respectively. By $P(m)$, $T(m)$ and $S(m)$ we denote a path, a tree and a star of size m , respectively. Recall that in a tree a vertex of degree at least 3 is a *branch vertex*, a vertex of degree 1 is a *leaf*. A *tripod* is a tree with exactly three leaves. In other words, every tripod has a unique branch vertex being the initial vertex of three elementary paths. Let $m \geq 3$, $p \geq 1$, $t \geq 1$ be integers. By $T(m, p, t)$ we mean a tripod of size m with paths of lengths p, t and $m - p - t$, starting from the branch vertex. These paths we will denote shortly: *p-path*, *t-path* and *(m - p - t)-path*, respectively.

We begin with a definition of $(A, 2B)$ -edge colouring. Let G be a connected graph and let $\mathcal{C} = \{A, B\}$ be a set of two colours. A graph G is $(A, 2B)$ -edge coloured if for every maximal (with respect to set inclusion) B -monochromatic subgraph H of G there exists a partition of H into edge disjoint paths of length 2. We have no restriction on the colour A , so $(A, 2B)$ -edge colouring exists for an arbitrary graph G . It is worth mentioning that the concept of $(A, 2B)$ -edge colouring is a special case of edge shade colouring of a graph (more details for edge shade colouring can be found in [1]).

Now, assume that \mathcal{F} is a family of all distinct $(A, 2B)$ -edge coloured graphs obtained by colouring of G , i.e. $\mathcal{F} = \{G^{(1)}, G^{(2)}, \dots, G^{(l)}\}$, where $l \geq 1$ and $G^{(p)}$ denotes a graph obtained by $(A, 2B)$ -edge colouring of the graph G for $p = 1, 2, \dots, l$. Let $\theta(G^{(p)})$, where $1 \leq p \leq l$, be the number of all partitions into edge disjoint paths of length 2 of all B -monochromatic subgraphs of $G^{(p)}$. If $G^{(p)}$ is A -monochromatic then we put $\theta(G^{(p)}) = 1$. Let us define a parameter $\sigma_{(A,2B)}(G)$ as follows:

$$\sigma_{(A,2B)}(G) = \sum_{p=1}^l \theta(G^{(p)}).$$

The parameter $\sigma_{(A,2B)}(G)$ was studied for different classes of graphs (see [1]). We recall the main result for trees.

Theorem 1.1 ([1]). *Let $T(m)$ be a tree of size $m, m \geq 1$. Then*

$$F_m \leq \sigma_{(A,2B)}(T(m)) \leq 1 + \sum_{j \geq 1} \binom{m}{2j} \prod_{p=0}^{j-1} [2j - (2p + 1)].$$

Moreover, $\sigma_{(A,2B)}(P(m)) = F_m$ and

$$\sigma_{(A,2B)}(S(m)) = 1 + \sum_{j \geq 1} \binom{m}{2j} \prod_{p=0}^{j-1} [2j - (2p + 1)].$$

From the above theorem follows that the graph $P(m)$ is the extremal tree achieving the minimum value of the parameter $\sigma_{(A,2B)}(T(m))$ and the graph $S(m)$ is the extremal tree achieving the maximum value of the parameter $\sigma_{(A,2B)}(T(m))$. It is natural to ask what are extremal trees achieving the second smallest and the second largest value of this parameter. In this paper we give the lower bound of the parameter $\sigma_{(A,2B)}(T(m))$ with the restriction that $T(m) \not\cong P(m)$ and we describe the extremal graph achieving the second smallest value of the parameter $\sigma_{(A,2B)}(T(m))$.

In the sequel we will use the following notation. If $e \in E(G)$ is a fixed edge that is coloured by the colour A then we will write $c(e) = A$. If e is coloured by the colour B then we will write $c(e) = 2B$ to indicate that there is an edge e' adjacent to e and coloured by the colour B . Moreover, we will write $\sigma_{A(e)}(G)$ (resp. $\sigma_{2B(e)}(G)$) to denote the number of all $(A, 2B)$ -edge colourings of G with $c(e) = A$ (resp. $c(e) = 2B$). The following lemmas give the basic rules for determining the parameter $\sigma_{(A,2B)}(G)$.

Lemma 1.2. *Let $e \in E(G)$ be a fixed edge. Then*

$$\sigma_{(A,2B)}(G) = \sigma_{A(e)}(G) + \sigma_{2B(e)}(G). \tag{1.1}$$

Lemma 1.3. *Let $G = H \cup T(l) \cup \{e\}$ be a connected graph, where H is a connected graph, $T(l)$ is a tree of size $l, l \geq 1$ and H and $T(l)$ are vertex disjoint. Assume that $e = uv$, where $u \in V(H), v \in V(T(l))$ and e is a bridge in G . Then*

$$\sigma_{(A,2B)}(G) \geq \sigma_{(A,2B)}(H \cup P(l) \cup \{e\}).$$

Moreover, the equality holds if $T(l) \cong P(l)$.

Proof. Let $G = H \cup T(l) \cup \{e\}$ and $e = uv \in E(G)$ with $u \in V(H)$ and $v \in V(T(l))$. We distinguish the following cases:

Case 1. $c(e) = A$.

Then $\sigma_{A(e)}(G) = \sigma_{(A,2B)}(H)\sigma_{(A,2B)}(T(l))$.

Case 2. $c(e) = 2B$.

Then there exists an edge, say $e' \in E(G)$, adjacent to the edge e , such that $\{e, e'\}$ belongs to a partition of $2B$ -monochromatic subgraph of G and $c(e') = 2B$. Clearly, either $e' \in E(H)$ or $e' \in E(T(l))$. Hence

$$\sigma_{2B(e)}(G) = \sigma_{2B(e)}(H \cup \{e\})\sigma_{(A,2B)}(T(l)) + \sigma_{(A,2B)}(H)\sigma_{2B(e)}(T(l) \cup \{e\}).$$

Consequently,

$$\begin{aligned} \sigma_{(A,2B)}(G) &= \sigma_{(A,2B)}(H)\sigma_{(A,2B)}(T(l)) + \sigma_{2B(e)}(H \cup \{e\})\sigma_{(A,2B)}(T(l)) \\ &\quad + \sigma_{(A,2B)}(H)\sigma_{2B(e)}(T(l) \cup \{e\}). \end{aligned}$$

By Theorem 1.1, we have $\sigma_{(A,2B)}(T(l)) > \sigma_{(A,2B)}(P(l))$. Hence

$$\begin{aligned} \sigma_{(A,2B)}(G) &\geq \sigma_{(A,2B)}(H)\sigma_{(A,2B)}(P(l)) + \sigma_{2B(e)}(H \cup \{e\})\sigma_{(A,2B)}(P(l)) \\ &\quad + \sigma_{(A,2B)}(H)\sigma_{2B(e)}(P(l+1)) = \sigma_{(A,2B)}(H \cup P(l) \cup \{e\}), \end{aligned}$$

which completes the proof. □

2. EXTREMAL TRIPODS WITH RESPECT TO $\sigma_{(A,2B)}(T(m, p, t))$

As it was mentioned earlier the path $P(m)$ is the extremal graph achieving the minimum value of the parameter $\sigma_{(A,2B)}(T(m))$ in the class of trees of size m . Looking for the second smallest value of the parameter $\sigma_{(A,2B)}$ in trees of size m let us consider the class of trees $T(m)$ such that $T(m) \not\cong P(m)$. This means that there exists at least one branch vertex in $T(m)$. We start with the class of tripods because results obtained for this class will be crucial for the main result. Assume that $\mathcal{T} = \{T(m, p, t); m \geq 3, p \geq 1, t \geq 1\}$ is the family of tripods.

Theorem 2.1. *Let $m \geq 3, p \geq 1, t \geq 1$ be integers. Then for an arbitrary tripod $T(m, p, t) \in \mathcal{T}$ holds*

$$\sigma_{(A,2B)}(T(m, p, t)) = F_{p+t}F_{m-t-p} + F_{m-t-p-1}(F_{p-1}F_t + F_pF_{t-1}). \tag{2.1}$$

Proof. Let $T(m, p, t) \in \mathcal{T}$. If $m = 3$ then $p = t = 1$ and by the simple observation we have

$$\sigma_{(A,2B)}(T(3, 1, 1)) = 4 = F_2F_1 + F_0(F_0F_1 + F_1F_0).$$

Let $m \geq 4$ and let $x \in V(T(m, p, t))$ be the unique branch vertex of a tripod. Clearly $p \geq 2$ or $t \geq 2$ or $m - p - t \geq 2$. Without loss of generality suppose that $m - p - t \geq 2$. Let $e \in E(T(m, p, t))$ be the edge incident with the vertex x and e belongs to $(m - p - t)$ -path of $T(m, p, t)$.

Consider the following cases:

Case 1. $c(e) = A$.

Then edges adjacent to e are coloured by A or $2B$. This means that we have exactly $F_{p+t}F_{m-t-p-1}$ distinct $(A, 2B)$ -edge colourings with $c(e) = A$.

Case 2. $c(e) = 2B$.

Then there exists an edge, say $e' \in E(T(m, p, t))$, adjacent to e and coloured by $2B$ and $\{e, e'\}$ belongs to a partition of $2B$ -monochromatic subgraph of $T(m, p, t)$. Clearly e' belongs to either p -path or t -path or $(m - t - p)$ -path. Considering all these possibilities we obtain $F_{p-1}F_tF_{m-t-p-1} + F_pF_{t-1}F_{m-t-p-1} + F_{p+t}F_{m-t-p-2}$ $(A, 2B)$ -edge colourings with $c(e) = 2B$.

Finally, by the above, by Lemma 1.2 and by simple calculations we obtain

$$\begin{aligned} \sigma_{(A,2B)}(T(m, p, t)) &= F_{p+t}F_{m-t-p-1} + F_{p-1}F_tF_{m-t-p-1} \\ &\quad + F_{t-1}F_pF_{m-t-p-1} + F_{p+t}F_{m-t-p-2} \\ &= F_{p+t}F_{m-t-p} + F_{m-t-p-1}(F_{p-1}F_t + F_pF_{t-1}), \end{aligned}$$

which ends the proof. □

Corollary 2.2. *Let $m \geq 3, t \geq 1$ be integers. Then*

- a) $\sigma_{(A,2B)}(T(m, 1, t)) = F_{t+1}F_{m-t}$,
- b) $\sigma_{(A,2B)}(T(m, 1, 1)) = 2F_{m-1}$.

Proof. a) Applying (2.1) for $p = 1$ and using the definition of Fibonacci numbers we have

$$\begin{aligned} \sigma_{(A,2B)}(T(m, 1, t)) &= F_{t+1}F_{m-t-1} + F_{m-t-2}(F_t + F_{t-1}) \\ &= F_{t+1}(F_{m-t-1} + F_{m-t-2}) = F_{t+1}F_{m-t}. \end{aligned}$$

b) Analogously by Theorem 2.1 for $p = t = 1$ we obtain

$$\sigma_{(A,2B)}(T(m, 1, 1)) = F_2F_{m-2} + 2F_{m-3} = 2F_{m-1}. \quad \square$$

Using the above results we can give the maximum value of the parameter $\sigma_{(A,2B)}(T(m, p, t))$. We will need the following well-known identities:

$$F_{m-1} = F_{p+t}F_{m-p-t-1} + F_{p+t-1}F_{m-p-t-2}, \tag{2.2}$$

$$F_{m+n} = F_mF_n + F_{m-1}F_{n-1}. \tag{2.3}$$

Theorem 2.3. *Let $m \geq 4, p \geq 1, t \geq 1$ be integers. Then for an arbitrary tripod $T(m, p, t) \in \mathcal{T}$ holds*

$$\sigma_{(A,2B)}(T(m, p, t)) \leq 2F_{m-1}.$$

Moreover, $\sigma_{(A,2B)}(T(m, p, t)) = 2F_{m-1}$ iff $T(m, p, t) \cong T(m, 1, 1)$.

Proof. It suffices to prove that

$$F_{p+t}F_{m-t-p} + F_{m-t-p-1}(F_{p-1}F_t + F_pF_{t-1}) - 2F_{m-1} \leq 0.$$

Applying (2.2) and the definition of Fibonacci numbers we have

$$\begin{aligned} &F_{p+t}F_{m-t-p} + F_{m-t-p-1}(F_{p-1}F_t + F_pF_{t-1}) - 2F_{p+t}F_{m-t-p-1} - 2F_{t+p-1}F_{m-t-p-2} \\ &= F_{p+t}F_{m-t-p-1} + F_{p+t}F_{m-t-p-2} + F_{m-t-p-1}(F_{p-1}F_t + F_pF_{t-1}) \\ &\quad - 2F_{p+t}F_{m-t-p-1} - 2F_{t+p-1}F_{m-t-p-2} \\ &= F_{m-t-p-1}(F_{p+t} + F_{p-1}F_t + F_pF_{t-1} - 2F_{p+t}) + F_{m-t-p-2}(F_{p+t} - 2F_{p+t-1}). \end{aligned}$$

By (2.3), we obtain

$$\begin{aligned} &F_{m-t-p-1}(F_{p-1}F_t + F_pF_{t-1} - F_pF_t - F_{p-1}F_{t-1}) \\ &\quad + F_{m-t-p-2}(F_{p+t-1} + F_{p+t-2} - 2F_{p+t-1}) \\ &= F_{m-t-p-1}(F_t(F_{p-1} - F_p) - F_{t-1}(F_{p-1} - F_p)) + F_{m-t-p-2}(F_{p+t-2} - F_{p+t-1}) \\ &= F_{m-t-p-1}(F_{p-1} - F_p)(F_t - F_{t-1}) + F_{m-t-p-2}(F_{p+t-2} - F_{p+t-1}) \leq 0 \end{aligned}$$

by $F_{p-1} \leq F_p$ and $F_{p+t-2} \leq F_{p+t-1}$.

Moreover, the equality holds if and only if $F_{p+t-2} = F_{p+t-1}$ and $F_{p-1} = F_p$ or $F_{p+t-2} = F_{p+t-1}$ and $F_t = F_{t-1}$. Clearly, $F_{p+t-2} = F_{p+t-1}$ only for $p+t-2 = 0$ and $p+t-1 = 1$. Hence $p+t = 2$, so $p = t = 1$. Consequently, $F_{p-1} = F_p$ and $F_t = F_{t-1}$, which immediately gives that the extremal tripod achieving the maximum value of the parameter $\sigma_{(A,2B)}(T(m, p, t))$ is only the tripod $T(m, 1, 1)$. \square

Now we give the recurrence rule for determining the parameter $\sigma_{(A,2B)}(T(m, p, t))$.

Theorem 2.4. *Let $m \geq 3, p \geq 1, t \geq 1$ be integers. Then for an arbitrary tripod $T(m, p, t) \in \mathcal{T}$ and $m - p - t \geq 3$ holds*

$$\sigma_{(A,2B)}(T(m, p, t)) = \sigma_{(A,2B)}(T(m - 1, p, t)) + \sigma_{(A,2B)}(T(m - 2, p, t)) \tag{2.4}$$

with initial conditions

$$\sigma_{(A,2B)}(T(p+t+1, p, t)) = F_{p+1}F_{t+1} \text{ and } \sigma_{(A,2B)}(T(p+t+2, p, t)) = F_{p+1}F_{t+1} + F_{p+t}.$$

Proof. Let $T(m, p, t) \in \mathcal{T}$. If $m = p + t + 1$ then $(m - p - t)$ -path has length 1. Let $e \in E(T(p + t + 1, p, t))$ be the unique edge of the $(m - p - t)$ -path. We distinguish the following cases.

Case 1. $c(e) = A$.

Then edges adjacent to e are coloured by A or $2B$. This means that $T(p+t+1, p, t) \setminus e \cong P(p+t)$ and, by Theorem 1.1, $\sigma_{A(e)}(T(p + t + 1, p, t)) = F_{p+t}$.

Case 2. $c(e) = 2B$.

Then there exists an edge $e' \in E(T(p+t+1, p, t) \setminus e)$ adjacent to e such that $c(e') = 2B$. Clearly e' belongs to either p -path or t -path, by $m = p + t + 1$. Considering these two possibilities we obtain that $\sigma_{2B(e)}(T(p + t + 1, p, t)) = F_{p-1}F_t + F_pF_{t-1}$.

Consequently, by (2.3), we have

$$\begin{aligned} \sigma_{(A,2B)}(T(p + t + 1, p, t)) &= F_{p+t} + F_{p-1}F_t + F_pF_{t-1} \\ &= F_pF_t + F_{p-1}F_{t-1} + F_{p-1}F_t + F_pF_{t-1} \\ &= F_t(F_p + F_{p-1}) + F_{t-1}(F_p + F_{p-1}) \\ &= F_{t+1}F_{p+1}. \end{aligned}$$

If $m = p + t + 2$ then, with respect to an edge $e \in E(T(m + p + t + 2, p, t))$ belonging to the $(m - p - t)$ -path and incident with a leaf, we obtain

$$\begin{aligned} \sigma_{(A,2B)}(T(p + t + 2, p, t)) &= F_{p+t} + \sigma_{(A,2B)}(T(p + t + 1, p, t)) \\ &= F_{p+t} + F_{t+1}F_{p+1}. \end{aligned}$$

Assume now $m - p - t \geq 3$. Let $e \in E(T(m, p, t))$ be an edge of the $(m - p - t)$ -path incident with a leaf. By analogy we obtain

$$\sigma_{(A,2B)}(T(m, p, t)) = \sigma_{(A,2B)}(T(m - 1, p, t)) + \sigma_{(A,2B)}(T(m - 2, p, t)),$$

which completes the proof. □

Solving the recurrence relation (2.4) we obtain the Binet formulas for the parameters $\sigma_{(A,2B)}(T(m, 1, 1))$ and $\sigma_{(A,2B)}(T(m, 2, 2))$.

$$\begin{aligned} \sigma_{(A,2B)}(T(m, 1, 1)) &= \frac{2\sqrt{5}}{5} \left[\left(\frac{1 + \sqrt{5}}{2} \right)^m - \left(\frac{1 - \sqrt{5}}{2} \right)^m \right] \quad \text{for } m \geq 3, \\ \sigma_{(A,2B)}(T(m, 2, 2)) &= \left(\frac{4\sqrt{5}}{5} - 1 \right) \left(\frac{1 + \sqrt{5}}{2} \right)^m \\ &\quad - \left(\frac{4\sqrt{5}}{5} + 1 \right) \left(\frac{1 - \sqrt{5}}{2} \right)^m \quad \text{for } m \geq 5. \end{aligned}$$

We shall show that $T(m, 2, 2)$ is the extremal tripod achieving the minimum value of the parameter $\sigma_{(A,2B)}(T(m, p, t))$ in the class \mathcal{T} . Consider non-isomorphic tripods of size $m = 5, 6$ (Figs 1 and 2).




Fig 1. All non-isomorphic tripods of size 5




Fig 2. All non-isomorphic tripods of size 6

For these tripods values of the parameter $\sigma_{(A,2B)}(T(m, p, t))$ are given in Table 1.

Table 1.

$T(m, p, t)$	$T(5, 2, 2)$	$T(5, 3, 1)$	$T(6, 2, 2)$	$T(6, 3, 2)$	$T(6, 4, 1)$
$\sigma_{(A,2B)}(T(m, p, t))$	9	10	14	15	16

Theorem 2.5. *Let $m \geq 5, p \geq 1, t \geq 1$ be integers. Then for an arbitrary tripod $T(m, p, t) \in \mathcal{T}$ holds*

$$\sigma_{(A,2B)}(T(m, p, t)) \geq F_{m-1} + 2F_{m-3}.$$

Moreover, the equality holds if $T(m, p, t) \cong T(m, 2, 2)$.

Proof (by induction on m). If $m = 5, 6$ then the result follows immediately from Table 1, Figures 1 and 2 and the definition of Fibonacci numbers.

Let $m \geq 7$. Assume that for all $n < m$ holds $\sigma_{(A,2B)}(T(n, p, t)) \geq F_{n-1} + 2F_{n-3}$. We shall show that the theorem is true for m . Since $m \geq 7$, we have that at least one path of tripod $T(m, p, t)$ has length at least 3. Without loss of generality we can assume that $m - p - t \geq 3$. Using Theorem 2.4 and the induction hypothesis we have

$$\begin{aligned} \sigma_{(A,2B)}(T(m, p, t)) &= \sigma_{(A,2B)}(T(m - 1, p, t)) + \sigma_{(A,2B)}(T(m - 2, p, t)) \\ &\geq F_{m-2} + 2F_{m-4} + F_{m-3} + 2F_{m-5} = F_{m-1} + 2F_{m-3} \end{aligned}$$

and the theorem follows.

Now we shall show that $\sigma_{(A,2B)}(T(m, 2, 2)) = F_{m-1} + 2F_{m-3}$. By Theorem 2.1 and by the definition of Fibonacci numbers, we obtain

$$\begin{aligned} \sigma_{(A,2B)}(T(m, 2, 2)) &= 5F_{m-4} + 4F_{m-5} = 4F_{m-3} + F_{m-4} \\ &= F_{m-2} + 3F_{m-3} = F_{m-1} + 2F_{m-3}, \end{aligned}$$

which completes the proof. □

3. MAIN RESULTS

In this section we determine the second smallest value of the parameter $\sigma_{(A,2B)}(T(m))$. We show that the tripod $T(m, 2, 2)$ realizes this second minimum value of $\sigma_{(A,2B)}(T(m))$.

Let $r \geq 1, \Delta \geq 3$ be integers. For $m \geq 3$ by a tree $S_r(m, \Delta)$ we mean a graph with a unique branch vertex obtained from the star with maximum degree Δ by inserting new vertices of degree 2 into some edges of the star such that in the resulting tree $S_r(m, \Delta)$ the longest path starting from the branch vertex has length r . In particular, $S_1(m, \Delta)$ is isomorphic to a star $S(m)$ and $S_r(m, 3)$ is isomorphic to a tripod $T(m, r, t)$, for some $t \geq 1$.

Theorem 3.1. *Let $m \geq 4, \Delta \geq 3$ be integers. Then*

$$\sigma_{(A,2B)}(S_2(m, \Delta)) = \begin{cases} \sigma_{(A,2B)}(S(m-1)) + \sigma_{(A,2B)}(S(m-2)), \\ \text{if } S_2(m, \Delta) \text{ has the unique 2-path,} \\ \sigma_{(A,2B)}(S_2(m-1, \Delta)) + \sigma_{(A,2B)}(S_2(m-2, \Delta-1)), \\ \text{otherwise.} \end{cases}$$

Proof. Let $m \geq 4$ and $\Delta \geq 3$ be integers. Consider two cases.

Case 1. There exists a unique 2-path in the tree $S_2(m, \Delta)$.

Let $e \in E(S_2(m, \Delta))$ be an edge which belongs to the 2-path and e is incident with a leaf. We have two possibilities.

Case 1.1. $c(e) = A$.

Then $\sigma_{A(e)}(S_2(m, \Delta)) = \sigma_{(A,2B)}(S(m-1))$.

Case 1.2. $c(e) = 2B$.

Then $\sigma_{2B(e)}(S_2(m, \Delta)) = \sigma_{(A,2B)}(S(m-2))$. Hence

$$\sigma_{(A,2B)}(S_2(m, \Delta)) = \sigma_{(A,2B)}(S(m-1)) + \sigma_{(A,2B)}(S(m-2)).$$

Case 2. There exist at least two 2-paths in the tree $S_2(m, \Delta)$.

Let $e \in E(S_2(m, \Delta))$ be an edge which belongs to any 2-path and e is incident with a leaf. We have two possibilities.

Case 2.1. $c(e) = A$.

Then $\sigma_{A(e)}(S_2(m, \Delta)) = \sigma_{(A,2B)}(S_2(m-1, \Delta))$.

Case 2.2. $c(e) = 2B$.

Then $\sigma_{2B(e)}(S_2(m, \Delta)) = \sigma_{(A,2B)}(S_2(m-2, \Delta-1))$. Hence

$$\sigma_{(A,2B)}(S_2(m, \Delta)) = \sigma_{(A,2B)}(S_2(m-1, \Delta)) + \sigma_{(A,2B)}(S_2(m-2, \Delta-1)),$$

which completes the proof. □

Theorem 3.2. *Let $m \geq 4, \Delta \geq 4, r \geq 1$ be integers. Then for an arbitrary tripod $T(m, p, t) \in \mathcal{T}$ holds*

$$\sigma_{(A,2B)}(S_r(m, \Delta)) > \sigma_{(A,2B)}(T(m, p, t)). \tag{3.1}$$

Proof. Let m, Δ, r be as in the statement of the theorem. We consider the following cases:

Case 1. There exists a unique r -path in the tree $S_r(m, \Delta)$.

Clearly $r \geq 2$ and $m \geq 5$. We use induction on m and r . If $m = 5$ then $r = 2$ and the result is obvious. Let $m \geq 6$ and $r \geq 2$ and assume that the inequality (3.1) holds for all $n < m$ and $k < r$. Let $e \in E(S_r(m, \Delta))$ belongs to the r -path and e is incident with a leaf of $S_r(m, \Delta)$. We need to consider two cases.

Case 1.1. $c(e) = A$.

Then $\sigma_{A(e)}(S_r(m, \Delta)) = \sigma_{(A,2B)}(S_{r-1}(m-1, \Delta))$.

Case 1.2. $c(e) = 2B$.

If $r = 2$ then the unique 2-path P is coloured by $2B$ and the graph $S_2(m, \Delta) \setminus P$ is isomorphic to a star $S(m-2)$. Hence $\sigma_{2B(e)}(S_2(m, \Delta)) = \sigma_{(A,2B)}(S(m-2))$.

If $r \geq 3$ then the graph $S_r(m, \Delta) \setminus P$ is isomorphic to a graph $S_{k < r}(m-2, \Delta)$.

If $r = 2$ then by the above and using the induction hypothesis we obtain

$$\begin{aligned} \sigma_{(A,2B)}(S_2(m, \Delta)) &= \sigma_{(A,2B)}(S_1(m-1, \Delta)) + \sigma_{(A,2B)}(S(m-2)) \\ &> \sigma_{(A,2B)}(T(m-1, p, t)) + \sigma_{(A,2B)}(S(m-2)). \end{aligned}$$

We shall show that for $T(m, p, t) \in \mathcal{T}$

$$\sigma_{(A,2B)}(T(m-1, p, t)) + \sigma_{(A,2B)}(S(m-2)) > \sigma_{(A,2B)}(T(m, p, t)).$$

It suffices to prove the following inequality

$$\sigma_{(A,2B)}(T(m-1, p, t)) + \sigma_{(A,2B)}(S(m-2)) - \sigma_{(A,2B)}(T(m, p, t)) > 0.$$

By Theorem 2.4, we have

$$\begin{aligned} &\sigma_{(A,2B)}(T(m-1, p, t)) + \sigma_{(A,2B)}(S(m-2)) - \sigma_{(A,2B)}(T(m-1, p, t)) \\ &- \sigma_{(A,2B)}(T(m-2, p, t)) > 0 \end{aligned}$$

because the star maximizes this parameter in trees.

If $r \geq 3$ then using the induction hypothesis we obtain

$$\begin{aligned} \sigma_{(A,2B)}(S_r(m, \Delta)) &= \sigma_{(A,2B)}(S_{r-1}(m-1, \Delta)) + \sigma_{(A,2B)}(S_{k < r}(m-2, \Delta)) \\ &> \sigma_{(A,2B)}(T(m-1, p, t)) + \sigma_{(A,2B)}(T(m-2, p, t)) \\ &= \sigma_{(A,2B)}(T(m, p, t)), \end{aligned}$$

which completes the proof of this case.

Case 2. There exist at least two r -paths in $S_r(m, \Delta)$.

For $r = 1$ the result is obvious since $S_1(m, \Delta)$ is isomorphic to the star. Let $r \geq 2$. Then $m \geq 6$. We now proceed by induction on m . If $m = 6$ then $r = 2$ and the result is obvious.

Let $m \geq 7$ and assume that for all $n < m$ the inequality holds. We distinguish two possibilities.

Case 2.1. $c(e) = A$.

Then $\sigma_{A(e)}(S_r(m, \Delta)) = \sigma_{(A,2B)}(S_r(m - 1, \Delta))$.

Case 2.2. $c(e) = 2B$.

If $r = 2$ then $S_2(m, \Delta) \setminus P$ is isomorphic to $S_2(m - 2, \Delta - 1)$. If $r \geq 3$ then $S_2(m, \Delta) \setminus P$ is isomorphic to $S_2(m - 2, \Delta)$. Let $r = 2$. Then from these possibilities and by the induction hypothesis we obtain

$$\begin{aligned} \sigma_{(A,2B)}(S_2(m, \Delta)) &= \sigma_{(A,2B)}(S_2(m - 1, \Delta)) + \sigma_{(A,2B)}(S_2(m - 2, \Delta - 1)) \\ &> \sigma_{(A,2B)}(T(m - 1, p, t)) + \sigma_{(A,2B)}(S_2(m - 2, \Delta - 1)). \end{aligned}$$

We shall show that

$$\sigma_{(A,2B)}(T(m - 1, p, t)) + \sigma_{(A,2B)}(S_2(m - 2, \Delta - 1)) > \sigma_{(A,2B)}(T(m, p, t))$$

for all $T(m, p, t) \in \mathcal{T}$. It suffices to prove that

$$\sigma_{(A,2B)}(T(m - 1, p, t)) + \sigma_{(A,2B)}(S_2(m - 2, \Delta - 1)) - \sigma_{(A,2B)}(T(m, p, t)) > 0.$$

Suppose that there exist l ($l \geq 2$) 2-paths in $S_2(m, \Delta)$. Then by (2.4) and applying the induction hypothesis in l steps we obtain

$$\begin{aligned} &\sigma_{(A,2B)}(T(m - 1, p, t)) + \sigma_{(A,2B)}(S_2(m - 2, \Delta - 1)) \\ &\quad - \sigma_{(A,2B)}(T(m - 1, p, t)) - \sigma_{(A,2B)}(T(m - 2, p, t)) \\ &= \sigma_{(A,2B)}(S_2(m - 2, \Delta - 1)) - \sigma_{(A,2B)}(T(m - 2, p, t)) > 0 \end{aligned}$$

in the first step. Consequently in the l th step

$$\sigma_{(A,2B)}(S_2(m - l - 1, \Delta - l)) - \sigma_{(A,2B)}(T(m - l - 1, p, t)) > 0$$

since $S_2(m - l - 1, \Delta - l)$ is isomorphic to the star $S(m - l - 1)$ and the result immediately follows.

Let $r \geq 3$. Then from Cases 2.1 and 2.2 and using the induction hypothesis we obtain

$$\begin{aligned} \sigma_{(A,2B)}(S_r(m, \Delta)) &= \sigma_{(A,2B)}(S_r(m - 1, \Delta)) + \sigma_{(A,2B)}(S_r(m - 2, \Delta)) \\ &> \sigma_{(A,2B)}(T(m - 1, p, t)) + \sigma_{(A,2B)}(T(m - 2, p, t)) \\ &= \sigma_{(A,2B)}(T(m, p, t)), \end{aligned}$$

and the proof is complete. □

Corollary 3.3. *Let $m \geq 4$, $\Delta \geq 4$, $r \geq 1$ be integers. Then*

$$\sigma_{(A,2B)}(S_r(m, \Delta)) > F_{m-1} + 2F_{m-3}.$$

Proof. By Theorems 3.2 and 2.5, we immediately obtain

$$\sigma_{(A,2B)}(S_r(m, \Delta)) > \sigma_{(A,2B)}(T(m, p, t)) \geq F_{m-1} + 2F_{m-3}. \quad \square$$

Theorem 3.4. *Let $T(m) \not\cong P(m)$ be a tree of the size m . Then*

$$\sigma_{(A,2B)}(T(m)) \geq F_{m-1} + 2F_{m-3}. \quad (3.2)$$

Moreover, $\sigma_{(A,2B)}(T(m)) = F_{m-1} + 2F_{m-3}$ if $T(m) \cong T(m, 2, 2)$.

Proof. Assume that $T(m)$ is a tree of size m non-isomorphic to the path $P(m)$. Since $T(m) \not\cong P(m)$, there exists in $T(m)$ at least one branch vertex, say x . If $T(m)$ has a unique branch vertex then the result follows by Theorem 3.2. Suppose that $T(m)$ has at least two branch vertices and let $u, v \in V(T(m))$ be such vertices. Let $e \in E(T(m))$ be an edge belonging to the path $u - v$ in $T(m)$. Then $T(m) = T_1(m_1) \cup T_2(m_2) \cup \{e\}$, where $T_i(m_i)$ for $i = 1, 2$ are trees of the size m_i , $m_i \geq 2$. Applying Lemma 1.3 we obtain

$$\begin{aligned} \sigma_{(A,2B)}(T(m)) &= \sigma_{(A,2B)}(T_1(m_1) \cup T_2(m_2) \cup \{e\}) \\ &\geq \sigma_{(A,2B)}(T_1(m_1) \cup P(m_2) \cup \{e\}). \end{aligned}$$

If $T_1(m_1) \cup P(m_2) \cup \{e\}$ is $S_r(m, \Delta)$, then by Theorem 3.2 the result follows. Otherwise, it has at least two branch vertices and we repeat the above procedure until we get a tree T^* of the same size m . By Theorem 3.2 we have $\sigma_{(A,2B)}(T^*) > \sigma_{(A,2B)}(T(m, p, t))$. In the class \mathcal{T} the minimum tripod $T(m, 2, 2)$ has the parameter $\sigma_{(A,2B)}(T(m, 2, 2)) = F_{m-1} + 2F_{m-3}$, which completes the proof. \square

REFERENCES

- [1] U. Bednarz, I. Włoch, M. Wołowicz-Musiał, *Total graph interpretation of numbers of the Fibonacci type*, J. Appl. Math. **2015** (2015), Article ID 837917.
- [2] C. Berge, *Principle of Combinatorics*, Academic Press, New York, London, 1971.
- [3] R. Diestel, *Graph Theory*, Springer-Verlag, Heidelberg, New York, 2005.
- [4] H. Prodinger, R.F. Tichy, *Fibonacci numbers of graphs*, Fibonacci Quart. **20** (1982) 1, 16–21.

Urszula Bednarz
ubednarz@prz.edu.pl

Rzeszow University of Technology
Faculty of Mathematics and Applied Physics
al. Powstańców Warszawy 12, 35-359 Rzeszów, Poland

Dorota Bród
dorotab@prz.edu.pl

Rzeszow University of Technology
Faculty of Mathematics and Applied Physics
al. Powstańców Warszawy 12, 35-359 Rzeszów, Poland

Anetta Szynal-Liana
aszynal@prz.edu.pl

Rzeszow University of Technology
Faculty of Mathematics and Applied Physics
al. Powstańców Warszawy 12, 35-359 Rzeszów, Poland

Iwona Włoch
iwloch@prz.edu.pl

Rzeszow University of Technology
Faculty of Mathematics and Applied Physics
al. Powstańców Warszawy 12, 35-359 Rzeszów, Poland

Małgorzata Wołowiec-Musiał
wołowiec@prz.edu.pl

Rzeszow University of Technology
Faculty of Mathematics and Applied Physics
al. Powstańców Warszawy 12, 35-359 Rzeszów, Poland

Received: April 12, 2016.

Revised: September 16, 2016.

Accepted: September 16, 2016.