

Wacław Pielichowski

**SOME REMARKS ON THE OPTIMIZATION
OF EIGENVALUE PROBLEMS
INVOLVING THE p -LAPLACIAN**

Abstract. Given a bounded domain $\Omega \subset \mathbb{R}^n$, numbers $p > 1$, $\alpha \geq 0$ and $A \in [0, |\Omega|]$, consider the optimization problem: find a subset $D \subset \Omega$, of measure A , for which the first eigenvalue of the operator $u \mapsto -\operatorname{div}(|\nabla u|^{p-2}\nabla u) + \alpha\chi_D|u|^{p-2}u$ with the Dirichlet boundary condition is as small as possible. We show that the optimal configuration D is connected with the corresponding positive eigenfunction u in such a way that there exists a number $t \geq 0$ for which $D = \{u \leq t\}$. We also give a new proof of symmetry of optimal solutions in the case when Ω is Steiner symmetric and $p = 2$.

Keywords: p -Laplacian, the first eigenvalue, Steiner symmetry.

Mathematics Subject Classification: 35P30, 35J65, 35J70.

1. INTRODUCTION

In this paper we obtain some results closely connected with those of [9], concerning the optimal pairs of an eigenvalue problem involving the p -Laplacian. The paper [9] is available online at www.im.uj.edu.pl/actamath. For the reader's convenience we shall recall the basic notation and terminology of [9], which in turn follow those of [1]. The paper [1] has originated research in the optimization of eigenvalues for the linear case of $p = 2$.

Let Ω be a bounded domain (i.e. open and connected set) in the space \mathbb{R}^n ($n \geq 1$) with the closure $\bar{\Omega}$ and boundary $\partial\Omega$. We denote by $|\Omega|$ the Lebesgue measure of Ω . Given numbers $p > 1$, $\alpha \geq 0$ and a measurable subset D of Ω , we shall be concerned with the eigenvalue problem of the form

$$\begin{cases} -\Delta_p(u) + \alpha\chi_D\varphi_p(u) = \lambda\varphi_p(u) & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases} \quad (1)$$

where Δ_p is the p -Laplacian, χ_D is the characteristic function of D , while φ_p is a function defined by

$$\varphi_p(u) := \begin{cases} |u|^{p-2}u, & \text{if } u \neq 0, \\ 0, & \text{if } u = 0. \end{cases}$$

The p -Laplacian is a nonlinear differential operator of the form

$$\Delta_p(u) = \operatorname{div}(|\nabla u|^{p-2}\nabla u) = \operatorname{div}(\varphi_p(\nabla u)),$$

which coincides with the Laplacian Δ for $p = 2$.

In this paper we deal with real function spaces only. In particular, we use standard Sobolev spaces $W^{1,p}(\Omega)$ and $W_0^{1,p}(\Omega)$, with $1 < p < \infty$. It is customary to use solutions of (1) in a weak sense. Any nontrivial function $u: \Omega \rightarrow \mathbb{R}$ is said to be an eigenfunction of problem (1) if and only if $u \in W_0^{1,p}(\Omega)$ and

$$\int_{\Omega} \varphi_p(\nabla u)\nabla v + \alpha \int_{\Omega} \chi_D \varphi_p(u)v = \lambda \int_{\Omega} \varphi_p(u)v, \quad \text{for any } v \in W_0^{1,p}(\Omega).$$

Let $\lambda(\alpha, D)$ stand for the lowest eigenvalue λ of problem (1). It is known that $\lambda(\alpha, D)$ is positive and its eigenfunction is unique up to a scalar multiple (see, e.g., [3, 10] and the references therein). Let us fix $A \in [0, |\Omega|]$ and define

$$\Lambda(\alpha, A) := \inf \{ \lambda(\alpha, D) : D \subset \Omega, |D| = A \}. \quad (2)$$

Any minimizer in (2) is called an *optimal configuration*. If u is an eigenfunction of problem (1) with $\lambda = \Lambda(\alpha, A)$ and with an optimal configuration D , then (u, D) is said to be an *optimal pair* (or *optimal solution*).

If u is an eigenfunction corresponding to the first eigenvalue of problem (1), then u does not change sign in Ω (see, e.g., [3, 10] and the references therein). From now on it will be chosen positive in Ω .

The above results were discussed in detail in our paper [9]. We shall also use the following lemmas:

Lemma 1. *Let $u \in W_{loc}^{1,1}(\Omega)$ and $t \in \mathbb{R}$. Then $\nabla u(x) = 0$ for almost every $x \in \{u=t\}$.*

In this connection refer to [5], Lemma 7.7, or [8], Theorem 6.19. We use the notation $\{u=t\} := \{x \in \Omega : u(x) = t\}$.

Lemma 2. *Assume that $u \in W_{loc}^{1,p}(\Omega)$ is a weak solution of the equation*

$$-\Delta_p(u) = f \quad \text{in } \Omega$$

with $p > 1$, $f \in L^q(\Omega)$, $q > \frac{n}{p}$, $q \geq 2$. Let

$$Z := \{x \in \Omega : \nabla u(x) = 0\}.$$

Then $|\nabla u|^{p-1} \in W_{loc}^{1,2}(\Omega)$ and $f(x) = 0$ for almost every $x \in Z$.

This result comes from [7] and is quoted in [2].

2. OPTIMAL PAIRS

We recall that Ω is any bounded domain in \mathbb{R}^n and $p \in (1, \infty)$. It is worth noting that we need no additional assumptions concerning the regularity of the boundary $\partial\Omega$.

Theorem 3. *For any $\alpha \geq 0$ and $A \in [0, |\Omega|]$ there exists an optimal pair.*

A proof of this result can be found in our paper [9].

Theorem 4. *Every optimal pair (u, D) has the following properties:*

- (a) $u \in W_0^{1,p}(\Omega) \cap L^\infty(\Omega)$ and ∇u is locally Hölder continuous, i.e., for every compact $K \subset \Omega$ there exists $\beta \in (0, 1)$ such that $\nabla u \in C^{0,\beta}(K)$,
- (b) there is a number $t \geq 0$ such that (up to a set of measure zero)

$$D = \{u \leq t\}. \tag{3}$$

As usual, we write $\{u < t\}$ instead of $\{x \in \Omega: u(x) < t\}$ and similarly we put $\{u \leq t\} := \{x \in \Omega: u(x) \leq t\}$.

Proof. The regularity properties of eigenfunctions, stated in assertion (a), are rather well known. In this connection see [9] and the references therein. Equality (3) in the case of $p = 2$ was stated in [1]. A lack of higher regularity of eigenfunctions is a source of difficulty in obtaining more general results.

We now claim that (3) holds for arbitrary $p > 1$. For $p \neq 2$ this is a new result. Let (u, D) be an optimal solution, corresponding to the optimal eigenvalue $\lambda_1 = \Lambda(\alpha, A)$ with $A > 0$ (the case of $A = 0$ is obvious). Note that according to [9], Theorem 1, there exists a number $t > 0$ such that

$$\{u < t\} \subset D \subset \{u \leq t\}. \tag{4}$$

In fact

$$t = \sup \{s: |\{u < s\}| \leq A\}.$$

In view of (4), it is sufficient to show that

$$|D^c \cap \{u = t\}| = 0,$$

where $D^c := \Omega \setminus D$. To begin with, let us introduce the critical set

$$Z := \{x \in \Omega: \nabla u(x) = 0\}.$$

According to Lemma 1, $\{u = t\} \subset Z$ and hence we see that $D^c \cap \{u = t\} \subset Z$. By Lemma 2,

$$-\Delta_p(u) = 0 \quad \text{in } Z, \tag{5}$$

(i.e. this equality holds almost everywhere in Z). On the other hand,

$$-\Delta_p(u) = (\lambda_1 - \alpha\chi_D)\varphi_p(u) = \lambda_1\varphi_p(u) \quad \text{in } D^c. \tag{6}$$

It now follows from (5) and (6) that

$$\lambda_1 \varphi_p(u) = 0 \quad \text{in } D^c \cap \{u = t\}.$$

Note that $\lambda_1 \neq 0$ and $\varphi_p(u) = t^{p-1} \neq 0$ in $\{u = t\}$. Thus

$$\lambda_1 \varphi_p(u) \neq 0 \quad \text{in } D^c \cap \{u = t\}.$$

This is only possible when $|D^c \cap \{u = t\}| = 0$, as desired. \square

Remark 5. *In a similar way, we can conclude that all level sets $\{u = s\}$ (with $s > 0$) have measure zero, provided that $\alpha \neq \Lambda(\alpha, A)$.*

Remark 6. *According to (3), our optimization of eigenvalue problem is equivalent to finding the smallest eigenvalue and an associated eigenfunction of the problem*

$$\begin{cases} -\Delta_p(u) + \alpha \chi_{\{u \leq t\}} \varphi_p(u) = \lambda \varphi_p(u) & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \\ |\{u \leq t\}| = A \end{cases}$$

with free variables u and t .

3. STEINER SYMMETRY

In this section we consider the linear case with $p = 2$ and we give another proof of a known result concerning the symmetry of optimal solutions (see [1], Theorem 4). Some ideas of [6] are adopted here.

From now on we shall assume that Ω satisfies the exterior cone condition at each point $x \in \partial\Omega$, which means that there exists a finite right circular cone $V = V_x$ with vertex x such that $\overline{\Omega} \cap V_x = \{x\}$.

Let us recall that a domain G of \mathbb{R}^n is Steiner symmetric with respect to a hyperplane P iff for any point $x = (x_1, \dots, x_n) \in G$ the segment connecting x and the point x^* reflected with respect to P is contained in G .

The next theorem is a key result of interesting book [4]. Theorem 3.6 of [4] may be stated as follows:

Theorem 7. *Let Ω be bounded, connected and Steiner symmetric relative to the hyperplane $P = \{x = (x_1, x') : x_1 = 0\}$. Assume that $u \in C(\overline{\Omega}) \cap C^1(\Omega)$ is a positive weak solution of the boundary value problem*

$$\begin{cases} -\Delta u = f_1(u) + f_2(u) & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases}$$

where $f_1: [0, \infty) \rightarrow \mathbb{R}$ is locally Lipschitz continuous, while $f_2: [0, \infty) \rightarrow \mathbb{R}$ is non-decreasing and is identically zero on an interval $[0, h]$ for some $h > 0$. Then

$$u(-x_1, x') = u(x_1, x') \quad \text{for } (x_1, x') \in \Omega.$$

Moreover,

$$\frac{\partial u}{\partial x_1}(x_1, x') < 0 \quad \text{if } (x_1, x') \in \Omega \text{ and } x_1 > 0.$$

A proof of Theorem 7 is an essential part of book [4]. We are now in a position to prove the following theorem.

Theorem 8. *Let $p = 2$. If the domain Ω is Steiner symmetric with respect to a hyperplane P , then for any optimal pair (u, D) both u and D are symmetric with respect to P , and D^c is Steiner symmetric with respect to P .*

Proof. Without loss of generality, we may assume that

$$P = \{x = (x_1, x') : x_1 = 0\}.$$

Let (u, D) be an optimal solution. It follows from assertion (a) of Theorem 4 that $u \in C^1(\Omega)$. Next, the assumption that Ω satisfies the exterior cone condition at each point of $\partial\Omega$ yields $u \in C(\bar{\Omega})$ (see, e.g., [5], Theorem 8.30). By statement (b) of Theorem 4, there exists t such that

$$D = \{u \leq t\} = \{u - t \leq 0\}.$$

Suppose that $t > 0$. Using the Heaviside function $H: \mathbb{R} \rightarrow \mathbb{R}$ defined by

$$H(s) := \begin{cases} 0 & \text{if } s < 0, \\ 1 & \text{if } s \geq 0, \end{cases}$$

we observe that

$$\chi_D = H(t - u) \quad \text{in } \Omega.$$

Since

$$\begin{cases} -\Delta u + \alpha \chi_D u = \lambda_1 u & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases}$$

where $\lambda_1 = \Lambda(\alpha, A)$, we see that the eigenfunction u is a weak solution of the problem

$$\begin{cases} -\Delta u = \lambda_1 u - \alpha H(t - u)u & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega. \end{cases}$$

An application of Theorem 7 with

$$f_1(u) = (\lambda_1 - \alpha)u, \quad f_2(u) = \alpha(1 - H(t - u))u$$

and $h = t$ gives the desired result. In the case of $t = 0$, corresponding to the assumption that $A = 0$, there is $|D| = 0$ and thus

$$\begin{cases} -\Delta u = \lambda_1 u & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases}$$

so that Theorem 7 may be applied again. This completes the proof. □

REFERENCES

- [1] S. Chanillo, D. Grieser, M. Imai, K. Kurata, I. Ohnishi, *Symmetry breaking and other phenomena in the optimization of eigenvalues for composite membranes*, Comm. Math. Phys. **214** (2000), 315–337.
- [2] L. Damascelli, B. Sciunzi, *Regularity, monotonicity and symmetry of positive solutions of m -Laplace equations*, J. Differential Equations **206** (2004), 483–515.
- [3] P. Drabek, *Strongly nonlinear degenerated and singular elliptic problems*, in: Nonlinear Partial Differential Equations (Fès, 1994), 112–146, Pitman Research Notes in Mathematics Series **343**, Longman, Harlow, 1996.
- [4] L.E. Fraenkel, *Introduction to Maximum Principles and Symmetry in Elliptic Problems*, Cambridge University Press, Cambridge, 2000.
- [5] D. Gilbarg, N.S. Trudinger, *Elliptic Partial Differential Equations of Second Order*, Spriger-Verlag, Berlin, 1983.
- [6] K. Kurata, M. Shibata, S. Sakamoto, *Symmetry-Breaking Phenomena in an Optimization Problem for Some Nonlinear Elliptic Equation*, Appl. Math. Optim. **50** (2004), 259–278.
- [7] H. Lou, On singular sets of local solutions to p -Laplace equations, preprint.
- [8] E.H. Lieb, M. Loss, *Analysis, 2nd ed.*, AMS Graduate Studies in Mathematics **14**, Providence 2001.
- [9] W. Pielichowski, *The optimization of eigenvalue problems involving the p -Laplacian*, Univ. Jagell. Acta. Math. **42** (2004), 109–122.
- [10] W. Pielichowski, *On the first eigenvalue of a quasilinear elliptic operator*, Selected Problems of Mathematics, Anniversary Issue **6**, Cracow University of Technology, Cracow, 1995, 235–241.

Wacław Pielichowski
wpielich@pk.edu.pl

Cracow University of Technology
Institute of Mathematics
ul. Warszawska 24, 31-155 Cracow, Poland

Received: December 7, 2007.

Revised: March 3, 2008.

Accepted: March 3, 2008.