

FIXED POINTS AND STABILITY IN NEUTRAL NONLINEAR DIFFERENTIAL EQUATIONS WITH VARIABLE DELAYS

Abdelouaheb Ardjouni and Ahcene Djoudi

Abstract. By means of Krasnoselskii's fixed point theorem we obtain boundedness and stability results of a neutral nonlinear differential equation with variable delays. A stability theorem with a necessary and sufficient condition is given. The results obtained here extend and improve the work of C.H. Jin and J.W. Luo [Nonlinear Anal. 68 (2008), 3307–3315], and also those of T.A. Burton [Fixed Point Theory 4 (2003), 15–32; Dynam. Systems Appl. 11 (2002), 499–519] and B. Zhang [Nonlinear Anal. 63 (2005), e233–e242]. In the end we provide an example to illustrate our claim.

Keywords: fixed points, stability, nonlinear neutral differential equation, integral equation, variable delays.

Mathematics Subject Classification: 34K20, 34K30, 34K40.

1. INTRODUCTION

Certainly, the Lyapunov direct method has been, for more than 100 years, the main tool for the study of stability properties of ordinary, functional and partial differential equations. Nevertheless, the application of this method to problems of stability in differential equations with delay has encountered serious difficulties if the delay is unbounded or if the equation has unbounded terms [1–3]. Recently, T.A. Burton and T. Furumochi have noticed that some of these difficulties vanish or might be overcome by means of fixed point theory (see [5–8]). The fixed point theory does not only solve the problem on stability but has a significant advantage over Lyapunov's direct method. The conditions of the former are often averages but those of the latter are usually pointwise (see [1]).

In this paper we consider the neutral nonlinear differential equation with variable delays

$$x'(t) = -a(t)x(t - r_1(t)) + b(t)x'(t - r_1(t)) + c(t)G(x^\gamma(t - r_2(t))), \quad (1.1)$$

with the initial condition

$$x(t) = \psi(t) \quad \text{for } t \in [m(0), 0],$$

where $\psi \in C([m(0), 0], \mathbb{R})$, $m_j(0) = \inf\{t - r_j(t), t \geq 0\}$, $m(0) = \min\{m_j(0), j = 1, 2\}$, $\gamma \in (0, 1)$ and γ is a quotient with odd positive integer denominator. Throughout this paper we assume that $a, c \in C(\mathbb{R}^+, \mathbb{R})$, $b \in C^1(\mathbb{R}^+, \mathbb{R})$ and $r_1 \in C^2(\mathbb{R}^+, \mathbb{R}^+)$, $r_2 \in C(\mathbb{R}^+, \mathbb{R}^+)$ with $t - r_j(t) \rightarrow \infty$ as $t \rightarrow \infty$, $j = 1, 2$. We also assume that $G(\cdot)$ is locally Lipschitz continuous in x . That is, there is a $L > 0$ so that if $|x|, |y| \leq 1$ then

$$|G(x) - G(y)| \leq L|x - y| \quad \text{and} \quad G(0) = 0.$$

Special cases of equation (1.1) have been previously considered and studied under various conditions. Particularly, T.A. Burton in [3] and B. Zhang in [12] have investigated the boundedness and stability of the linear equation

$$x'(t) = -a(t)x(t - r_1(t)).$$

In [7], T.A. Burton and T. Furumochi have studied the boundedness and the asymptotic stability by using Krasnoselskii fixed point theorem for the following equation:

$$x'(t) = -a(t)x(t - r_1) + b(t)x^{\frac{1}{3}}(t - r_2(t)),$$

with $r_1 \geq 0$ is a constant and $a \in C(\mathbb{R}^+, (0, \infty))$. By letting $\gamma = 1/3$, $G(x) = x$ and $b(t) = 0$ in equation (1.1), C.H. Jin and J.W. Luo [9] studied, by means of Krasnoselskii's fixed point theorem, the boundedness and stability, under appropriate conditions, of the following equation:

$$x'(t) = -a(t)x(t - r_1(t)) + b(t)x^{\frac{1}{3}}(t - r_2(t)),$$

and generalized the results claimed in [3, 7, 12].

Our purpose here is to give, by using Krasnoselskii fixed point theorem, boundedness and stability results for the nonlinear neutral differential equation with variable delays (1.1).

In Section 2, we present the inversion of equation (1.1) and we state the hybrid Krasnoselskii's fixed point theorem. For details on Krasnoselskii theorem we refer the reader to [1, 11]. We present our main results on stability in Section 3 and at the end we provide an example to illustrate this work.

2. INVERSION OF EQUATION (1.1)

We have to invert equation (1.1). For this, we use the variation of parameter formula to rewrite the equation as an integral equation suitable for Krasnoselskii theorem. Besides, the integration by parts will be applied. In our consideration we suppose that

$$r_1'(t) \neq 1, \quad \forall t \in \mathbb{R}^+. \quad (2.1)$$

Lemma 2.1. *Let $g : [m(0), \infty) \rightarrow \mathbb{R}^+$ be an arbitrary continuous function and suppose that (2.1) holds. Then x is a solution of (1.1) if and only if*

$$\begin{aligned}
 x(t) = & \left(x(0) - \frac{b(0)}{1 - r_1'(0)} x(-r_1(0)) - \int_{-r_1(0)}^0 g(u)x(u)du \right) e^{-\int_0^t g(u)du} + \\
 & + \frac{b(t)}{1 - r_1'(t)} x(t - r_1(t)) + \int_{t-r_1(t)}^t g(u)x(u)du - \\
 & - \int_0^t e^{-\int_s^t g(u)du} g(s) \left(\int_{s-r_1(s)}^s g(u)x(u)du \right) ds + \\
 & + \int_0^t e^{-\int_s^t g(u)du} [g(s - r_1(s))(1 - r_1'(s)) - a(s) - \mu(s)] x(s - r_1(s)) ds + \\
 & + \int_0^t e^{-\int_s^t g(u)du} c(s)G(x^\gamma(s - r_2(s))) ds,
 \end{aligned} \tag{2.2}$$

where

$$\mu(s) = \frac{(b'(s) + b(s)g(s))(1 - r_1'(s)) + r_1''(s)b(s)}{(1 - r_1'(s))^2}. \tag{2.3}$$

Proof. Let x be a solution of (1.1). Rewrite equation (1.1) as

$$\begin{aligned}
 x'(t) = & -g(t)x(t) + (d/dt) \int_{t-r_1(t)}^t g(s)x(s)ds + \\
 & + [g(t - r_1(t))(1 - r_1'(t)) - a(t)] x(t - r_1(t)) + \\
 & + b(t)x'(t - r_1(t)) + c(t)G(x^\gamma(t - r_2(t))).
 \end{aligned}$$

Multiply both sides of the above equation by $e^{\int_0^t g(s)ds}$ and then integrate from 0 to t to obtain

$$\begin{aligned}
 x(t) = & x(0)e^{-\int_0^t g(s)ds} + \int_0^t e^{-\int_s^t g(u)du} d \left(\int_{s-r_1(s)}^s g(u)x(u)du \right) + \\
 & + \int_0^t e^{-\int_s^t g(u)du} [g(s - r_1(s))(1 - r_1'(s)) - a(s)] x(s - r_1(s)) ds + \\
 & + \int_0^t e^{-\int_s^t g(u)du} b(s)x'(s - r_1(s)) ds + \int_0^t e^{-\int_s^t g(u)du} c(s)G(x^\gamma(s - r_2(s))) ds.
 \end{aligned} \tag{2.4}$$

Letting

$$\begin{aligned} & \int_0^t e^{-\int_s^t g(u)du} b(s) x'(s - r_1(s)) ds = \\ & = \int_0^t e^{-\int_s^t g(u)du} \frac{b(s)}{(1 - r_1'(s))} (1 - r_1'(s)) x'(s - r_1(s)) ds. \end{aligned}$$

By performing an integration by parts, we have

$$\begin{aligned} & \int_0^t e^{-\int_s^t g(u)du} b(s) x'(s - r_1(s)) ds = \\ & = \frac{b(t)}{1 - r_1'(t)} x(t - r_1(t)) - \frac{b(0)}{1 - r_1'(0)} x(-r_1(0)) e^{-\int_0^t g(u)du} - \\ & - \int_0^t e^{-\int_s^t g(u)du} \mu(s) x(s - r_1(s)) ds, \end{aligned} \tag{2.5}$$

where $\mu(s)$ is given by (2.3), and

$$\begin{aligned} & \int_0^t e^{-\int_s^t g(u)du} d\left(\int_{s-r_1(s)}^s g(u)x(u)du \right) = \\ & = -e^{-\int_0^t g(u)du} \int_{-r_1(0)}^0 g(u)x(u)du + \int_{t-r_1(t)}^t g(u)x(u)du - \\ & - \int_0^t e^{-\int_s^t g(u)du} g(s) \left(\int_{s-r_1(s)}^s g(u)x(u)du \right) ds. \end{aligned} \tag{2.6}$$

Finally, substituting (2.5) and (2.6) into (2.4) ends the proof. \square

Lastly in this section, we state Krasnoselskii's fixed point theorem which enables us to prove the stability of the zero solution. For its proof we refer the reader to [1,11]. Krasnoselskii's theorem is a captivating tool it is not only used to solve old problems of existence in nonlinear analysis but it does, as well, solve stability of hard problems which have frustrated investigators for many years using others method.

Theorem 2.2 (Krasnoselskii). *Let M be a closed convex nonempty subset of a Banach space $(S, \|\cdot\|)$. Suppose that A and B map M into S such that:*

- (i) $x, y \in M$, implies $Ax + By \in M$,
- (ii) A is continuous and AM is contained in a compact set,
- (iii) B is a contraction with constant $\alpha < 1$.

Then there exists $z \in M$ with $z = Az + Bz$.

3. STABILITY BY KRASNOSELSKII FIXED POINT THEOREM

From existence theory, which can be found in [1], we conclude that for each continuous initial function $\psi : [m(0), 0] \rightarrow \mathbb{R}$, there is a continuous solution $x(t, 0, \psi)$ on an interval $[0, T)$ for some $T > 0$ and $x(t, 0, \psi) = \psi(t)$ on $[m(0), 0]$. For stability definitions we refer to [1].

Theorem 3.1. *Let (2.1) holds and suppose that there are constants $\alpha \in (0, 1)$, $k_1, k_2 > 0$ and a function $g \in C([m(0), \infty), \mathbb{R}^+)$ such that for $|t_2 - t_1| \leq 1$,*

$$\left| \int_{t_1}^{t_2} |c(u)| du \right| \leq k_1 |t_1 - t_2|, \tag{3.1}$$

and

$$\left| \int_{t_1}^{t_2} g(u) du \right| \leq k_2 |t_1 - t_2|, \tag{3.2}$$

while for $t \geq 0$

$$\begin{aligned} & \left| \frac{b(t)}{1 - r_1'(t)} \right| + \int_{t-r_1(t)}^t g(u) du + \int_0^t e^{-\int_s^t g(u) du} g(s) \left(\int_{s-r_1(s)}^s g(u) du \right) ds + \\ & + \int_0^t e^{-\int_s^t g(u) du} \{ |g(s - r_1(s)) (1 - r_1'(s)) - a(s) - \mu(s)| + L|c(s)| \} ds \leq \alpha. \end{aligned} \tag{3.3}$$

If ψ is a given continuous initial function which is sufficiently small, then there is a solution $x(t, 0, \psi)$ of (1.1) on \mathbb{R}^+ with $|x(t, 0, \psi)| \leq 1$.

Proof. For $\alpha \in (0, 1)$, find an appropriate $\delta > 0$ such that

$$\left(1 + \left| \frac{b(0)}{1 - r_1'(0)} \right| + \int_{-r_1(0)}^0 g(u) du \right) e^{-\int_0^t g(s) ds} \delta + \alpha \leq 1.$$

Let $\psi : [m(0), 0] \rightarrow \mathbb{R}$ be a given small bounded initial function with $\|\psi\| < \delta$. In the same context as in papers [1, 7, 9], let $h : [m(0), \infty) \rightarrow [1, \infty)$ be any strictly increasing and continuous function with $h(m(0)) = 1$, $h(s) \rightarrow \infty$ as $s \rightarrow \infty$, such that

$$\begin{aligned} & \left| \frac{b(t)}{1 - r_1'(t)} \right| + \int_{t-r_1(t)}^t g(u) h(u) / h(t) du + \int_0^t e^{-\int_s^t g(u) du} g(s) \left(\int_{s-r_1(s)}^s g(u) h(u) / h(t) du \right) ds + \\ & + \int_0^t e^{-\int_s^t g(u) du} |g(s - r_1(s)) (1 - r_1'(s)) - a(s) - \mu(s)| h(s - r_1(s)) / h(t) ds \leq \alpha. \end{aligned} \tag{3.4}$$

Let $(S, |\cdot|_h)$ be the Banach space of continuous $\varphi : [m(0), \infty) \rightarrow \mathbb{R}$ with

$$|\varphi|_h := \sup_{t \geq m(0)} |\varphi(t)/h(t)| < \infty,$$

and define the set S_ψ by

$$S_\psi = \{\varphi \in S : |\varphi(t)| \leq 1 \text{ for } t \in [m(0), \infty) \text{ and } \varphi(t) = \psi(t) \text{ if } t \in [m(0), 0]\}.$$

Define the mappings $A, B : S_\psi \rightarrow S_\psi$ by

$$(A\varphi)(t) = \int_0^t e^{-\int_s^t g(u)du} c(s)G(\varphi^\gamma(s - r_2(s))) ds, \quad (3.5)$$

and

$$\begin{aligned} (B\varphi)(t) = & \left(\varphi(0) - \frac{b(0)}{1 - r_1'(0)} \varphi(-r_1(0)) - \int_{-r_1(0)}^0 g(u)\varphi(u)du \right) e^{-\int_0^t g(s)ds} + \\ & + \frac{b(t)}{1 - r_1'(t)} \varphi(t - r_1(t)) + \int_{t-r_1(t)}^t g(u)\varphi(u)du - \\ & - \int_0^t e^{-\int_s^t g(u)du} g(s) \left(\int_{s-r_1(s)}^s g(u)\varphi(u)du \right) ds + \\ & + \int_0^t e^{-\int_s^t g(u)du} [g(s - r_1(s))(1 - r_1'(s)) - a(s) - \mu(s)] \varphi(s - r_1(s)) ds. \end{aligned} \quad (3.6)$$

That A maps S_ψ into itself can be deduced from condition (3.3).

We now show that $\varphi, \phi \in S_\psi$ implies that $A\varphi + B\phi \in S_\psi$. When doing this we see that also B maps S_ψ into itself by letting $\varphi = 0$ in the preceding sum. Now, let $\|\cdot\|$ be the supremum norm on $[m(0), \infty)$ of $\varphi \in S_\psi$ if φ is bounded. Note that if $\varphi, \phi \in S_\psi$

then

$$\begin{aligned}
 & |(A\varphi)(t) + (B\phi)(t)| \leq \\
 & \leq \left(1 + \left| \frac{b(0)}{1 - r_1'(0)} \right| + \int_{-r_1(0)}^0 g(u)du \right) e^{-\int_0^t g(s)ds} \|\psi\| + \\
 & + \|\phi\| \left| \frac{b(t)}{1 - r_1'(t)} \right| + \\
 & + \|\phi\| \int_{t-r_1(t)}^t g(u)du + \\
 & + |\phi| \int_0^t e^{-\int_s^t g(u)du} g(s) \left(\int_{s-r_1(s)}^s g(u)du \right) ds + \\
 & + |\phi| \int_0^t e^{-\int_s^t g(u)du} \{ |g(s - r_1(s))(1 - r_1'(s)) - a(s) - \mu(s)| \} ds + \\
 & + \|\varphi\|^\gamma \int_0^t e^{-\int_s^t g(u)du} L|c(s)|ds \leq \\
 & \leq \left(1 + \left| \frac{b(0)}{1 - r_1'(0)} \right| + \int_{-r_1(0)}^0 g(u)du \right) e^{-\int_0^t g(s)ds} \delta + \alpha \leq 1.
 \end{aligned}$$

Next, we show that AS_ψ is equicontinuous. If $\varphi \in S_\psi$ and $0 \leq t_1 < t_2$ with $t_2 - t_1 < 1$

$$\begin{aligned}
& |(A\varphi)(t_2) - (A\varphi)(t_1)| = \\
& = \left| \int_0^{t_2} e^{-\int_s^{t_2} g(u)du} c(s) G(\varphi^\gamma(s - r_2(s))) ds - \right. \\
& \quad \left. - \int_0^{t_1} e^{-\int_s^{t_1} g(u)du} c(s) G(\varphi^\gamma(s - r_2(s))) ds \right| \leq \\
& \leq \left| \int_{t_1}^{t_2} e^{-\int_s^{t_2} g(u)du} c(s) G(\varphi^\gamma(s - r_2(s))) ds \right| + \\
& \quad + \left| \int_0^{t_1} \left[e^{-\int_s^{t_2} g(u)du} - e^{-\int_s^{t_1} g(u)du} \right] c(s) G(\varphi^\gamma(s - r_2(s))) ds \right| \leq \\
& \leq L \int_{t_1}^{t_2} e^{-\int_s^{t_2} g(u)du} d \left(\int_{t_1}^s |c(s)| ds \right) + \\
& \quad + L \left| e^{-\int_{t_1}^{t_2} g(u)du} - e^{-\int_{t_1}^{t_1} g(u)du} \right| e^{\int_0^{t_1} g(u)du} \int_0^{t_1} e^{-\int_s^{t_1} g(u)du} |c(s)| ds \leq \\
& \leq L \int_{t_1}^{t_2} |c(u)| du \left(1 + \int_{t_1}^{t_2} e^{-\int_s^{t_2} g(u)du} g(s) ds \right) + \alpha \left| e^{-\int_{t_1}^{t_2} g(u)du} - 1 \right| \leq \\
& \leq 2L \int_{t_1}^{t_2} |c(u)| du + \alpha \left| \int_{t_1}^{t_2} g(u) du \right| \leq \\
& \leq (2Lk_1 + \alpha k_2) |t_2 - t_1|,
\end{aligned}$$

by (3.1)–(3.3). In the space $(S, |\cdot|_h)$, the set AS_ψ resides in a compact set.

Now we show that B is a contraction with constant α . We have

$$\begin{aligned}
& |(B\phi_1)(t) - (B\phi_2)(t)|/h(t) \leq \\
& \leq \left| \frac{b(t)}{1 - r_1'(t)} \right| |\phi_1(t) - \phi_2(t)|/h(t) + \int_{t-r_1(t)}^t g(u) |\phi_1(u) - \phi_2(u)|/h(t) du + \\
& + \int_0^t e^{-\int_s^t g(u) du} g(s) \left(\int_{s-r_1(s)}^s g(u) |\phi_1(u) - \phi_2(u)|/h(t) du \right) ds + \\
& + \int_0^t e^{-\int_s^t g(u) du} |g(s - r_1(s)) (1 - r_1'(s)) - a(s) - \mu(s)| \times \\
& \times |\phi_1(s - r_1(s)) - \phi_2(s - r_1(s))|/h(t) ds \leq \\
& \leq |\phi_1 - \phi_2|_h \left\{ \left| \frac{b(t)}{1 - r_1'(t)} \right| + \int_{t-r_1(t)}^t g(u) h(u)/h(t) du + \right. \\
& + \int_0^t e^{-\int_s^t g(u) du} g(s) \left(\int_{s-r_1(s)}^s g(u) h(u)/h(t) du \right) ds + \\
& \left. + \int_0^t e^{-\int_s^t g(u) du} |g(s - r_1(s)) (1 - r_1'(s)) - a(s) - \mu(s)| h(s - r_1(s)) / h(t) ds \right\} \leq \\
& \leq \alpha |\phi_1 - \phi_2|_h,
\end{aligned}$$

by (3.4).

Finally, we need to show that A is continuous. Let $\epsilon > 0$ be given and let $\varphi \in S_\psi$. Now x^γ is uniformly continuous on $[-1, 1]$ so for a fixed $T > 0$ with $4/h(t) < \epsilon$ there is an $\eta > 0$ such that $|x_1 - x_2| < \eta h(t)$ implies $|x_1^\gamma - x_2^\gamma| < \epsilon/2$. Thus for $|\varphi(t) - \phi(t)| < \eta h(t)$ and for $t > T$ we have

$$\begin{aligned}
& |(A\varphi)(t) - (A\phi)(t)|/h(t) \leq \\
& \leq (1/h(t)) \int_0^t e^{-\int_s^t g(u)du} |c(s)| |G(\varphi^\gamma(s - r_2(s))) - G(\phi^\gamma(s - r_2(s)))| ds \leq \\
& \leq L(1/h(t)) \int_0^t e^{-\int_s^t g(u)du} |c(s)| |\varphi^\gamma(s - r_2(s)) - \phi^\gamma(s - r_2(s))| ds \leq \\
& \leq L(1/h(t)) \left\{ \int_0^T e^{-\int_s^t g(u)du} |c(s)| |\varphi^\gamma(s - r_2(s)) - \phi^\gamma(s - r_2(s))| ds + \right. \\
& \left. + 2 \int_T^t |c(s)| e^{-\int_s^t g(u)du} ds \right\} \leq L \{(\alpha\epsilon)/(2h(t)) + 2\alpha/h(t)\} \leq \\
& \leq L \{(\alpha\epsilon/2) + (2\alpha/h(t))\} < L\alpha\epsilon.
\end{aligned}$$

The conditions of Krasnoselskii's theorem are satisfied and there is a fixed point. This completes the proof. \square

Letting $r_1(t) = r_1$, a constant, and $g(t) = a(t + r_1)$ with $a \in C(\mathbb{R}^+, \mathbb{R}^+)$, we obtain the following corollary.

Corollary 3.2. *Let (3.1) and (3.2) hold and (3.3) be replaced by*

$$\begin{aligned}
|b(t)| + \int_{t-r_1}^t a(u+r_1)du + \int_0^t e^{-\int_s^t a(u+r_1)du} a(s+r_1) \left(\int_{s-r_1}^s a(u+r_1)du \right) ds + \\
+ \int_0^t e^{-\int_s^t a(u+r_1)du} (|b'(s) + b(s)a(s+r_1)| + L|c(s)|) ds \leq \alpha.
\end{aligned} \tag{3.7}$$

If ψ is a given continuous initial function which is sufficiently small, then there is a solution $x(t, 0, \psi)$ of (1.1) on \mathbb{R}^+ with $|x(t, 0, \psi)| \leq 1$.

Letting $\gamma = 1/3$, $b(t) = 0$ and $G(x) = x$, we have the following corollary.

Corollary 3.3. *Let (3.1) and (3.2) hold and (3.3) be replaced by*

$$\begin{aligned}
\int_{t-r_1(t)}^t g(u)du + \int_0^t e^{-\int_s^t g(u)du} g(s) \left(\int_{s-r_1(s)}^s g(u)du \right) ds + \\
+ \int_0^t e^{-\int_s^t g(u)du} \{ |g(s - r_1(s))(1 - r_1'(s)) - a(s)| + |c(s)| \} ds \leq \alpha.
\end{aligned} \tag{3.8}$$

If ψ is a given continuous initial function which is sufficiently small, then there is a solution $x(t, 0, \psi)$ of (1.1) on \mathbb{R}^+ with $|x(t, 0, \psi)| \leq 1$.

Remark 3.4. Obviously Corollary 3.3 reduces to Theorem 2.1 of [9]. Thus, Theorem 3.1 above is a generalization of Theorem 2.1 of [9].

Theorem 3.5. *Let (2.1) and (3.1)–(3.3) hold and assume that*

$$\int_0^t e^{-\int_s^t g(u)du} |c(s)| ds \rightarrow 0 \quad \text{as } t \rightarrow \infty. \tag{3.9}$$

If ψ is a given continuous initial function which is sufficiently small, then (1.1) has a solution $x(t, 0, \psi) \rightarrow 0$ as $t \rightarrow \infty$ if and only if

$$\int_0^t g(s) ds \rightarrow \infty \quad \text{as } t \rightarrow \infty. \tag{3.10}$$

Proof. First, suppose that (3.10) holds. We set

$$N = \sup_{t \geq 0} \left\{ e^{-\int_0^t g(s) ds} \right\}. \tag{3.11}$$

All of the calculations in the proof of Theorem 3.1 hold with $h(t) = 1$ when $|\cdot|_h$ is replaced by the supremum norm $\|\cdot\|$. For $\varphi \in S_\psi$, we have

$$|(A\varphi)(t)| \leq L \int_0^t e^{-\int_s^t g(u)du} |c(s)| ds =: q(t), \tag{3.12}$$

where $q(t) \rightarrow 0$ as $t \rightarrow \infty$ by (3.9).

Add to S_ψ the condition that $\varphi \in S_\psi$ implies that $\varphi(t) \rightarrow 0$ as $t \rightarrow \infty$. We can see that for $\varphi \in S_\psi$ then $(A\varphi)(t) \rightarrow 0$ as $t \rightarrow \infty$ by (3.12) and $(B\phi)(t) \rightarrow 0$ as $t \rightarrow \infty$ by (3.10). Since AS_ψ has been shown to be equicontinuous, A maps S_ψ into a compact subset of S_ψ (see [1], Theorem 1.2.2 on p. 20). By Krasnoselskii’s theorem there is an $x \in S_\psi$ with $Ax + Bx = x$. As $x \in S_\psi$, $x(t) \rightarrow 0$ as $t \rightarrow \infty$.

Conversely, suppose (3.10) fails. Then there exists a sequence $\{t_n\}$, $t_n \rightarrow \infty$ as $n \rightarrow \infty$ such that $\lim_{n \rightarrow \infty} \int_0^{t_n} g(u)du = l$ for some $l \in \mathbb{R}$. We may also choose a positive constant J satisfying

$$-J \leq \int_0^{t_n} g(s) ds \leq J,$$

for all $n \geq 1$. To simplify the expression, we define

$$\omega(s) = |g(s - r_1(s))(1 - r_1'(s)) - a(s) - \mu(s)| + L|c(s)| + g(s) \int_{s-r_1(s)}^s g(u)du,$$

for all $s \geq 0$. By (3.3), we have

$$\int_0^{t_n} e^{-\int_s^{t_n} g(u)du} \omega(s) ds \leq \alpha.$$

This yields

$$\int_0^{t_n} e^{\int_0^s g(u)du} \omega(s) ds \leq \alpha e^{\int_0^{t_n} g(u)du} \leq J.$$

The sequence $\left\{ \int_0^{t_n} e^{\int_0^s g(u)du} \omega(s) ds \right\}$ is bounded, so there exists a convergent subsequence. For brevity of notation, we may assume

$$\lim_{n \rightarrow \infty} \int_0^{t_n} e^{\int_0^s g(u)du} \omega(s) ds = \lambda,$$

for some $\lambda \in \mathbb{R}^+$ and choose a positive integer m so large that

$$\int_{t_m}^{t_n} e^{\int_0^s g(u)du} \omega(s) ds < \delta_0/4N,$$

for all $n \geq m$, where $\delta_0 > 0$ satisfies $2\delta_0 N e^J + \alpha \leq 1$.

We now consider the solution $x(t) = x(t, t_m, \psi)$ of (1.1) with $\psi(t_m) = \delta_0$ and $|\psi(s)| \leq \delta_0$ for $s \leq t_m$. We may choose ψ so that $|x(t)| \leq 1$ for $t \geq t_m$ and

$$\psi(t_m) - \frac{b(t_m)}{1 - r_1'(t_m)} \psi(t_m - r_1(t_m)) - \int_{t_m - r_1(t_m)}^{t_m} g(s) \psi(s) ds \geq \frac{1}{2} \delta_0.$$

It follows from (3.5) and (3.6) with $x(t) = (Ax)(t) + (Bx)(t)$ that for $n \geq m$

$$\begin{aligned}
 & \left| x(t_n) - \frac{b(t_n)}{1 - r_1'(t_n)} x(t_n - r_1(t_n)) - \int_{t_n - r_1(t_n)}^{t_n} g(s)x(s)ds \right| \geq \\
 & \geq \frac{1}{2} \delta_0 e^{-\int_{t_m}^{t_n} g(u)du} - \int_{t_m}^{t_n} e^{-\int_s^{t_n} g(u)du} \omega(s)ds = \\
 & = \frac{1}{2} \delta_0 e^{-\int_{t_m}^{t_n} g(u)du} - e^{-\int_0^{t_n} g(u)du} \int_{t_m}^{t_n} e^{\int_0^s g(u)du} \omega(s)ds = \\
 & = e^{-\int_{t_m}^{t_n} g(u)du} \left(\frac{1}{2} \delta_0 - e^{-\int_0^{t_m} g(u)du} \int_{t_m}^{t_n} e^{\int_0^s g(u)du} \omega(s)ds \right) \geq \\
 & \geq e^{-\int_{t_m}^{t_n} g(u)du} \left(\frac{1}{2} \delta_0 - N \int_{t_m}^{t_n} e^{\int_0^s g(u)du} \omega(s)ds \right) \geq \\
 & \geq \frac{1}{4} \delta_0 e^{-\int_{t_m}^{t_n} g(u)du} \geq \frac{1}{4} \delta_0 e^{-2J} > 0.
 \end{aligned} \tag{3.13}$$

On the other hand, if the solution of (1.1) $x(t) = x(t, t_m, \psi) \rightarrow 0$ as $t \rightarrow \infty$, since $t_n - r_1(t_n) \rightarrow \infty$ as $n \rightarrow \infty$ and (3.3) holds, we have

$$x(t_n) - \frac{b(t_n)}{1 - r_1'(t_n)} x(t_n - r_1(t_n)) - \int_{t_n - r_1(t_n)}^{t_n} g(s)x(s)ds \rightarrow 0 \text{ as } n \rightarrow \infty,$$

which contradicts (3.13). Hence condition (3.10) is necessary in order that (1.1) has a solution $x(t, 0, \psi) \rightarrow 0$ as $t \rightarrow \infty$. The proof is complete. \square

Letting $r_1(t) = r_1$, a constant, and $g(t) = a(t + r_1)$ with $a \in C(\mathbb{R}^+, \mathbb{R}^+)$, we have the following corollary.

Corollary 3.6. *Let (3.1), (3.2), and (3.9) hold and (3.3) be replaced by (3.7). If ψ is a given continuous initial function which is sufficiently small, then (1.1) has a solution $x(t, 0, \psi) \rightarrow 0$ as $t \rightarrow \infty$ if and only if*

$$\int_0^t g(s)ds \rightarrow \infty \text{ as } t \rightarrow \infty.$$

For the case $\gamma = 1/3$, $b(t) = 0$ and $G(x) = x$, we have the following corollary.

Corollary 3.7. *Let (3.1), (3.2), and (3.9) hold and (3.3) be replaced by (3.8). If ψ is a given continuous initial function which is sufficiently small, then (1.1) has a solution $x(t, 0, \psi) \rightarrow 0$ as $t \rightarrow \infty$ if and only if*

$$\int_0^t g(s) ds \rightarrow \infty \quad \text{as } t \rightarrow \infty.$$

Remark 3.8. Corollary 3.7 reduces to Theorem 2.2 of [9].

Example 3.9. Let

$$x'(t) = -a(t)x(t - r_1(t)) + b(t)x'(t - r_1(t)) + c(t)G(x^\gamma(t - r_2(t))), \quad (3.14)$$

where $\gamma = 1/3$, $G(x) = \sin(x)$, $r_1(t) = 0.232t$, $r_2 \in C(\mathbb{R}^+, \mathbb{R}^+)$ with $t - r_2(t) \rightarrow \infty$ as $t \rightarrow \infty$, $a(t) = 0.768/(0.768t + 1)$, $b(t) = t/(16t + 16)$, $c(t) = 1/(8(t + 1)^2)$.

Then for any small continuous initial function ψ , every solution $x(t, 0, \psi)$ of the nonlinear neutral differential equation (3.14) goes to 0 as $t \rightarrow \infty$.

Indeed, clearly $G(0) = 0$ and $G(x)$ is locally Lipschitz continuous in x . Let $|x|, |y| \leq 1$, then

$$|G(x) - G(y)| = |\sin(x) - \sin(y)| \leq |x - y|.$$

Choosing $g(t) = 1/(t + 1)$, we have

$$\begin{aligned} \int_{t-r_1(t)}^t g(s) ds &= \int_{0.768t}^t 1/(s+1) ds = \ln\left(\frac{t+1}{0.768t+1}\right) < 0.264, \\ \int_0^t e^{-\int_s^t g(u) du} g(s) \left(\int_{s-r_1(s)}^s g(u) du \right) ds &< 0.264, \\ \int_0^t e^{-\int_s^t g(u) du} |g(s-r_1(s))(1-r_1'(s)) - a(s) - \mu(s)| ds &= \\ &= (1/(16 \cdot 0.768)) \int_0^t e^{-\int_s^t \frac{1}{u+1} du} \frac{1}{s+1} ds < 0.082, \\ \int_0^t e^{-\int_s^t g(u) du} L|c(s)| ds &\leq 0.125, \end{aligned}$$

and

$$\left| \frac{b(t)}{1-r_1'(t)} \right| < 0.082.$$

Let $\alpha = 0.082 + 0.264 + 0.264 + 0.082 + 0.125 = 0.817 < 1$, then by Theorem 3.5, every solution $x(t, 0, \psi)$ of (3.14) with small continuous initial function ψ , goes to zero as $t \rightarrow \infty$.

REFERENCES

- [1] T.A. Burton, *Stability by Fixed Point Theory for Functional Differential Equations*, Dover Publications, New York, 2006.
- [2] T.A. Burton, *Liapunov functionals, fixed points, and stability by Krasnoselskii's theorem*, *Nonlinear Stud.* **9** (2001), 181–190.
- [3] T.A. Burton, *Stability by fixed point theory or Liapunov's theory: A comparison*, *Fixed Point Theory* **4** (2003), 15–32.
- [4] T.A. Burton, *Fixed points and stability of a nonconvolution equation*, *Proc. Amer. Math. Soc.* **132** (2004), 3679–3687.
- [5] T.A. Burton, T. Furumochi, *A note on stability by Schauder's theorem*, *Funkcial. Ekvac.* **44** (2001), 73–82.
- [6] T.A. Burton, T. Furumochi, *Fixed points and problems in stability theory*, *Dynam. Systems Appl.* **10** (2001), 89–116.
- [7] T.A. Burton, T. Furumochi, *Asymptotic behavior of solutions of functional differential equations by fixed point theorems*, *Dynam. Systems Appl.* **11** (2002), 499–519.
- [8] T.A. Burton, T. Furumochi, *Krasnoselskii's fixed point theorem and stability*, *Nonlinear Anal.* **49** (2002), 445–454.
- [9] C.H. Jin, J.W. Luo, *Stability in functional differential equations established using fixed point theory*, *Nonlinear Anal.* **68** (2008), 3307–3315.
- [10] Y.N. Raffoul, *Stability in neutral nonlinear differential equations with functional delays using fixed-point theory*, *Math. Comput. Modelling* **40** (2004), 691–700.
- [11] D.R. Smart, *Fixed Point Theorems*, Cambridge Tracts in Mathematics, No. 66. Cambridge University Press, London-New York, 1974.
- [12] B. Zhang, *Fixed points and stability in differential equations with variable delays*, *Nonlinear Anal.* **63** (2005), e233–e242.

Abdelouaheb Ardjouni
abd_ardjouni@yahoo.fr

University of Annaba
Department of Mathematics
P.O. Box 12, Annaba 23000, Algeria

Ahcene Djoudi
adjoudi@yahoo.com

University of Annaba
Department of Mathematics
P.O. Box 12, Annaba 23000, Algeria

Received: May 11, 2011.

Accepted: June 13, 2011.