

This paper is dedicated with great esteem  
and admiration to Professor Leon Mikołajczyk.

## REMARKS FOR ONE-DIMENSIONAL FRACTIONAL EQUATIONS

Massimiliano Ferrara and Giovanni Molica Bisci

*Communicated by Marek Galewski*

**Abstract.** In this paper we study a class of one-dimensional Dirichlet boundary value problems involving the Caputo fractional derivatives. The existence of infinitely many solutions for this equations is obtained by exploiting a recent abstract result. Concrete examples of applications are presented.

**Keywords:** fractional differential equations, Caputo fractional derivatives, variational methods.

**Mathematics Subject Classification:** 34A08, 26A33, 35A15.

### 1. INTRODUCTION

The aim of this short note is to study nonlinear fractional boundary value problems whose general form is given by

$$\begin{cases} \Delta_{F,\alpha}u(t) + f(t, u(t)) = 0 & \text{a.e. } t \in [0, T], \\ u(0) = u(T) = 0, \end{cases}$$

where

$$\Delta_{F,\alpha}u(t) := \frac{d}{dt} \left( {}_0D_t^{\alpha-1} ({}^c_0D_t^\alpha u(t)) - {}_tD_T^{\alpha-1} ({}^c_tD_T^\alpha u(t)) \right),$$

$\alpha \in (1/2, 1]$ ,  ${}_0D_t^{\alpha-1}$  and  ${}_tD_T^{\alpha-1}$  are the left and right Riemann-Liouville fractional integrals of order  $1 - \alpha$  respectively,  ${}^c_0D_t^\alpha$  and  ${}^c_tD_T^\alpha$  are the left and right Caputo fractional derivatives of order  $\alpha$  respectively, and  $f : [0, T] \times \mathbb{R} \rightarrow \mathbb{R}$  is a continuous function.

Following [12], denote by  $C_0^\infty([0, T], \mathbb{R})$  the set of all functions  $g \in C^\infty([0, T], \mathbb{R})$  with  $g(0) = g(T) = 0$ . The fractional derivative Hilbert space  $E_0^\alpha$  is defined by the closure of  $C_0^\infty([0, T], \mathbb{R})$  with respect to the norm

$$\|u\| := \left( \int_0^T |{}_0^c D_t^\alpha u(t)|^2 dt + \int_0^T |u(t)|^2 dt \right)^{1/2}$$

for every  $u \in E_0^\alpha$ .

For basic facts and usual notation on the variational setting adopted here we refer the reader to [12, 18]. Let us denote

$$\begin{aligned} \kappa &:= \frac{T^{\alpha-\frac{1}{2}}}{\Gamma(\alpha)\sqrt{2\alpha-1}}, \\ C(T, \alpha) &:= \int_0^{T/4} t^{2-2\alpha} dt + \int_{T/4}^{3T/4} \left[ t^{1-\alpha} - \left( t - \frac{T}{4} \right)^{1-\alpha} \right]^2 dt \\ &\quad + \int_{3T/4}^T \left[ t^{1-\alpha} - \left( t - \frac{T}{4} \right)^{1-\alpha} - \left( t - \frac{3T}{4} \right)^{1-\alpha} \right]^2 dt, \end{aligned}$$

and

$$B^0 := \limsup_{\xi \rightarrow 0^+} \frac{\int_{T/4}^{3T/4} F(t, \xi) dt}{\xi^2},$$

where  $F$  is the potential of  $f$  defined by

$$F(t, \xi) := \int_0^\xi f(t, x) dx, \quad (t, \xi) \in [0, T] \times \mathbb{R}.$$

With the above notation, in [2, Theorem 3.2], exploiting a quoted critical point theorem established by Ricceri in [17], the following result has been shown.

**Theorem 1.1.** *Let  $f : [0, T] \times \mathbb{R} \rightarrow \mathbb{R}$  be a continuous function such that*

(f<sub>1</sub>)  $F(t, \xi) \geq 0$  for every  $(t, \xi) \in ([0, \frac{T}{4}] \cup [\frac{3T}{4}, T]) \times \mathbb{R}$ .

*Assume that there exist two real sequences  $\{c_n\}$  and  $\{d_n\}$  in  $[0, +\infty)$ , with  $\lim_{n \rightarrow \infty} d_n = 0$ , satisfying the conditions:*

(h<sub>3</sub>) *for some  $n_0 \in \mathbb{N}$  one has*

$$c_n < \frac{T |\cos(\pi\alpha)| \Gamma(2-\alpha)}{4\kappa \sqrt{C(T, \alpha)}} d_n$$

*for each  $n \geq n_0$ ,*

$$(h_4) \mathcal{A}_0 := \lim_{n \rightarrow \infty} \varphi(c_n, d_n, \alpha, T) < \frac{B^0}{16\kappa^2 C(T, \alpha)}, \text{ where}$$

$$\varphi(c_n, d_n, \alpha, T) := \frac{\int_0^T \max_{|\xi| \leq d_n} F(t, \xi) dt - \int_{T/4}^{3T/4} F(t, c_n) dt}{T^2 |\cos(\pi\alpha)|^2 \Gamma^2(2 - \alpha) d_n^2 - 16\kappa^2 c_n^2 C(T, \alpha)}.$$

Then, for each

$$\lambda \in \left( \frac{16C(T, \alpha)}{T^2 \Gamma^2(2 - \alpha) |\cos(\pi\alpha)| B^0}, \frac{1}{\kappa^2 T^2 \Gamma^2(2 - \alpha) |\cos(\pi\alpha)| \mathcal{A}_0} \right),$$

problem

$$\begin{cases} \Delta_{F, \alpha} u(t) + \lambda f(t, u(t)) = 0 & \text{a.e. } t \in [0, T], \\ u(0) = u(T) = 0, \end{cases}$$

admits a sequence of non-zero solutions which strongly converges to zero in  $E_0^\alpha$ .

The aim of this paper is to prove the following remarkable consequence of Theorem 1.1.

**Theorem 1.2.** *Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be a continuous function such that  $f|_{(-\infty, 0]} \equiv 0$  and  $\inf_{\xi \geq 0} F(\xi) = 0$ . Further, let  $h \in C^0([0, T])$  with*

$$(a_0) \min_{t \in [0, T]} h(t) > 0.$$

Suppose that there exist two sequences  $\{c_n\}$  and  $\{d_n\}$  in  $(0, +\infty)$ , with  $c_n < d_n$  for every  $n \geq \nu$ , and  $\lim_{n \rightarrow \infty} d_n = 0$ , such that:

$$(a_1) \lim_{n \rightarrow \infty} \frac{d_n}{c_n} = +\infty,$$

$$(a_2) \max_{x \in [c_n, d_n]} f(x) \leq 0 \text{ for every } n \geq \nu,$$

$$(a_3) \frac{16C(T, \alpha)}{T^2 \Gamma^2(2 - \alpha) |\cos(\pi\alpha)| \int_{\frac{T}{4}}^{\frac{3T}{4}} h(t) dt} < \limsup_{\xi \rightarrow 0^+} \frac{F(\xi)}{\xi^2} < +\infty.$$

Then, the following problem

$$\begin{cases} \Delta_{F, \alpha} u(t) + h(t)f(u(t)) = 0 & \text{a.e. } t \in [0, T], \\ u(0) = u(T) = 0 \end{cases}$$

admits a sequence of non-zero solutions which strongly converges to zero in  $E_0^\alpha$ .

For several results on fractional differential equations, one can see, for example, the monographs of Miller and Ross [15], Samko *et al.* [18], Podlubny [16], Hilfer [11], Kilbas *et al.* [13] and the papers [1, 3–7]

We cite a recent monograph by Kristály, Rădulescu and Varga [14] as a general reference on variational methods adopted here.

2. PROOF OF THE MAIN RESULT

Our aim is to apply Theorem 1.1. First of all observe that, by (a<sub>0</sub>), condition (f<sub>1</sub>) holds. Further, if {c<sub>n</sub>} and {d<sub>n</sub>} are two real sequences satisfying our assumptions, we have that there exists n<sub>0</sub> ≥ ν such that

$$\frac{c_n^2}{d_n^2} < \frac{T|\cos(\pi\alpha)|\Gamma(2-\alpha)}{4\kappa\sqrt{C(T,\alpha)}},$$

for every n ≥ n<sub>0</sub>. Hence the hypothesis (h<sub>3</sub>) in Theorem 1.1 is verified. We will prove that

$$\mathcal{A}_0 := \lim_{n \rightarrow \infty} \frac{\|h\|_{L^1([0,T])} \max_{|\xi| \leq d_n} F(\xi) - \left( \int_{\frac{T}{4}}^{\frac{3T}{4}} h(t) dt \right) F(c_n)}{T^2|\cos(\pi\alpha)|^2\Gamma^2(2-\alpha)d_n^2 - 16\kappa^2c_n^2C(T,\alpha)} = 0.$$

Set

$$h_n := \|h\|_{L^1([0,T])} \frac{\max_{|\xi| \leq d_n} F(\xi)}{c_n^2} - \left( \int_{\frac{T}{4}}^{\frac{3T}{4}} h(t) dt \right) \frac{F(c_n)}{c_n^2}$$

for every n ≥ n<sub>0</sub> and observe that hypothesis (a<sub>2</sub>) yields

$$\max_{|\xi| \leq d_n} F(\xi) = \max_{|\xi| \leq c_n} F(\xi). \tag{2.1}$$

Thus, since

$$\frac{\int_{\frac{T}{4}}^{\frac{3T}{4}} h(t) dt}{\|h\|_{L^1([0,T])}} \leq 1 \quad \text{and} \quad F(c_n) \geq 0,$$

by (2.1), we can write

$$\frac{\max_{|\xi| \leq d_n} F(\xi)}{c_n^2} = \frac{\max_{|\xi| \leq c_n} F(\xi)}{c_n^2} \geq \frac{F(c_n)}{c_n^2} \geq \frac{\int_{\frac{T}{4}}^{\frac{3T}{4}} h(t) dt}{\|h\|_{L^1([0,T])}} \frac{F(c_n)}{c_n^2}.$$

for every n ≥ n<sub>0</sub>.

Since h<sub>n</sub> ≥ 0 for every n ≥ n<sub>0</sub>, one easily gets

$$0 \leq \limsup_{n \rightarrow \infty} h_n.$$

Further, by (a<sub>3</sub>) we have

$$0 < \limsup_{\xi \rightarrow 0^+} \frac{F(\xi)}{\xi^2} < +\infty, \tag{2.2}$$

and consequently (note that  $c_n \searrow 0^+$  as  $n \rightarrow \infty$ ) we obtain

$$0 \leq \limsup_{n \rightarrow \infty} \frac{F(c_n)}{c_n^2} < +\infty. \tag{2.3}$$

Now, let  $\xi_n \in (0, c_n]$  be a sequence such that  $F(\xi_n) := \max_{|\xi| \leq c_n} F(\xi)$  for every  $n \geq n_0$ .

Thus

$$\limsup_{n \rightarrow \infty} \frac{\max_{|\xi| \leq d_n} F(\xi)}{c_n^2} = \limsup_{n \rightarrow \infty} \frac{\max_{|\xi| \leq c_n} F(\xi)}{c_n^2} = \limsup_{n \rightarrow \infty} \frac{F(\xi_n)}{c_n^2} \leq \limsup_{n \rightarrow \infty} \frac{F(\xi_n)}{\xi_n^2}.$$

The above inequalities and (2.2) yield

$$0 \leq \limsup_{n \rightarrow \infty} \frac{\max_{|\xi| \leq d_n} F(\xi)}{c_n^2} \leq \limsup_{n \rightarrow \infty} \frac{F(\xi_n)}{\xi_n^2} < +\infty.$$

Hence, there exists a constant  $\beta$  such that

$$0 \leq \limsup_{n \rightarrow \infty} h_n = \beta. \tag{2.4}$$

Then, by (a<sub>1</sub>) and (2.4), one has

$$\mathcal{A}_0 = \limsup_{n \rightarrow \infty} \frac{h_n}{\left( T^2 |\cos(\pi\alpha)|^2 \Gamma^2(2 - \alpha) \frac{d_n^2}{c_n^2} - 16\kappa^2 C(T, \alpha) \right)} = 0.$$

Concluding, hypothesis (h<sub>4</sub>) holds. Finally, bearing in mind condition (a<sub>3</sub>), one has

$$1 \in \left( \frac{16C(T, \alpha)}{T^2 \Gamma^2(2 - \alpha) |\cos(\pi\alpha)| B^0}, +\infty \right).$$

Thanks to Theorem 1.1, the thesis is achieved. The next result is a direct consequence of Theorem 1.2.

**Proposition 2.1.** *Let  $h \in C^0([0, T])$  satisfying condition (a<sub>0</sub>). Also let  $\{c_n\}$  and  $\{d_n\}$  be two sequences in  $(0, +\infty)$  such that  $d_{n+1} < c_n < d_n$  for every  $n \geq \nu$ ,  $\lim_{n \rightarrow \infty} d_n = 0$ , and  $\lim_{n \rightarrow \infty} \frac{d_n}{c_n} = +\infty$ . Moreover, let  $\varphi \in C^1([0, 1])$  be a nonnegative function such that  $\varphi(0) = \varphi(1) = \varphi'(0) = \varphi'(1) = 0$  and*

$$\max_{s \in [0, 1]} \varphi(s) > \frac{16C(T, \alpha)}{T^2 \Gamma^2(2 - \alpha) |\cos(\pi\alpha)| \int_{\frac{T}{4}}^{\frac{3T}{4}} h(t) dt}.$$

Further, let  $g : \mathbb{R} \rightarrow \mathbb{R}$  be the function defined by

$$g(t) := \begin{cases} \varphi\left(\frac{t - d_{n+1}}{c_n - d_{n+1}}\right) & \text{if } t \in \bigcup_{n \geq \nu} [d_{n+1}, c_n], \\ 0 & \text{otherwise.} \end{cases}$$

Then, problem

$$\begin{cases} \Delta_{F,\alpha} u(t) + h(t)y(u(t)) = 0 & \text{a.e. } t \in [0, T], \\ u(0) = u(T) = 0, \end{cases}$$

where

$$y(u(t)) := |u(t)|(2g(u(t)) + ug'(u(t))),$$

admits a sequence of non-zero solutions which strongly converge to zero in  $E_0^\alpha$ .

*Proof.* Let  $\{c_n\}$  and  $\{d_n\}$  be two positive sequences satisfying our assumptions. We claim that all the hypotheses of Theorem 1.2 are verified. Indeed, one has

$$F(\xi) := \int_0^\xi y(t)dt = \xi^2 g(\xi) \quad \text{for all } \xi \in \mathbb{R}^+.$$

Moreover, direct computations ensure that

$$\max_{x \in [c_{n+1}, d_{n+1}]} y(x) = 0$$

for every  $n \geq \nu$ , and

$$\limsup_{\xi \rightarrow 0^+} \frac{F(\xi)}{\xi^2} = \max_{s \in [0,1]} \varphi(s) > \frac{16C(T, \alpha)}{T^2 \Gamma^2(2 - \alpha) |\cos(\pi\alpha)| \int_{\frac{T}{4}}^{\frac{3T}{4}} h(t) dt}.$$

The assertion follows by Theorem 1.2. □

In conclusion we present a concrete example of the application of Proposition 2.1.

**Example 2.2.** Let  $h \in C^0([0, T])$  satisfying condition  $(a_0)$ . Take the positive real sequences

$$a_n := \frac{1}{n!n} \quad \text{and} \quad b_n := \frac{1}{n!}$$

for every  $n \geq 2$ . Now, define  $\varphi \in C^1([0, 1])$  as follows

$$\varphi(s) := \zeta e^{\frac{1}{s(s-1)} + 4} \quad (\text{for all } s \in [0, 1]),$$

and set

$$\widehat{g}(t) := \begin{cases} \varphi\left(\frac{t - 1/(n+1)!}{1/(n!n) - 1/(n+1)!}\right) & \text{if } t \in A, \\ 0 & \text{otherwise,} \end{cases}$$

where

$$A := \bigcup_{n \geq 2} \left[ \frac{1}{(n+1)!}, \frac{1}{(n!n)} \right].$$

If

$$\zeta > \frac{16C(T, \alpha)}{T^2 \Gamma^2(2 - \alpha) |\cos(\pi\alpha)| \int_{\frac{T}{4}}^{\frac{3T}{4}} h(t) dt},$$

then problem

$$\begin{cases} \Delta_{F, \alpha} u(t) + h(t)y(u(t)) = 0 & \text{a.e. } t \in [0, T], \\ u(0) = u(T) = 0, \end{cases}$$

where

$$y(u(t)) := |u(t)|(2\widehat{g}(u(t)) + u\widehat{g}'(u(t))),$$

admits a sequence of non-zero solutions which strongly converges to zero in  $E_0^\alpha$ .

We just mention, for completeness, that related variational arguments have been used recently in [8] proving the existence of at least one non-zero solution for one dimensional fractional equations. See also [9, 10] for related topics.

### Acknowledgments

*The paper is realized with the auspices of the INdAM – GNAMPA Project 2014 titled: Proprietà geometriche ed analitiche per problemi non-locali.*

### REFERENCES

- [1] R.P. Agarwal, M. Benchohra, S. Hamani, *A survey on existence results for boundary value problems of nonlinear fractional differential equations and inclusions*, Acta Appl. Math. **109** (2010), 973–1033.
- [2] G.A. Afrouzi, A. Hadjian, G. Molica Bisci, *Some results for one dimensional fractional problems* (submitted).
- [3] B. Ahmad, J.J. Nieto, *Existence results for a coupled system of nonlinear fractional differential equations with three-point boundary conditions*, Comput. Math. Appl. **58** (2009), 1838–1843.
- [4] C. Bai, *Impulsive periodic boundary value problems for fractional differential equation involving Riemann-Liouville sequential fractional derivative*, J. Math. Anal. Appl. **384** (2011), 211–231.
- [5] C. Bai, *Solvability of multi-point boundary value problem of nonlinear impulsive fractional differential equation at resonance*, Electron. J. Qual. Theory Differ. Equ. **89** (2011), 1–19.
- [6] Z. Bai, H. Lu, *Positive solutions for boundary value problem of nonlinear fractional differential equation*, J. Math. Anal. Appl. **311** (2005), 495–505.
- [7] M. Benchohra, S. Hamani, S.K. Ntouyas, *Boundary value problems for differential equations with fractional order and nonlocal conditions*, Nonlinear Anal. **71** (2009), 2391–2396.

- [8] M. Galewski, G. Molica Bisci, *Existence results for one-dimensional fractional equations* (submitted).
- [9] S. Heidarkhani, *Multiple solutions for a nonlinear perturbed fractional boundary value problem*, *Dynamic Systems and Applications* (to appear).
- [10] S. Heidarkhani, *Infinitely many solutions for nonlinear perturbed fractional boundary value problems*, preprint.
- [11] R. Hilfer, *Applications of Fractional Calculus in Physics*, World Scientific, Singapore, 2000.
- [12] F. Jiao, Y. Zhou, *Existence of solutions for a class of fractional boundary value problems via critical point theory*, *Comput. Math. Appl.* **62** (2011), 1181–1199.
- [13] A.A. Kilbas, H.M. Srivastava, J.J. Trujillo, *Theory and Applications of Fractional Differential Equations*, Elsevier, Amsterdam, 2006.
- [14] A. Kristály, V. Rădulescu, Cs. Varga, *Variational Principles in Mathematical Physics, Geometry, and Economics: Qualitative Analysis of Nonlinear Equations and Unilateral Problems*, *Encyclopedia of Mathematics and its Applications*, No. 136, Cambridge University Press, Cambridge, 2010.
- [15] K.S. Miller, B. Ross, *An Introduction to the Fractional Calculus and Fractional Differential Equations*, Wiley, New York, 1993.
- [16] I. Podlubny, *Fractional Differential Equations*, Academic Press, San Diego, 1999.
- [17] B. Ricceri, *A general variational principle and some of its applications*, *J. Comput. Appl. Math.* **113** (2000), 401–410.
- [18] S.G. Samko, A.A. Kilbas, O.I. Marichev, *Fractional Integral and Derivatives: Theory and Applications*, Gordon and Breach, Longhorne, PA, 1993.

Massimiliano Ferrara  
massimiliano.ferrara@unirc.it

University of Reggio Calabria  
and CRIOS University Bocconi of Milan  
Via dei Bianchi presso Palazzo Zani  
89127 Reggio Calabria, Italy

Giovanni Molica Bisci  
gmolica@unirc.it

Dipartimento P.A.U.  
Università degli Studi Mediterranea di Reggio Calabria  
Salita Melissari - Feo di Vito  
89124 Reggio Calabria, Italy

*Received: February 3, 2014.*  
*Revised: February 27, 2014.*  
*Accepted: February 28, 2014.*