

## MATRIX POLYNOMIALS ORTHOGONAL WITH RESPECT TO A NON-SYMMETRIC MATRIX OF MEASURES

Marcin J. Zygmunt

*Communicated by P.A. Cojuhari*

**Abstract.** The paper focuses on matrix-valued polynomials satisfying a three-term recurrence relation with constant matrix coefficients. It is shown that they form an orthogonal system with respect to a matrix of measures, not necessarily symmetric. Moreover, it is stated the condition on the coefficients of the recurrence formula for which the matrix measure is symmetric.

**Keywords:** matrix orthogonal polynomials, recurrence formula, matrix of measures, block Jacobi matrices.

**Mathematics Subject Classification:** 47B36, 15A57, 39B42, 42C05.

### 1. INTRODUCTION

The theory of matrix orthogonal polynomials (which will be denoted as MOP) began with the works of Berezanski [1] and Krein [9]. The turn of millennia saw a flurry of articles, among them for instance [2, 4–7, 11, 13, 14]. They consider only positive definite matrices of measures of orthogonality. MOPs – which are known to satisfy a three-term recurrence relation of type (1.3) – are investigated for example in the theory of random walks on graphs or in the theory of multiple birth-and-death processes (cf. [5, 8]), where such three term formulae appear. Many properties of random walk or birth-and-death processes can be obtained from the matrix of measures of orthogonality. Investigation of more general classes of such processes require application of matrices of measures which are non symmetric, see [15]. Theorem 1.2 shows conditions on coefficients in (1.3) to provide the positive definiteness.

The case of non symmetric matrices of measure of orthogonality was discussed for example already by Naimark [10], although not in an explicit form. But still there is very little on the subject in the literature. In this paper we show that for constant terms

in a three-term recurrence relation of type (1.3) there exists a matrix of measures, which are the measures of orthogonality for polynomials generated by (1.3).

Let  $\{P_n\}$  be a family of matrix-valued polynomials such that  $\deg P_n = n$  and the leading coefficient of  $P_n$  is invertible. In the common sense the family  $\{P_n\}$  is said to be orthogonal if there is a matrix of measures  $\Sigma$  supported on a bounded set  $\Delta$  such that the integral

$$\int_{\Delta} P_n(x)\Sigma(dx)P_m(x)^* = 0 \quad (1.1)$$

for  $n \neq m$ . In the literature on the subject – as it was said above – they consider mostly positive definite matrices of measures of orthogonality. In [6] was shown that in this case the generalization of Favard's theorem holds which can be stated as follows (cf. [6, Theorem 3.1]).

**Theorem 1.1.** *Let matrix valued polynomials  $P_n$  satisfy the three-term recurrence formula*

$$xP_n(x) = A_{n+1}P_{n+1}(x) + B_nP_n(x) + A_n^*P_{n-1}(x), \quad (1.2)$$

where matrices  $A_n$  are invertible and  $B_n$  self-adjoint. Then there exists a positive definite matrix of measures  $\Sigma$  satisfying (1.1). Moreover, those polynomials are orthonormal, i.e. the following equality holds

$$\int_{\mathbb{R}} P_n(x)\Sigma(dx)P_n(x)^* = I,$$

where  $I$  stands for the identity matrix (note that in this case the orthogonality matrix measure is supported on the real line  $\mathbb{R}$ ).

If the normalization is not required, polynomials  $P_n$  fulfill more general recurrence formula (cf. [4, 5, 9]):

$$xP_n(x) = A_nP_{n+1}(x) + B_nP_n(x) + C_nP_{n-1}(x),$$

where all matrices  $A_n$  and  $C_n$  are invertible. But not for all sets of matrices  $A_n, B_n, C_n$  there exists a positive definite matrix of measures which satisfy the orthogonality relation (1.1). In fact the following theorem holds to be true (the proof is given in the next section).

**Theorem 1.2.** *Let  $P_n$  be the family of matrix polynomials satisfying the recurrence relation*

$$xP_n(x) = A_nP_{n+1}(x) + B_nP_n(x) + C_nP_{n-1}(x), \quad (1.3)$$

with  $P_0(x) = I$  and  $P_{-1}(x) = 0$ , where all matrices  $A_n$  and  $C_n$  are invertible. Then there exists a positive definite matrix of measure, with respect to which the polynomials  $P_n$  are orthogonal, if and only if matrices  $A_0 \cdots A_n C_{n+1} \cdots C_1$  and  $A_0 \cdots A_{n-1} B_n C_n \cdots C_1$  are self-adjoint for every  $n$ .

In this paper we will use another way of defining orthogonality, which is of course equivalent in the scalar case: we will say that polynomials  $P_n$  are orthogonal with respect to the matrix of measures  $\Sigma$  if

$$\int_{\Delta} x^k P_n(x) \Sigma(dx) = 0 \tag{1.4}$$

for all  $n > k \geq 0$ . The condition above means that polynomial  $P_n$  is orthogonal to the subspace of all polynomials of degree less than or equal to  $n - 1$ . The reader can verify that both ways to define orthogonality used for example in such textbooks on classical orthogonal polynomials as [3] and [12].

In the matrix case the family of matrix valued polynomials, given by the Gramm–Schmidt procedure from the system of monomials  $\{I, x, x^2, \dots\}$  orthogonalized with respect to a positive definite matrix of measures, are not only orthogonal to the subspace of polynomials of lower degree, but such polynomials are orthogonal to each other, i.e. they satisfy (1.1). This is due to the fact that the matrix valued bilinear form in this case is symmetric in the sense that

$$\left( \int_{\mathbb{R}} Q(x) \Sigma(dx) R(x) \right)^* = \int_{\mathbb{R}} R(x)^* \Sigma(dx) Q(x)^*, \tag{1.5}$$

for any polynomials  $Q$  and  $R$ .

In the whole generality the matrix of measures does not need to be symmetric, so (1.5) does not hold any more. What we can have is only (1.1) for  $n > m$ . Indeed, if (1.4) holds, then (1.1) for  $n > m$  holds too, as every polynomial  $P_m^*$  has degree lower than  $n$ . And vice-versa, every monomial  $x^m$  is a linear combination of polynomials  $P_0^*, P_1^*, \dots, P_m^*$ , hence (1.4) holds if only (1.1) holds for  $n > m$ . So in their full generality the formulas (1.1) and (1.4) do not need to be equivalent.

## 2. PROOF OF THE THEOREM 1.2

*Proof of the Theorem 1.2.* Let  $P_n$  be MOP, orthogonal with respect to a positive definite matrix of measures  $\Sigma$  and let them satisfy the recurrence formula (1.3). The condition that all coefficients  $A_n$  and  $C_n$  are invertible provides the fact that integral  $\int Q(x) \Sigma(dx) Q(x)^*$  is a positive (strictly) definite matrix for all non zero polynomials  $Q$ , which do not have a non trivial constant null subspace (cf. §2 of [4], in particular Lemma 2.1).

In this case the corresponding orthonormal matrix polynomials (which we denote here by  $P_n^o$ ) satisfy  $P_n(x) = V_n P_n^o(x)$ , where

$$V_n V_n^* = \left( \int_{\Delta} P_n(x) \Sigma(dx) P_n(x)^* \right)^{-1}.$$

If we require  $P_0 = I$ , then we need the condition  $V_0 = I$ . But if  $P_0 \neq I$ , then we can consider polynomials  $\hat{P}_n^o = P_n^o P_0$  which are orthonormal with respect to the measure  $P_0^{-1} \Sigma P_0^{*-1}$ . Hence we can assume that  $V_0 = I$ .

Orthonormal polynomials  $P_n^o$  fulfill the recurrence formula

$$xP_n^o(x) = A_n^o P_{n+1}^o(x) + B_n^o P_n^o(x) + (A_{n-1}^o)^* P_{n-1}^o(x), \text{ with } P_0^o(x) = I, P_{-1}^o(x) = 0,$$

with suitable coefficients  $A_n^o$  and  $B_n^o$ . Putting  $V_n P_n^o(x)$  into (1.2) leads to the formula

$$xP_n^o(x) = V_n^{-1} A_n V_{n+1} P_{n+1}^o(x) + V_n^{-1} B_n V_n P_n^o(x) + V_n^{-1} C_n V_{n-1} P_{n-1}^o(x),$$

so

$$A_n^o = V_{n-1}^{-1} A_{n-1} V_n = (V_n^{-1} C_n V_{n-1})^*$$

and

$$B_n^o = V_n^{-1} B_n V_n = (V_n^{-1} B_n V_n)^*. \quad (2.1)$$

Hence

$$V_n V_n^* = A_{n-1}^{-1} V_{n-1} V_{n-1}^* C_n^* = \dots = A_{n-1}^{-1} \dots A_0^{-1} C_1^* \dots C_n^*.$$

But the left-hand side is a hermitian matrix, so the right-hand side is hermitian as well. Thus

$$A_{n-1}^{-1} \dots A_0^{-1} C_1^* \dots C_n^* = C_n \dots C_1 (A_0^{-1})^* \dots (A_{n-1}^{-1})^*,$$

which leads to

$$A_0 \dots A_{n-1} C_n \dots C_1 = (A_0 \dots A_{n-1} C_n \dots C_1)^*.$$

Now (2.1) gives

$$B_n (V_n V_n^*) = (V_n V_n^*) B_n^*.$$

Combining it with the previous one shows that

$$A_0 \dots A_{n-1} B_n C_n \dots C_1 = (A_0 \dots A_{n-1} B_n C_n \dots C_1)^*.$$

To prove the implication in the opposite direction assume now that the matrices  $A_n$ ,  $B_n$  and  $C_n$  fulfill the conditions in the thesis, i.e.

$$A_0 \dots A_{n-1} C_n \dots C_1 = (A_0 \dots A_{n-1} C_n \dots C_1)^* = C_n^* \dots C_1^* A_{n-1}^* \dots A_0^*$$

and

$$A_0 \dots A_{n-1} B_n C_n \dots C_1 = (A_0 \dots A_{n-1} B_n C_n \dots C_1)^* = C_n^* \dots C_1^* B_n^* A_{n-1}^* \dots A_0^*.$$

Then

$$C_n \dots C_1 (A_0^{-1})^* \dots (A_{n-1}^{-1})^* = A_{n-1}^{-1} \dots A_0^{-1} C_1^* \dots C_n^*$$

and

$$B_n C_n \cdots C_1 (A_0^{-1})^* \cdots (A_{n-1}^{-1})^* = A_{n-1}^{-1} \cdots A_0^{-1} C_n^* \cdots C_1^* B_n^*,$$

which shows that the left-hand sides of both equations are self-adjoint and invertible matrices. Hence the first one can be rewritten as a product  $W_n W_n^*$ , where  $W_n$  is an invertible matrix satisfying  $W_n W_n^* = A_{n-1}^{-1} W_{n-1} W_{n-1}^* C_n^*$  and the other as  $B_n (W_n W_n^*) = (W_n W_n^*) B_n^*$ . This shows that the polynomials  $W_n^{-1} P_n$  fulfill the recurrence formula (1.2). By Theorem 1.1, polynomials  $P_n$  are orthonormal with respect to a positive definite matrix of measures. So polynomials  $P_n$  are also orthogonal with respect to the same matrix-valued measure.  $\square$

Now a question arises: In a more general case, is there any matrix-valued measure  $\Sigma$  supported on an appropriate set  $\Delta$  such that

$$\int_{\Delta} x^m P_n(x) \Sigma(dx) = 0$$

for  $n > m \geq 0$ ?

### 3. GENERAL CASE OF CONSTANT COEFFICIENTS

#### 3.1. MATRIX OF MEASURES

Let  $P_n$  be polynomials satisfying the recurrence formula of type (1.3) with constant coefficients, i.e.

$$xP_n(x) = AP_{n+1}(x) + BP_n(x) + CP_{n-1}(x) \tag{3.1}$$

with  $P_0 = I$  and  $P_1(x) = A^{-1}(x - B)$ , where  $A$  and  $C$  are invertible. Hence  $\deg P_n = n$ . We will extend the equality (3.1) to all  $n \geq 0$  assuming that  $P_{-1}(x) = 0$ , i.e., a zero matrix.

In the case when  $C = A^*$  and  $B$  is a symmetric matrix, we already know that polynomials  $P_n$  are MOP with respect to a positive definite matrix of measures supported on the real line  $\mathbb{R}$ . In the following we will show that there exists a matrix of measures – of course not necessarily symmetric or with its support outside real line  $\mathbb{R}$  – which is an orthogonality measure for  $P_n$  in the sense of (1.4).

Let us now introduce new polynomials  $R_n$  which satisfy

$$xR_n(x) = R_{n+1}(x)C + R_n(x)B + R_{n-1}(x)A \tag{3.2}$$

for  $n \geq 1$ , where  $R_0(x) = I$  and  $R_{-1}(x) = 0$ . One can put the initial condition  $R_1(x) = (x - B)C^{-1}$  instead of  $R_{-1}$ .

**Lemma 3.1.** *Let matrix valued polynomials  $P_n$  and  $R_n$  satisfy (3.1) and (3.2), respectively. Then we have*

$$\begin{aligned} &P_n(z)A^{-1}R_{m+1}(z) - P_{n+1}(z)C^{-1}R_m(z) \\ &= P_{n+k}(z)A^{-1}R_{m+k+1}(z) - P_{n+k+1}(z)C^{-1}R_{m+k}(z) \end{aligned} \tag{3.3}$$

for any  $m, n, k \in \mathbb{N}$ .

*Proof.* Let

$$\mathcal{X}(x) = A^{-1}(x - B), \quad \mathcal{D} = A^{-1}C,$$

and

$$\hat{\mathcal{X}}(x) = (x - B)C^{-1}, \quad \hat{\mathcal{D}} = AC^{-1}.$$

The formulas (3.1) and (3.2) become as follows:

$$P_{n+1}(x) = \mathcal{X}(x)P_n(x) - \mathcal{D}P_{n-1}(x), \quad R_{n+1}(x) = R_n(x)\hat{\mathcal{X}}(x) - R_{n-1}(x)\hat{\mathcal{D}}.$$

But we have also

$$\begin{aligned} P_1(x) &= \mathcal{X}(x) = P_0(x)\mathcal{X}(x), \\ P_2(x) &= \mathcal{X}(x)^2 - \mathcal{D} = P_1(x)\mathcal{X}(x)^2 - P_0(x)\mathcal{D} \end{aligned}$$

and

$$\begin{aligned} R_1(x) &= \hat{\mathcal{X}}(x) = \hat{\mathcal{X}}(x)R_0(x), \\ R_2(x) &= \hat{\mathcal{X}}(x)^2 - \hat{\mathcal{D}} = \hat{\mathcal{X}}(x)R_1(x) - \hat{\mathcal{D}}R_0(x). \end{aligned}$$

By simply induction on the degree of polynomials one can see that

$$P_{n+1}(x) = P_n(x)\mathcal{X}(x) - P_{n-1}(x)\mathcal{D} \quad \text{and} \quad R_{n+1}(x) = \hat{\mathcal{X}}(x)R_n(x) - \hat{\mathcal{D}}R_{n-1}(x). \quad (3.4)$$

The equality (3.3) is a matter of applying (3.4), and simple calculations. □

Let us fix  $n$  for a while and let  $\Sigma_{(n)}$  be a matrix of measures supported on zeros of  $P_n$  (denoted by  $\Delta_n = \{\xi \in \mathbb{C} : \det(P_n(\xi)) = 0\}$ ) such that

$$\int_{\Delta_n} \frac{1}{z - \xi} \Sigma_{(n)}(d\xi) = P_n(z)^{-1}P_{n-1}(z)A^{-1}.$$

This means that

$$\Sigma_{(n)} = \sum_{\xi \in \Delta_n} W_{\xi}^{(n)} \delta_{\{\xi\}},$$

where  $\delta_{\{\xi\}}$  is a well-known Dirac measure at  $\xi$ . The matrix coefficients  $W_{\xi}^{(n)}$  can be calculated for instance from the Cauchy integral formula

$$W_{\xi_o}^{(n)} = \frac{1}{2\pi i} \int_{\Gamma_{\xi_o}} P_n(\zeta)^{-1}P_{n-1}(\zeta)A^{-1} d\zeta$$

provided that  $\Gamma_{\xi_o}$  is a simple closed curve enclosing  $\xi_o$  and leaving outside its contour other zeroes of  $P_n$  (note that  $\xi_o$  does not need to be a real number, as Examples 4.3 and 4.4 show). Indeed, as

$$P_n(z)^{-1}P_{n-1}(z)A^{-1} = \sum_{\xi \in \Delta_n} \frac{1}{z - \xi} W_{\xi}^{(n)},$$

we get

$$\begin{aligned} \frac{1}{2\pi i} \int_{\Gamma_{\xi_o}} P_n(\zeta)^{-1} P_{n-1}(\zeta) A^{-1} d\zeta &= \sum_{\xi \in \Delta_n} \frac{1}{2\pi i} \int_{\Gamma_{\xi_o}} \frac{1}{\zeta - \xi} W_{\xi}^{(n)} d\zeta \\ &= W_{\xi_o}^{(n)} \frac{1}{2\pi i} \int_{\Gamma_{\xi_o}} \frac{d\zeta}{\zeta - \xi_o} = W_{\xi_o}^{(n)}. \end{aligned}$$

Let us now denote the moments of  $\Sigma_{(n)}$  as

$$M_k^{(n)} = \int_{\Delta} \xi^k \Sigma_{(n)}(d\xi) = \sum_{\xi \in \Delta_n} \xi^k W_{\xi}^{(n)}$$

and let

$$F_{(n)}(z) = \int_{\Delta} \frac{1}{z - \xi} \Sigma_{(n)}(d\xi) = \sum_{\xi \in \Delta_n} \frac{1}{z - \xi} W_{\xi}^{(n)} = \sum_{k=0}^{+\infty} \frac{z^k}{z^{k+1}} M_k^{(n)}.$$

**Lemma 3.2.** *In the above notation*

$$F_{(n)}(z) = C^{-1} R_{n-1}(z) R_n(z)^{-1}$$

for all  $n \geq 1$ .

*Proof.* From Lemma 3.1 we have

$$P_{n-1}(z) A^{-1} R_n(z) - P_n(z) C^{-1} R_{n-1}(z) = P_0(z) A^{-1} R_1(z) - P_1(z) C^{-1} R_0(z) = 0.$$

Hence

$$P_n(z) F_{(n)}(z) R_n(z) = P_{n-1}(z) A^{-1} R_n(z) = P_n(z) C^{-1} R_{n-1}(z),$$

which leads to the conclusion. □

Let now  $\Delta_{n,\infty} = \overline{\bigcup_{k \geq n} \Delta_k}$ . It is clear that

$$\int_{\Delta_{n,\infty}} \xi^k P_n(\xi) \Sigma_{(n)}(d\xi) = 0.$$

We shall show the following result.

**Theorem 3.3.** *Let  $P_n$  and  $\Sigma_{(n)}$  be as above. Then*

$$\int_{\Delta_{n,\infty}} \xi^k P_m(\xi) \Sigma_{(n)}(d\xi) = 0$$

for all  $0 \leq k < m \leq n$ .

*Proof.* Fix  $n \geq 1$ . Now

$$\begin{aligned} F_{(n+1)}(z) - F_{(n)}(z) &= P_{n+1}(z)^{-1}P_n(z)A^{-1} - C^{-1}R_{n-1}(z)R_n(z)^{-1} \\ &= P_{n+1}(z)^{-1}\left(P_n(z)A^{-1}R_n(z) - P_{n+1}(z)C^{-1}R_{n-1}(z)\right)R_n(z)^{-1} \end{aligned}$$

and by Lemma 3.1

$$= P_{n+1}(z)^{-1}A^{-1}R_n(z)^{-1}.$$

Hence

$$\lim_{z \rightarrow \infty} z^k (F_{(n+1)}(z) - F_{(n)}(z)) = 0$$

for  $0 \leq k \leq 2n$ . This means that  $M_k^{(n+1)} = M_k^{(n)}$  for such  $k$ . Now the thesis of the Theorem follows from a simple induction argument and the fact that  $\Delta_{n+1,\infty} \subset \Delta_{n,\infty}$ .  $\square$

Now we are able to show the following result.

**Theorem 3.4.** *Let  $P_n$  be polynomials satisfying the recurrence formula (3.1). Then there exists a matrix of measures  $\Sigma$  such that*

$$\int_{\Delta} \xi^k P_n(\xi) \Sigma(d\xi) = 0$$

for all integers  $n > k \geq 0$ , where

$$\Delta = \bigcap_{n=1}^{\infty} \Delta_{n,\infty} = \bigcap_{n=1}^{\infty} \overline{\bigcup_{m \geq n} \left\{ \xi \in \mathbb{C} : \det(P_m(\xi)) = 0 \right\}}.$$

*Proof.* Let us fix  $m \in \mathbb{N}$ . Every matrix-valued polynomial  $P$  of degree  $m$  has the unique decomposition

$$P = \sum_{k=0}^m C_{k,P} P_k.$$

Hence, by Theorem 3.3, we have

$$\int_{\Delta_{n,\infty}} P(\xi) \Sigma_{(n)}(d\xi) = C_{0,P}$$

for all  $n > m$ . So the sequence of measures  $\Sigma_{(n)}$  is weakly convergent on the space of all matrix-valued polynomials. Thus, by the Banach–Steinhaus theorem, it has the weak limit  $\Sigma$  such that we have

$$\int_{\Delta_{n,\infty}} P(\xi) \Sigma(d\xi) = C_{0,P}$$



for all  $n > m$ . It is clear that  $\text{supp}\Sigma \subset \Delta_{n,\infty}$  for all  $n \in \mathbb{N}$ , hence

$$\text{supp}\Sigma \subset \bigcap_{n=1}^{\infty} \Delta_{n,\infty}.$$

Now let us fix  $n, k \in \mathbb{N}$ ,  $n > k \geq 0$ . Then

$$\int_{\Delta} \xi^k P_n(\xi) \Sigma(d\xi) = \lim_{m \rightarrow \infty} \int_{\Delta_{m,\infty}} \xi^k P_n(\xi) \Sigma_{(m)}(d\xi) = 0,$$

where the last equality holds by Theorem 3.3. □

### 3.2. CAUCHY TRANSFORM OF THE MATRIX OF MEASURES OF ORTHOGONALITY

Let us now define

$$F(z) = \int_{\Delta} \frac{1}{z - \xi} \Sigma(d\xi).$$

Function  $F$  is often called the *Cauchy Transform* of  $\Sigma$ . Knowing the function  $F$  allows us to find an explicit formula for the orthogonality measure in many cases.

**Theorem 3.5.** *Matrix valued function  $F$  satisfies the equation*

$$F(z) = (z - B - AF(z)C)^{-1}.$$

*Proof.* We have

$$\lim_{n \rightarrow \infty} F_{(n)}(z) = F(z)$$

for all  $z$  in a neighborhood of infinity. By analytic continuation the previous equality holds for all  $z \in \mathbb{C} \setminus \Delta$ .

Now, by (3.4), we have

$$P_n(z)^{-1}P_{n+1}(z) = \mathcal{X}(z) - P_n(z)^{-1}P_{n-1}(z)\mathcal{D},$$

hence

$$F_{(n+1)}(z)^{-1} = z - B - AF_{(n)}(z)C.$$

Taking the limit as  $n$  tends to the infinity leads to the thesis. □

**Corollary 3.6.** *We have*

$$F(z) = \frac{1}{z - B - A \frac{1}{z - B - A \frac{1}{z - B - A \frac{1}{z - B - \dots}} C} C}.$$

## 4. INSTRUCTIVE EXAMPLES

What follows are four examples which show really unexpected behaviour and properties of the orthogonality measure. We will always assume that  $P_0(x) = I$ ,  $P_{-1}(x) = 0$ .

The first two examples are taken from [13].

**Example 4.1.** Let  $P_n$  be polynomials satisfying the recurrence formula

$$xP_n(x) = AP_{n+1}(x) + BP_n(x) + CP_{n-1}(x)$$

with

$$A = C = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \quad \text{and} \quad B = \begin{pmatrix} 0 & \gamma \\ \gamma & 0 \end{pmatrix}.$$

In this case it is not difficult to compute the corresponding Cauchy transform<sup>1)</sup>, i.e.

$$F(z) = \frac{1}{2} \left( 1 - \sqrt{\frac{z^2 - \gamma^2 - 4}{z^2 - \gamma^2}} \right) \begin{pmatrix} z & \gamma \\ \gamma & z \end{pmatrix}.$$

It is shown in [13] that polynomials  $R_n$  are orthogonal with respect to the matrix of measures

$$\Sigma_1(dx) = \frac{1}{2\pi} \sqrt{\frac{\gamma^2 + 4 - x^2}{x^2 - \gamma^2}} \begin{pmatrix} |x| & \gamma \\ \gamma & |x| \end{pmatrix} dx,$$

which is supported on the set

$$\Delta = \left[ -\sqrt{\gamma^2 + 4}, -\gamma \right] \cup \left[ \gamma, \sqrt{\gamma^2 + 4} \right].$$

This result can be achieved also by the Stieltjes-Perron's inversion formula from function  $F$ .

**Example 4.2.** Let  $P_n$  be polynomials satisfying the recurrence formula

$$xP_n(x) = AP_{n+1}(x) + CP_{n-1}(x)$$

with

$$A = \begin{pmatrix} 0 & \beta \\ \alpha & 0 \end{pmatrix} \quad \text{and} \quad C = \begin{pmatrix} 0 & \alpha \\ \beta & 0 \end{pmatrix}, \quad \text{where } \alpha > \beta > 0.$$

In this case the midterm coefficient  $B$  is equal to the zero matrix.

Direct calculation shows that

$$F(z) = \begin{pmatrix} \frac{\alpha^2 - \beta^2}{2z\alpha^2} & 0 \\ 0 & \frac{\beta^2 - \alpha^2}{2z\beta^2} \end{pmatrix} + \frac{z^2 - \sqrt{(z - \alpha - \beta)(z + \alpha - \beta)(z - \alpha + \beta)(z + \alpha + \beta)}}{2z} \begin{pmatrix} \frac{1}{\alpha^2} & 0 \\ 0 & \frac{1}{\beta^2} \end{pmatrix}.$$

<sup>1)</sup> This and other formulas for  $F(z)$  in this section were calculated in *Mathematica* v.10.0, licence L4887-2121

Author in [13] shows that polynomials  $P_n$  are orthogonal with respect to the matrix of measures  $\Sigma = \Sigma_0 + \Sigma_1$ , where

$$\Sigma_0 = \begin{pmatrix} 1 - \left(\frac{\beta}{\alpha}\right)^2 & 0 \\ 0 & 0 \end{pmatrix} \delta_0,$$

where  $\delta_0$  is Dirac measure at 0, and

$$\Sigma_1(dx) = \frac{\sqrt{((\alpha + \beta)^2 - x^2)(x^2 - (\alpha - \beta)^2)}}{4\pi|x|} \begin{pmatrix} \frac{1}{\alpha} & 0 \\ 0 & \frac{1}{\beta} \end{pmatrix} dx.$$

The measure  $\Sigma$  is supported on the set

$$\Delta = [-(\alpha + \beta), -(\alpha - \beta)] \cup \{0\} \cup [\alpha - \beta, \alpha + \beta].$$

Hence, the orthogonality measure can contain a non-empty atomic part.

**Example 4.3.** Let now  $P_n$  be polynomials satisfying the recurrence formula

$$xP_n(x) = AP_{n+1}(x) + CP_{n-1}(x)$$

with

$$A = C = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}.$$

It can be easily seen that multiplying (4.3) on the left-hand side by matrix  $\Lambda$  and on the right-hand side by  $\Lambda^{-1}$ , where

$$\Lambda = \begin{pmatrix} 1 & i \\ i & 1 \end{pmatrix} \quad \text{and} \quad \Lambda^{-1} = \frac{1}{2} \begin{pmatrix} 1 & -i \\ -i & 1 \end{pmatrix},$$

leads to the formula

$$x\tilde{P}_n(x) = \tilde{A}\tilde{P}_{n+1}(x) + \tilde{C}\tilde{P}_{n-1}(x)$$

with

$$\tilde{A} = \tilde{C} = \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix} \quad \text{and} \quad \tilde{P}_n(x) = \Lambda P_n(x) \Lambda^{-1}.$$

Hence  $\tilde{P}_n$  has non-zero terms only on the diagonal, i.e.  $\tilde{P}_n(x) = \begin{pmatrix} \tilde{p}_n^{1,1}(x) & 0 \\ 0 & \tilde{p}_n^{2,2}(x) \end{pmatrix}$ , and those satisfy

$$-\frac{1}{2}ix\tilde{p}_n^{1,1}(x) = \frac{1}{2}\tilde{p}_{n+1}^{1,1}(x) + \frac{1}{2}\tilde{p}_{n-1}^{1,1}(x), \quad \frac{1}{2}ix\tilde{p}_n^{2,2}(x) = \frac{1}{2}\tilde{p}_{n+1}^{2,2}(x) + \frac{1}{2}\tilde{p}_{n-1}^{2,2}(x).$$

So

$$\tilde{p}_n^{1,1}(x) = U_n\left(-\frac{1}{2}ix\right) \quad \text{and} \quad \tilde{p}_n^{2,2}(x) = U_n\left(-\frac{1}{2}ix\right),$$

where scalar polynomials  $U_n(x)$  are Chebyshev polynomials of the second kind and are known to be orthogonal with respect to the measure  $\mu_u(dx) = \frac{2}{\pi}\sqrt{1-x^2} dx$  supported

on the interval  $[-1, 1]$  (ie. polynomials  $U_n(\frac{1}{2}ix)$  are orthogonal with respect to the measure  $\mu_+(dx) = \frac{\sqrt{4+x^2}}{2\pi} dx$  supported on the segment of the line between points  $-2i$  and  $2i$  on complex plane  $\mathbb{C}$ , i.e. the set  $\{z \in \mathbb{C} : z = -2i\lambda + 2i(1 - \lambda), \text{ for } \lambda \in [0, 1]\}$ , which we call later the interval  $[-2i, 2i]$ ). So the polynomials  $P_n$  are equal to

$$P_n(x) = \begin{pmatrix} \frac{1}{2}(U_n(\frac{1}{2}ix) + U_n(-\frac{1}{2}ix)) & -\frac{i}{2}(U_n(\frac{1}{2}ix) - U_n(-\frac{1}{2}ix)) \\ \frac{i}{2}(U_n(\frac{1}{2}ix) - U_n(-\frac{1}{2}ix)) & \frac{1}{2}(U_n(\frac{1}{2}ix) + U_n(-\frac{1}{2}ix)) \end{pmatrix}$$

and they are orthogonal with respect to the matrix of measures  $\Sigma = I\tilde{\mu}$ , where  $\tilde{\mu}(dx) = \frac{1}{2\pi}\sqrt{4+x^2} dx$ . In this case the Cauchy transform equals to

$$F(z) = \frac{z - \sqrt{z^2 + 4}}{2} I.$$

**Example 4.4.** Let  $P_n$  be polynomials satisfying the recurrence formula

$$xP_n(x) = AP_{n+1}(x) + BP_n(x) + CP_{n-1}(x) \tag{4.1}$$

with

$$A = C = I \quad \text{and} \quad B = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}.$$

As in Example 4.3 multiplying (4.1) by matrices  $\Lambda$  and  $\Lambda^{-1}$  leads to

$$x\hat{P}_n(x) = \hat{P}_{n+1}(x) + \hat{B}\hat{P}_n(x) + \hat{P}_{n-1}(x),$$

with  $\hat{B} = \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix}$ . Hence the solutions is

$$\hat{P}_n(x) = \begin{pmatrix} U_n(\frac{x-i}{2}) & 0 \\ 0 & U_n(\frac{x+i}{2}) \end{pmatrix}.$$

This shows that polynomials  $P_n$  are equal to

$$P_n(x) = \begin{pmatrix} \frac{1}{2}(U_n(\frac{x+i}{2}) + U_n(\frac{x-i}{2})) & -\frac{i}{2}(U_n(\frac{x+i}{2}) - U_n(\frac{x-i}{2})) \\ \frac{i}{2}(U_n(\frac{x+i}{2}) - U_n(\frac{x-i}{2})) & \frac{1}{2}(U_n(\frac{x+i}{2}) + U_n(\frac{x-i}{2})) \end{pmatrix}$$

and they are orthogonal with

$$\Sigma = \begin{pmatrix} \frac{1}{2}(\hat{\mu}_+ + \hat{\mu}_-) & \frac{i}{2}(\hat{\mu}_+ - \hat{\mu}_-) \\ -\frac{i}{2}(\hat{\mu}_+ - \hat{\mu}_-) & \frac{1}{2}(\hat{\mu}_+ + \hat{\mu}_-) \end{pmatrix}$$

supported on the set  $\Delta = [-2+i, 2+i] \cup [-2-i, -2+i]$  on complex plane  $\mathbb{C}$ , where

$$\hat{\mu}_+(dx) = \frac{1}{2\pi}\sqrt{5+2ix-x^2} dx \quad \text{and} \quad \hat{\mu}_-(dx) = \frac{1}{2\pi}\sqrt{5-2ix-x^2} dx$$

are supported respectively on intervals  $[-2+i, 2+i]$  and  $[-2-i, -2+i]$ . In this example the Cauchy transform equals to

$$F(z) = \begin{pmatrix} \frac{2z - \sqrt{z^2 + 2iz - 5} - \sqrt{z^2 - 2iz - 5}}{4} & \frac{-2 - i\sqrt{z^2 + 2iz - 5} + i\sqrt{z^2 - 2iz - 5}}{4} \\ \frac{2 + i\sqrt{z^2 + 2iz - 5} - i\sqrt{z^2 - 2iz - 5}}{4} & \frac{2z - \sqrt{z^2 + 2iz - 5} - \sqrt{z^2 - 2iz - 5}}{4} \end{pmatrix}.$$

**Example 4.5.** Let  $P_n$  be polynomials satisfying the recurrence formula

$$xP_n(x) = AP_{n+1}(x) + BP_n(x) + CP_{n-1}(x)$$

with

$$A = C = I \quad \text{and} \quad B = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}.$$

Straightforward computation of the Cauchy transform in this example shows that

$$F(z) = \begin{pmatrix} \frac{z - \sqrt{z^2 - 4}}{2} & \frac{1}{2} \left( \frac{z}{\sqrt{z^2 - 4}} - 1 \right) \\ 0 & \frac{z - \sqrt{z^2 - 4}}{2} \end{pmatrix}.$$

This shows that the solution of the recurrence formula is an upper triangular matrix of polynomials

$$P_n(x) = \begin{pmatrix} p_n^{1,1}(x) & p_n^{1,2}(x) \\ 0 & p_n^{2,2}(x) \end{pmatrix}.$$

Scalar polynomials  $p_n^{1,1}, p_n^{1,2}, p_n^{2,2}$ , satisfy the following recurrences:

$$\begin{aligned} x p_n^{1,1}(x) &= p_{n+1}^{1,1}(x) + p_{n-1}^{1,1}(x) && \text{with } p_0^{1,1}(x) = 1, p_{-1}^{1,1}(x) = 0, \\ x p_n^{1,2}(x) &= p_{n+1}^{1,2}(x) + p_n^{2,2}(x) + p_{n-1}^{1,2}(x) && \text{with } p_0^{1,2}(x) = 0, p_{-1}^{1,2}(x) = 0, \\ x p_n^{2,2}(x) &= p_{n+1}^{2,2}(x) + p_{n-1}^{2,2}(x) && \text{with } p_0^{2,2}(x) = 1, p_{-1}^{2,2}(x) = 0. \end{aligned}$$

This shows that

$$p_n^{1,1}(x) = p_n^{2,2}(x) = U_n\left(\frac{1}{2}x\right).$$

Differentiation of the third equation above gives

$$x \frac{d}{dx} p_n^{2,2}(x) = \frac{d}{dx} p_{n+1}^{2,2}(x) + \frac{d}{dx} p_{n-1}^{2,2}(x) - p_n^{2,2}(x),$$

so

$$p_n^{1,2}(x) = -\frac{d}{dx} p_n^{2,2}(x) = -\frac{1}{2} U'_n\left(\frac{1}{2}x\right).$$

Now simple calculation proves that polynomials  $P_n$  are orthogonal with respect to the matrix of measures

$$\Sigma = \begin{pmatrix} \mu_U & \mu'_U \\ 0 & \mu_U \end{pmatrix},$$

where  $\mu'_U(dx) = \frac{x dx}{2\pi\sqrt{4-x^2}}$  and all measures are supported on the interval  $[-2, 2]$  on the real line  $\mathbb{R}$ . This shows that in this case  $\Sigma$  cannot be transformed neither to a positive definite matrix of measures, nor to a symmetric one.

## REFERENCES

- [1] Yu.M. Berezanski, *Expansions in Eigenfunctions of Selfadjoint Operators*, Translations of Mathematical Monographs 17, American Mathematical Society, R.I., 1968.
- [2] C. Berg, A.J. Duran, *Orthogonal polynomials and analytic functions associated to positive definite matrices*, J. Math. Anal. Appl. **315** (2006), 54–67.
- [3] T. Chihara, *An Introduction to Orthogonal Polynomials*, Mathematics and Its Applications 13, Gordon and Breach, New York, 1978.
- [4] D. Damanik, A. Pushnitski, B. Simon, *The analytic theory of matrix orthogonal polynomials*, Surveys in Approx. Theory **4** (2008), 1–85.
- [5] H. Dette, B. Reuther, W.J. Studden, M.J. Zygmunt, *Matrices measures and random walks with a block tridiagonal transition matrix*, SIAM J. Matrix Anal. Appl. **29** (2006) 1, 117–142.
- [6] A.J. Duran, *A generalization of Favard’s theorem for polynomials satisfying a recurrence relation*, J. Approx. Theory **74** (1993), 83–109.
- [7] A.J. Duran, *On orthogonal polynomials with respect to a positive definite matrix of measures*, Canad. J. Math. **47** (1995), 88–112.
- [8] F.A. Grünbaum, M.D. de la Iglesia, *Matrix valued orthogonal polynomials arising from group representation theory and a family of quasi-birth-and-death processes*, SIAM J. Matrix Anal. Applic. **30** (2008) 2, 741–761.
- [9] M.G. Krein, *Infinite  $J$ -matrices and a matrix moment problem*, Dokl. Akad. Nauk SSSR **69** (1949), 125–128 [in Russian].
- [10] M.A. Naimark, *Investigation of the spectrum and the expansion in eigenfunctions of a non-selfadjoint operator of second order on a semi-axis*, Trudy Mosk. Mat. Obs. **3** (1954), 181–270 [in Russian].
- [11] A. Sinap, W. Van Assche, *Orthogonal matrix polynomials and applications*, J. Comput. Appl. Math. **66** (1996), 27–52.
- [12] G. Szegő, *Orthogonal Polynomials*, AMS Coll. Pub., vol. 23, AMS, Providence, 1975 (4th edition).
- [13] M.J. Zygmunt, *Matrix orthogonal polynomials and continued fractions*, Linear Alg. Appl. **340**, (2002) 1–3, 155–168.
- [14] M.J. Zygmunt, *Jacobi block matrices with constant matrix terms*, Oper. Th.: Adv. & Appl. **154** (2004), 233–238.
- [15] M.J. Zygmunt, *Non symmetric random walk on infinite graph*, Opuscula Math. **31**, (2011) 4, 669–674.

Marcin J. Zygmunt  
zygmunt@agh.edu.pl

AGH University of Science and Technology  
Faculty of Applied Mathematics  
al. A. Mickiewicza 30, 30-059 Krakow, Poland

*Received: March 8, 2015.*

*Revised: November 25, 2015.*

*Accepted: November 28, 2015.*