

TOWARDS THEORY OF \mathcal{C} -SYMMETRIES

S. Kuzhel and V. Sudilovskaya

Communicated by Aurelian Gheondea

Abstract. The concept of \mathcal{C} -symmetry originally appeared in \mathcal{PT} -symmetric quantum mechanics is studied within the Krein spaces framework.

Keywords: Krein space, J -self-adjoint operator, J -symmetric operator, Friedrichs extension, \mathcal{C} -symmetry.

Mathematics Subject Classification: 47A55, 47B25, 47A57, 81Q15

1. INTRODUCTION

Let \mathfrak{H} be a Hilbert space with inner product (\cdot, \cdot) linear in the first argument and let J be a non-trivial fundamental symmetry, i.e., $J = J^*$, $J^2 = I$, and $J \neq \pm I$.

The space \mathfrak{H} endowed with the indefinite inner product

$$[f, g] = (Jf, g) \tag{1.1}$$

is called a Krein space $(\mathfrak{H}, [\cdot, \cdot])$.

A linear operator H acting in \mathfrak{H} is called J -self-adjoint if $JH = H^*J$ and J -symmetric if $JH \subseteq H^*J$. The condition of J -symmetricity of H is equivalent to the relation

$$[Hf, g] = [f, Hg], \quad f, g \in D(H). \tag{1.2}$$

During last two decades, the theory of J -self-adjoint operators has found successful applications in studying nonself-adjoint Hamiltonians of \mathcal{PT} -symmetric quantum mechanics (PTQM). The Hamiltonians of PTQM are not self-adjoint with respect to the initial Hilbert space's inner product (\cdot, \cdot) but they possess a certain “more physical property” of symmetry, which does not depend on the choice of inner product (like \mathcal{PT} -symmetry property, see [6,7] and references therein). Typically, such nonself-adjoint operators can be interpreted as J -self-adjoint or J -symmetric operators for a suitable choice of Krein space. However, the self-adjointness of a Hamiltonian H in a Krein space cannot be satisfactory because it does not guarantee the unitarity of the dynamics

generated by H . In PTQM-related studies, this problem is solved by finding a new symmetry represented by a linear operator \mathcal{C} and such that the new inner product $(\cdot, \cdot)_{\mathcal{C}} := [\mathcal{C}\cdot, \cdot]$ ensures the self-adjointness of H in a (possible) new Hilbert space endowed with $(\cdot, \cdot)_{\mathcal{C}}$.

The description of a symmetry \mathcal{C} for a given nonself-adjoint Hamiltonian H is one of the key points in PTQM. Because of the complexity of the problem (since \mathcal{C} depends on the choice of H), it is not surprising that the majority of the available formulae are approximative [8, 11] and the relevant mathematical study of the concept of \mathcal{C} -symmetry is not yet developed well.

The concept of \mathcal{C} -symmetry in PTQM can be easily reformulated in the Krein space setting [1]. The crucial point here is that each operator \mathcal{C} in the Krein space $(\mathfrak{H}, [\cdot, \cdot])$ can be completely determined by an J -orthogonal pair of maximal positive \mathfrak{L}_+ and maximal negative \mathfrak{L}_- subspaces of \mathfrak{H} , where positivity (negativity) is understood with respect to the indefinite inner product $[\cdot, \cdot]$. The existence of a \mathcal{C} -symmetry for a J -symmetric operator H means that H has a maximal dual pair $\{\mathfrak{L}_+, \mathfrak{L}_-\}$ of invariant subspaces. This is an underlying mathematical structure that allows one to develop mathematically rigorous study of \mathcal{C} -symmetries.

It should be noted that operators associated with maximal dual pairs $\{\mathfrak{L}_+, \mathfrak{L}_-\}$ in the sense mentioned above have been considered in the mathematical literature before the development of PTQM. In particular, an unbounded operator \mathcal{C} was used in [5, Example 3.31] as a curious example of a J -self-adjoint and J -positive operator whose spectrum coincides with \mathbb{C} . \mathcal{C} -symmetries were also considered in [13, 21] (in these papers the concepts of positive symmetry and \sharp -positive \sharp unitary operators were used, respectively) in connection with some geometric aspects of Krein spaces and studies of J -unitary operators. In particular, an explicit formula for operators \mathcal{C} in terms of angular operators was obtained [13].

The present paper can be considered as an introduction to the full-scale theory of \mathcal{C} -symmetries inspired by the development of PTQM, where the concept of operator \mathcal{C} is consistently studied with the use of Krein spaces approach.

This paper is organized as follows. In Section 2, after preliminary results related to the Krein space theory, we discuss various definitions of operators \mathcal{C} which are typically used in literature. Further, the relation between operators \mathcal{C} and the concept of operator angles is studied and the difference between bounded and unbounded operators \mathcal{C} is explained. Section 3 deals with the case of J -self-adjoint operators with \mathcal{C} -symmetry. We show that J -self-adjoint operators with non-zero resolvent set may only have a property of bounded \mathcal{C} -symmetry, that is equivalent to the similarity of those operators to self-adjoint ones. Our presentation of Sections 2 and 3 is built in part on previous works [1, 10, 14, 17] where some properties of \mathcal{C} -operators have been investigated.

In Section 4, the method of construction of \mathcal{C} operators for J -symmetric operators with complete set of eigenfunctions corresponding to real eigenvalues is presented. We characterize (possible) nonuniqueness of the corresponding unbounded operators \mathcal{C} with the use of extension theory of symmetric operators. These results were inspired by [8, 9] where the non uniqueness of unbounded operator \mathcal{C} was discovered with the use of formal methods.

Finally, we remark that the area of applicability of the theory of \mathcal{C} -symmetries does not restricted to PTQM-related studies. In particular, the concept of \mathcal{C} -symmetry can be useful for the theory of operator differential equations in Krein spaces and for the solution of operator Riccati equations [14].

In what follows $D(H)$ and $\ker H$ denote, respectively, the domain and the kernel space of a linear operator H . The symbol $H \upharpoonright \mathcal{D}$ means the restriction of H onto a set \mathcal{D} . Let \mathfrak{H} be a complex Hilbert space. Sometimes, it is useful to specify the inner product (\cdot, \cdot) endowed with \mathfrak{H} . In that case the notation $(\mathfrak{H}, (\cdot, \cdot))$ will be used.

2. DEFINITION AND GENERAL PROPERTIES OF OPERATORS \mathcal{C}

2.1. ELEMENTS OF THE KREIN SPACE THEORY

Let $(\mathfrak{H}, [\cdot, \cdot])$ be a Krein space with a fundamental symmetry J . For the theory of Krein spaces and operators acting therein we refer the interested reader to [5].

A (closed) subspace \mathfrak{L} of the Hilbert space \mathfrak{H} is called *nonnegative*, *positive*, *uniformly positive* with respect to the indefinite inner product $[\cdot, \cdot]$ if, respectively, $[f, f] \geq 0$, $[f, f] > 0$, $[f, f] \geq \alpha \|f\|^2$, ($\alpha > 0$) for all $f \in \mathfrak{L} \setminus \{0\}$. Nonpositive, negative and uniformly negative subspaces are introduced similarly.

In each of the above mentioned classes we can define maximal subspaces. For instance, a closed positive subspace \mathfrak{L} is called *maximal positive* if \mathfrak{L} is not a proper subspace of a positive subspace in \mathfrak{H} . The concept of maximality for other classes is defined similarly.

Let \mathfrak{L}_+ be a maximal positive subspace. Then its J -orthogonal complement (i.e., orthogonal complement with respect to the indefinite inner product $[\cdot, \cdot]$)

$$\mathfrak{L}_- = \mathfrak{L}_+^{\perp} = \{f \in \mathfrak{H} : [f, g] = 0, \text{ for all } g \in \mathfrak{L}_+\} \quad (2.1)$$

is a maximal negative subspace of \mathfrak{H} , and the direct sum

$$\mathcal{D} = \mathfrak{L}_+ \dot{+} \mathfrak{L}_- \quad (2.2)$$

is a *dense set* in the Hilbert space \mathfrak{H} . The brackets in (2.2) indicates that the subspaces \mathfrak{L}_+ and \mathfrak{L}_- are J -orthogonal.

Denote

$$\mathfrak{H}_+ = \frac{1}{2}(I + J)\mathfrak{H}, \quad \mathfrak{H}_- = \frac{1}{2}(I - J)\mathfrak{H}, \quad (2.3)$$

where J is the operator of fundamental symmetry which determines indefinite inner product in (1.1). It is easy to see that the J -orthogonal subspaces \mathfrak{H}_\pm are maximal uniformly positive/negative with respect to $[\cdot, \cdot]$. Moreover, they are also orthogonal with respect to the initial inner product (\cdot, \cdot) and

$$\mathfrak{H} = \mathfrak{H}_+ \dot{+} \mathfrak{H}_-. \quad (2.4)$$

The decomposition (2.4) is called *the fundamental decomposition* of \mathfrak{H} .

The subspaces \mathfrak{L}_\pm in (2.2) are maximal positive/negative and their “deviation” from \mathfrak{H}_\pm are described by a self-adjoint strong contraction¹⁾ T , which anticommutes with J . Precisely,

$$\mathfrak{L}_+ = (I + T)\mathfrak{H}_+, \quad \mathfrak{L}_- = (I + T)\mathfrak{H}_-. \quad (2.5)$$

The operator T is called *an operator of transition* from the fundamental decomposition (2.4) to the direct sum (2.2) [14].

The collection of operators of transition admits a simple “external” description. Namely, a self-adjoint operator T in \mathfrak{H} is an operator of transition if and only if

$$\|Tf\| < \|f\| \quad (\text{for all } f \in \mathfrak{H}, f \neq 0), \quad JT = -TJ. \quad (2.6)$$

The subspaces $J\mathfrak{L}_\pm$ are maximal positive/negative and their J -orthogonal direct sum

$$JD = J\mathfrak{L}_+[\dot{+}]J\mathfrak{L}_- \quad (2.7)$$

is called *dual* to (2.2). The transition operator corresponding to the dual decomposition (2.7) coincides with $-T$.

Denote by $P_{\mathfrak{L}_\pm} : \mathcal{D} \rightarrow \mathfrak{L}_\pm$ the projection operators onto \mathfrak{L}_\pm with respect to the decomposition (2.2). The operators $P_{\mathfrak{L}_\pm}$ are defined on $\mathcal{D} = \mathfrak{L}_+[\dot{+}]\mathfrak{L}_-$ and

$$P_{\mathfrak{L}_\pm}f = P_{\mathfrak{L}_\pm}(f_{\mathfrak{L}_+} + f_{\mathfrak{L}_-}) = f_{\mathfrak{L}_\pm}, \quad f = f_{\mathfrak{L}_+} + f_{\mathfrak{L}_-} \in \mathcal{D}, \quad f_{\mathfrak{L}_\pm} \in \mathfrak{L}_\pm.$$

It is known [1] that

$$P_{\mathfrak{L}_+} = (I - T)^{-1}(P_+ - TP_-), \quad P_{\mathfrak{L}_-} = (I - T)^{-1}(P_- - TP_+), \quad (2.8)$$

where $P_+ = \frac{1}{2}(I + J)$ and $P_- = \frac{1}{2}(I - J)$ are orthogonal projection operators on \mathfrak{H}_+ and \mathfrak{H}_- , respectively.

2.2. DEFINITION AND GENERAL PROPERTIES OF OPERATORS \mathcal{C} .

Let a maximal positive subspace \mathfrak{L}_+ be given. Then its J -orthogonal complement $\mathfrak{L}_- = \mathfrak{L}_+^{\perp}$ is a maximal negative subspace and we can consider the direct sum (2.2).

The operator \mathcal{C} associated with the direct sum (2.2) is defined as follows: according to (2.2), any element $f \in \mathcal{D}$ admits the decomposition

$$f = f_{\mathfrak{L}_+} + f_{\mathfrak{L}_-}, \quad f_{\mathfrak{L}_\pm} \in \mathfrak{L}_\pm. \quad (2.9)$$

Then,

$$\mathcal{C}f = \mathcal{C}(f_{\mathfrak{L}_+} + f_{\mathfrak{L}_-}) = f_{\mathfrak{L}_+} - f_{\mathfrak{L}_-}. \quad (2.10)$$

The operator \mathcal{C} associated with the direct sum (2.2) is *J -self-adjoint and J -positive* [5, Example 3.31]. The latter means that $[\mathcal{C}f, f] > 0$ for all nonzero $f \in D(\mathcal{C}) = \mathcal{D}$.

¹⁾ “Strong contraction” means that $\|Tf\| < \|f\|$, $f \neq 0$.

The set of operators \mathcal{C} in the Krein space $(\mathfrak{H}, [\cdot, \cdot])$ is in one-to-one correspondence with the set of all possible J -orthogonal direct sums (2.2). If \mathcal{C} is given, then the corresponding maximal positive/negative subspaces \mathfrak{L}_\pm in (2.2) are recovered by the formula

$$\mathfrak{L}_\pm = \frac{1}{2}(I \pm \mathcal{C})\mathfrak{H}. \quad (2.11)$$

In view of (2.8) and (2.10)

$$\begin{aligned} \mathcal{C}f &= (P_{\mathfrak{L}_+} - P_{\mathfrak{L}_-})f = (I - T)^{-1}(P_+ - TP_- - P_- + TP_+)f \\ &= (I - T)^{-1}(J + TJ)f = J(I + T)^{-1}(I - T)f \end{aligned} \quad (2.12)$$

for all $f \in D(\mathcal{C})$.

The definition of \mathcal{C} with the use of (2.10) is not always convenient because it requires a description of the direct sum (2.2). An additional analysis of (2.12) allows one to establish:

Theorem 2.1. *The following statements are equivalent:*

- (i) *an operator \mathcal{C} is determined by the direct sum (2.2) with the use of the formula (2.10);*
- (ii) *an operator \mathcal{C} has the form $\mathcal{C} = Je^Q$, where Q is a self-adjoint operator in $(\mathfrak{H}, (\cdot, \cdot))$ such that $JQ = -QJ$;*
- (iii) *an operator \mathcal{C} satisfies the relation $\mathcal{C}^2 f = f$ for all $f \in D(\mathcal{C})$ and $J\mathcal{C}$ is a positive self-adjoint operator in \mathfrak{H} .*

The particular cases (the proof of equivalence of various relations between the items (i)–(iii) of the theorem) have been considered in [1], [10], and [17].

In what follows we will use assertions (i)–(iii) of Theorem 2.1 as (*equivalent definitions of operator \mathcal{C}*).

This means that each operator \mathcal{C} has the form $\mathcal{C} = Je^Q$ and, simultaneously, the same operator \mathcal{C} can be determined by (2.10) for a special choice of the direct sum (2.2).

Let us find the relationship between the operator of transition T from the fundamental decomposition (2.4) to (2.2) and the self-adjoint operator Q . In view of (2.12),

$$e^Q = (I + T)^{-1}(I - T)$$

or, after elementary transformations, $T = (I - e^Q)(I + e^Q)^{-1}$. Hence,

$$T = -\frac{e^{Q/2} - e^{-Q/2}}{2} \left(\frac{e^{Q/2} + e^{-Q/2}}{2} \right)^{-1} = -\frac{\sinh(Q/2)}{\cosh(Q/2)} = -\tanh \frac{Q}{2}.$$

Corollary 2.2. *The set of all operators \mathcal{C} is closed with respect to the calculation of the adjoint operator. The operator \mathcal{C}^* is determined by the dual direct sum (2.7).*

Proof. In view of statement (ii), the adjoint $\mathcal{C}^* = [Je^Q]^* = e^Q J = Je^{-Q}$ belongs to the set of operators \mathcal{C} . The operator e^{-Q} corresponds to the transition operator $-T$ from the fundamental decomposition (2.4) to (2.7). Therefore, \mathcal{C}^* is determined by the dual direct sum (2.7). \square

2.3. RELATION BETWEEN OPERATORS \mathcal{C} AND OPERATOR ANGLES.

Recall that the operator angle $\Theta(\mathfrak{M}, \mathfrak{N})$ between subspaces \mathfrak{M} and \mathfrak{N} measured relative to \mathfrak{M} is given by (see, e.g., [2])

$$\Theta(\mathfrak{M}, \mathfrak{N}) = \arcsin \sqrt{I_{\mathfrak{M}} - P_{\mathfrak{M}}P_{\mathfrak{N}}|_{\mathfrak{M}}},$$

where $I_{\mathfrak{M}}$ denotes the identity operator on \mathfrak{M} , and $P_{\mathfrak{M}}$ and $P_{\mathfrak{N}}$ stand for the orthogonal projections in \mathfrak{H} onto \mathfrak{M} and \mathfrak{N} , respectively.

By definition, the operator angle $\Theta(\mathfrak{M}, \mathfrak{N})$ is a bounded non-negative operator on \mathfrak{M} and

$$0 \leq \Theta(\mathfrak{M}, \mathfrak{N}) \leq \frac{\pi}{2}I.$$

This inequality can be specified for the cases where $\mathfrak{M} = \mathfrak{H}_{\pm}$ and $\mathfrak{N} = \mathfrak{L}_{\pm}$.

Lemma 2.3. *Let \mathfrak{H}_{\pm} and \mathfrak{L}_{\pm} be determined by (2.3) and (2.5), respectively. Then*

$$\Theta(\mathfrak{H}_{\pm}, \mathfrak{L}_{\pm}) = \arcsin \sqrt{T^2(I + T^2)^{-1}} = \arcsin |T| \sqrt{(I + |T|^2)^{-1}},$$

where $|T| = \sqrt{T^2}$ is the modulus of T .

Proof. Let $\mathfrak{M} = \mathfrak{H}_{+}$ and $\mathfrak{N} = \mathfrak{L}_{+}$. Denote by $\widehat{P}_{\mathfrak{L}_{+}}$ the orthogonal projection in \mathfrak{H} onto \mathfrak{L}_{+} . The orthogonal projection P_{+} in \mathfrak{H} onto \mathfrak{H}_{+} has the form $P_{+} = \frac{1}{2}(I + J)$.

By virtue of (2.1), $\mathfrak{H} = \mathfrak{L}_{+} \oplus J\mathfrak{L}_{-}$. Hence, any $f_{+} \in \mathfrak{H}_{+}$ can be decomposed:

$$f_{+} = (I + T)y_{+} + J(I + T)z_{-} = (I + T)y_{+} - (I - T)z_{-}, \quad y_{+} \in \mathfrak{H}_{+}, \quad z_{-} \in \mathfrak{H}_{-}.$$

Separation of the elements belonging to \mathfrak{H}_{\pm} , gives the system of equations $y_{+} - Tz_{-} = f_{+}$, $T_{+}y_{+} + z_{-} = 0$, which enables one to establish that $(I + T^2)y_{+} = f_{+}$ and $y_{+} = (I + T^2)^{-1}f_{+}$. Using the decomposition of f_{+} , we obtain

$$P_{+}\widehat{P}_{\mathfrak{L}_{+}}f_{+} = P_{+}(I + T)y_{+} = y_{+} = (I + T^2)^{-1}f_{+}.$$

Therefore,

$$(I - P_{+}\widehat{P}_{\mathfrak{L}_{+}})f_{+} = f_{+} - (I + T^2)^{-1}f_{+} = T^2(I + T^2)^{-1}f_{+}$$

that completes the proof for $\Theta(\mathfrak{H}_{+}, \mathfrak{L}_{+})$. The case $\Theta(\mathfrak{H}_{-}, \mathfrak{L}_{-})$ is considered similarly. \square

Corollary 2.4. *The operator angles $\Theta(\mathfrak{H}_{\pm}, \mathfrak{L}_{\pm})$ satisfy the inequality*

$$0 \leq \Theta(\mathfrak{H}_{\pm}, \mathfrak{L}_{\pm}) \leq \frac{\pi}{4}I. \quad (2.13)$$

The operator \mathcal{C} corresponding to the direct sum (2.2) of the subspaces \mathfrak{L}_{\pm} is bounded if and only if

$$\|\Theta(\mathfrak{H}_{\pm}, \mathfrak{L}_{\pm})\| = \max \left\{ \lambda : \lambda \in \sigma(\Theta(\mathfrak{H}_{\pm}, \mathfrak{L}_{\pm})) \right\} < \frac{\pi}{4}.$$

Proof. In view of Lemma 2.3, $\Theta(\mathfrak{H}_\pm, \mathfrak{L}_\pm)$ are the functions of a nonnegative self-adjoint contractions $|T|_\pm = |T| \upharpoonright_{\mathfrak{H}_\pm}$ acting in \mathfrak{H}_\pm . Precisely,

$$\Theta(\mathfrak{H}_\pm, \mathfrak{L}_\pm) = g(|T|_\pm), \quad g(\lambda) = \arcsin \left(\frac{\lambda}{\sqrt{1 + \lambda^2}} \right). \quad (2.14)$$

This formula immediately leads to (2.13) since $0 \leq g(\lambda) \leq \frac{\pi}{4}$ for $\lambda \in [0, 1] \supset \sigma(|T|_\pm)$. The second assertion of corollary 2.4 also follows from (2.14) and the fact that an operator \mathcal{C} is bounded if and only if the corresponding operator of transition T is an uniformly strong contraction (i.e., $\|T\| < 1$). \square

An elementary calculation shows (see, e.g., [15]) that

$$|T|_\pm = \tan \Theta(\mathfrak{H}_\pm, \mathfrak{L}_\pm).$$

On the other hand,

$$|T| = \left| \tanh \frac{Q}{2} \right| = |T|_+ P_+ + |T|_- P_-.$$

Therefore, the relationship between the self-adjoint operator Q in the formula $\mathcal{C} = J e^Q$ and the operator angles $\Theta(\mathfrak{H}_\pm, \mathfrak{L}_\pm)$ where \mathfrak{L}_\pm are defined by \mathcal{C} is:

$$\left| \tanh \frac{Q}{2} \right| = \tan \Theta(\mathfrak{H}_+, \mathfrak{L}_+) P_+ + \tan \Theta(\mathfrak{H}_-, \mathfrak{L}_-) P_-.$$

2.4. BOUNDED AND UNBOUNDED OPERATORS \mathcal{C}

There is an essential difference between the properties of bounded and unbounded operators \mathcal{C} . In particular, the spectrum of a bounded operator \mathcal{C} coincides with eigenvalues $\{-1, 1\}$, while the spectrum of unbounded \mathcal{C} additionally has the continuous part on $\mathbb{C} \setminus \{-1, 1\}$ ([1]).

If \mathcal{C} is bounded, then: $\mathcal{C} = J e^Q$, where Q is a bounded self-adjoint operator in \mathfrak{H} ; the subspaces \mathfrak{L}_\pm in (2.2) turns out to be *maximal uniformly* positive/negative in the Krein space $(\mathfrak{H}, [\cdot, \cdot])$; the direct sum (2.2) gives the decomposition of the whole space \mathfrak{H} :

$$\mathfrak{H} = \mathfrak{L}_+ [+] \mathfrak{L}_-; \quad (2.15)$$

the given indefinite inner product $[\cdot, \cdot]$ generates infinitely many equivalent inner products $(\cdot, \cdot)_{\mathcal{C}}$ in the Hilbert space $(\mathfrak{H}, (\cdot, \cdot))$, which are determined by the choice of bounded operators \mathcal{C} :

$$(\cdot, \cdot)_{\mathcal{C}} = [\mathcal{C}\cdot, \cdot] = (J\mathcal{C}\cdot, \cdot) = (e^Q \cdot, \cdot); \quad (2.16)$$

the subspaces \mathfrak{L}_\pm in (2.15) are mutually orthogonal with respect to $(\cdot, \cdot)_{\mathcal{C}}$ and \mathcal{C} turns out to be the fundamental symmetry operator in the Hilbert space $(\mathfrak{H}, (\cdot, \cdot)_{\mathcal{C}})$.

If \mathcal{C} is unbounded, then: $\mathcal{C} = J e^Q$, where Q is an unbounded self-adjoint operator in \mathfrak{H} ; the subspaces \mathfrak{L}_\pm in (2.2) are *maximal* positive/negative in the Krein space $(\mathfrak{H}, [\cdot, \cdot])$; the direct sum (2.2) is a dense subset of \mathfrak{H} ; the formula (2.16) determines

infinitely many inner products $(\cdot, \cdot)_{\mathcal{C}}$ defined on various sets $\mathcal{D} = D(\mathcal{C})$; the linear spaces \mathcal{D} endowed with inner products $(\cdot, \cdot)_{\mathcal{C}}$ are pre-Hilbert spaces; the completion $\mathfrak{H}_{\mathcal{C}}$ of \mathcal{D} with respect to $(\cdot, \cdot)_{\mathcal{C}}$ does not coincide with \mathfrak{H} ; the Hilbert space $(\mathfrak{H}_{\mathcal{C}}, (\cdot, \cdot)_{\mathcal{C}})$ has the decomposition

$$\mathfrak{H}_{\mathcal{C}} = \hat{\mathfrak{L}}_+ \oplus_{\mathcal{C}} \hat{\mathfrak{L}}_-, \quad (2.17)$$

where the mutually orthogonal²⁾ subspaces $\hat{\mathfrak{L}}_{\pm}$ are the completion of \mathfrak{L}_{\pm} with respect to $(\cdot, \cdot)_{\mathcal{C}}$; the decomposition (2.17) gives rise to the new Krein space $(\mathfrak{H}_{\mathcal{C}}, [\cdot, \cdot]')$ with the indefinite inner product

$$[f, g]' = (f_{\hat{\mathfrak{L}}_+}, g_{\hat{\mathfrak{L}}_+})_{\mathcal{C}} - (f_{\hat{\mathfrak{L}}_-}, g_{\hat{\mathfrak{L}}_-})_{\mathcal{C}}, \quad f = f_{\hat{\mathfrak{L}}_+} + f_{\hat{\mathfrak{L}}_-}, \quad g = g_{\hat{\mathfrak{L}}_+} + g_{\hat{\mathfrak{L}}_-},$$

which coincides with the original indefinite inner product $[\cdot, \cdot]$ on \mathcal{D} .

3. J -SELF-ADJOINT OPERATORS WITH PROPERTY OF \mathcal{C} -SYMMETRY

Definition 3.1. A densely defined operator H in \mathfrak{H} has the property of \mathcal{C} -symmetry if there exists an operator \mathcal{C} with properties described in Theorem 2.1 and such that

$$HCf = CHf, \quad f \in D(H). \quad (3.1)$$

We will say that H has *bounded (unbounded) \mathcal{C} -symmetry* if the corresponding operator \mathcal{C} is bounded (unbounded).

The commutation identity (3.1) requires an additional explanation because \mathcal{C} may be unbounded. Precisely, if (3.1) is satisfied, then

$$D(\mathcal{C}) \supset D(H), \quad \mathcal{C} : D(H) \rightarrow D(H), \quad H : D(H) \rightarrow D(\mathcal{C}).$$

If \mathcal{C} is a bounded operator, then the first and third relations are trivial since $D(\mathcal{C}) = \mathfrak{H}$.

The existence of \mathcal{C} -symmetry for H means that the J -orthogonal direct sum (2.2), where $\mathfrak{L}_{\pm} = \frac{1}{2}(I \pm \mathcal{C})\mathfrak{H}$ decomposes the operator H into the sum

$$H = H_+ \dot{+} H_-, \quad D(H) = D(H_+) \dot{+} D(H_-), \quad D(H_{\pm}) \subset \mathfrak{L}_{\pm} \quad (3.2)$$

of operators $H_{\pm} : D(H_{\pm}) \rightarrow \mathfrak{L}_{\pm}$ acting in \mathfrak{L}_{\pm} .

If H is a J -self-adjoint (J -symmetric), then its components H_{\pm} in (3.2) should be J -self-adjoint (J -symmetric) operators in \mathfrak{L}_{\pm} . In this case H turns out to be a self-adjoint (symmetric) operator with respect to the new inner product $(\cdot, \cdot)_{\mathcal{C}}$. Therefore, *the description of \mathcal{C} -symmetry for a J -self-adjoint (J -symmetric) operator H leads to the explicit construction of the new inner product $(\cdot, \cdot)_{\mathcal{C}}$ which ensures the self-adjointness (the symmetricity) of H .*

Proposition 3.2. *Let H be a J -self-adjoint operator with non-empty resolvent set $\rho(H) \neq \emptyset$. Then H may only have a property of bounded \mathcal{C} -symmetry.*

²⁾ with respect to the inner product $(\cdot, \cdot)_{\mathcal{C}}$

Proof. If H is J -self-adjoint, then H should be a closed operator in \mathfrak{H} . Let $\lambda \in \rho(H)$. Then the operator $H - \lambda I$ commutes with the same operator \mathcal{C} and $D(\mathcal{C}) \supseteq \mathcal{R}(H - \lambda) = \mathfrak{H}$. Therefore, \mathcal{C} is a bounded operator. \square

Remark 3.3.

1. Proposition 3.2 may seem paradoxical only at the first glance. Indeed, if \mathcal{C} is unbounded, then the new inner product $(\cdot, \cdot)_{\mathcal{C}}$ is singular with respect to the original inner product (\cdot, \cdot) . Therefore, the closedness of H with respect to (\cdot, \cdot) together with the condition $\rho(H) \neq \emptyset$ do not fit the singular inner product $(\cdot, \cdot)_{\mathcal{C}}$. For this reason, the property of unbounded \mathcal{C} -symmetry is more natural for the case of closable (but not closed) operators.

2. The condition $\rho(H) \neq \emptyset$ is important in Proposition 3.2. Indeed, every unbounded operator $H = \mathcal{C}$ is closed (since \mathcal{C} is J -self-adjoint), its spectrum coincides with \mathbb{C} and, evidently, this operator has the property of unbounded \mathcal{C} -symmetry.

The next result is a direct consequence of Theorem 1.9 in [1] and Proposition 3.2.

Theorem 3.4. *Let H be a J -self-adjoint operator. Then the following assertions are equivalent:*

- (i) H has the property of \mathcal{C} -symmetry;
- (ii) H is a self-adjoint operator with respect to a new (equivalent to (\cdot, \cdot)) inner product in the Hilbert space $(\mathfrak{H}, (\cdot, \cdot))$;
- (iii) H is similar³⁾ to a self-adjoint operator in $(\mathfrak{H}, (\cdot, \cdot))$.

The item (iii) of Theorem 3.4 allows one to characterize the (possible) existence of \mathcal{C} -symmetry for a J -self-adjoint operator H with the use of a well-known criterion of similarity (see, e.g., [20]). Precisely, a J -self-adjoint operator H has the property of \mathcal{C} -symmetry if and only if it has a real spectrum and there exists a constant M such that

$$\sup_{\varepsilon > 0} \varepsilon \int_{-\infty}^{\infty} \|(H - zI)^{-1} f\|^2 d\xi \leq M \|f\|^2, \quad f \in \mathfrak{H}, \quad (3.3)$$

where the integral is taken along the line $z = \xi + i\varepsilon$ ($\varepsilon > 0$ is fixed).

However, the relation (3.3) does not answer how to construct the corresponding operator \mathcal{C} .

The explicit formulas for \mathcal{C} can be derived for special classes of J -self-adjoint operators:

I. Let us assume that a J -self-adjoint operator H is J -nonnegative, i.e., $[Hf, f] \geq 0$ for all $f \in D(H)$. If the resolvent set $\rho(H)$ is nonempty, then the spectrum of H is real [5, Theorem 3.27]. Moreover, H has a spectral function $E_H(\cdot)$ with properties similar to that of a spectral function of a self-adjoint operator [19]. The main difference is the occurrence of critical points $\{0, \infty\}$. Note that ∞ is always a critical point of H , and 0 may be its critical point. Significantly different behavior of the spectral function $E_H(\cdot)$ occurs at *singular* critical points in any neighborhood of which $E_H(\cdot)$ is unbounded.

The next statement is a direct consequence of [19, Theorem 5.7]

³⁾ An operator H is called similar to a self-adjoint operator A if there exists a bounded and boundedly invertible operator Z such that $ZA = HZ$.

Proposition 3.5. *Let H be a J -nonnegative and J -self-adjoint operator. Assume that the resolvent set $\rho(H)$ is non-empty, 0 and ∞ are not singular critical points and 0 is not an eigenvalue of H . Then H has a bounded \mathcal{C} -symmetry $\mathcal{C} = E_H(\mathbb{R}_+) - E_H(\mathbb{R}_-)$, where $E_H(\cdot)$ is the spectral function of H .*

The operator

$$H = (\operatorname{sgn} x) \frac{d^2}{dx^2}$$

defined on its maximal domain in $L_2(\mathbb{R})$ is J -nonnegative and J -self-adjoint with the fundamental symmetry $Jf = (\operatorname{sgn} x)f$ in $L_2(\mathbb{R})$. It was showed [12] that H satisfies all assumptions of Proposition 3.5. Hence, H has a bounded \mathcal{C} -symmetry.

II. Let H be a J -self-adjoint operator with Riesz basis of eigenvectors $\{f_n\}$ corresponding to real simple eigenvalues. Then [1, Theorem 6.3.7], H has the property of \mathcal{C} -symmetry, the corresponding bounded operator \mathcal{C} is determined uniquely and

$$\mathcal{C}f = \sum_{n=1}^{\infty} [f, f_n] f_n, \quad f \in \mathfrak{H}.$$

(Without loss of generality, in the last formula, we assume that $\{f_n\}$ are *normalized with respect to the indefinite metric* $[\cdot, \cdot]$, i.e., $[f_n, f_n]^2 = 1$.)

4. \mathcal{C} -SYMMETRY FOR J -SYMMETRIC OPERATORS WITH A COMPLETE SET OF EIGENVECTORS

4.1. DESCRIPTION OF OPERATORS \mathcal{C} ASSOCIATED WITH COMPLETE SET OF EIGENVECTORS

Let a J -symmetric operator H have a complete set of eigenvectors $\{f_n\}_{n=1}^{\infty}$ corresponding to real simple eigenvalues $\{\lambda_n\}_{n=1}^{\infty}$. It should be noted that $[f_n, f_n] \neq 0$ for any eigenfunction f_n . This fact is crucial for our investigations. Let us verify it. Indeed, it is well known that eigenfunctions corresponding to different real eigenvalues of a J -symmetric operator are J -orthogonal [5]. Hence, if there exists an eigenfunction f_m such that $[f_m, f_m] = 0$, then f_m will be J -orthogonal to the linear span \mathcal{S} of $\{f_n\}$. The latter means that Jf_m is orthogonal to the dense set \mathcal{S} in the Hilbert space $(\mathfrak{H}, (\cdot, \cdot))$. Therefore, $Jf_m = 0$ and $f_m = 0$ that is impossible.

Separating the sequence of eigenvectors $\{f_n\}$ by the signs of $[f_n, f_n]$:

$$f_n = \begin{cases} f_n^+ & \text{if } [f_n, f_n] > 0, \\ f_n^- & \text{if } [f_n, f_n] < 0 \end{cases} \quad (4.1)$$

we obtain two sequences of positive $\{f_n^+\}$ and negative $\{f_n^-\}$ elements of the Krein space $(\mathfrak{H}, [\cdot, \cdot])$.

Let \mathfrak{L}_+^0 and \mathfrak{L}_-^0 be the closure (with respect to the initial inner product (\cdot, \cdot)) of the linear spans generated by the sets $\{f_n^+\}$ and $\{f_n^-\}$, respectively. By construction, \mathfrak{L}_{\pm}^0 are J -orthogonal positive/negative subspaces in $(\mathfrak{H}, [\cdot, \cdot])$ and the direct sum

$$\mathcal{D}_0 = \mathfrak{L}_+^0 [+] \mathfrak{L}_-^0 \quad (4.2)$$

is a dense set in \mathfrak{H} (since \mathcal{D}_0 contains \mathcal{S}).

Let \mathcal{C}_0 be an operator associated with (4.2):

$$\mathcal{C}_0 f = f_{\mathfrak{L}_+^0} - f_{\mathfrak{L}_-^0}, \quad f \in D(\mathcal{C}_0) = \mathcal{D}_0. \quad (4.3)$$

It follows from (4.3) that the operator \mathcal{C}_0 is densely defined and $\mathcal{C}_0^2 f = f$ for any $f \in D(\mathcal{C}_0)$. However, in general case, the operator \mathcal{C}_0 *cannot be considered* as an operator of \mathcal{C} -symmetry. The matter is that the subspaces \mathfrak{L}_\pm^0 in (4.2) may be *proper subspaces of a maximal definite subspaces* \mathfrak{L}_\pm .

If at least one of subspaces \mathfrak{L}_\pm^0 loses the property of being maximal in the classes of definite subspaces (positive or negative), then the sum (4.2) *cannot be domain of definition* of an operator of \mathcal{C} -symmetry with properties described in Theorem 2.1. In other words, the operator \mathcal{C}_0 cannot be presented as Je^Q , where e^Q is a *self-adjoint operator*. Of course, this phenomenon is possible only in the case of unbounded operators \mathcal{C} .

In order to construct a proper operator \mathcal{C} associated with (4.2) we have to extend (4.2) to a J -orthogonal direct sum $\mathfrak{L}_+[\dot{+}]\mathfrak{L}_-$, where $\mathfrak{L}_\pm \supset \mathfrak{L}_\pm^0$ are *maximal* definite subspaces (positive and negative). Obviously, the operator \mathcal{C} determined by the direct sum $\mathfrak{L}_+[\dot{+}]\mathfrak{L}_-$ will be an extension of \mathcal{C}_0 .

In general, the mentioned extension

$$\mathfrak{L}_+^0[\dot{+}]\mathfrak{L}_-^0 \rightarrow \mathfrak{L}_+[\dot{+}]\mathfrak{L}_- \quad (4.4)$$

is not determined uniquely⁴⁾. This leads to the *nonuniqueness* of unbounded operators of \mathcal{C} -symmetry $\mathcal{C} \supset \mathcal{C}_0$ associated with $\mathfrak{L}_+^0[\dot{+}]\mathfrak{L}_-^0$. Therefore, the complete set of eigenvectors $\{f_n\}$ does not always determine a unique operator \mathcal{C} .

In order to describe all possible extensions (4.4) we introduce the operator $G_0 = J\mathcal{C}_0$, where \mathcal{C}_0 is defined by (4.3).

Lemma 4.1. *The operator G_0 is a closed positive symmetric operator in $(\mathfrak{H}, (\cdot, \cdot))$ and*

$$JG_0JG_0f = f, \quad f \in \mathcal{D}_0 = D(G_0) = D(\mathcal{C}_0). \quad (4.5)$$

Proof. Repeating the proof of Lemma 6.2.3 in [1] we decide that \mathcal{C}_0 is a closed operator in \mathfrak{H} . Therefore, the operator $G_0 = J\mathcal{C}_0$ is closed too. The positivity of G_0 follows from the fact that

$$(G_0f, f) = [\mathcal{C}_0f, f] = [f_{\mathfrak{L}_+^0}, f_{\mathfrak{L}_+^0}] - [f_{\mathfrak{L}_-^0}, f_{\mathfrak{L}_-^0}] > 0$$

for all nonzero $f \in \mathcal{D}_0$. The “boundary condition” (4.5) is the direct consequence of the relation $\mathcal{C}_0^2 = I$ on $D(\mathcal{C}_0)$. \square

The next result reduces the description of all possible operators \mathcal{C} associated with the initial direct sum (4.2) to the description of a special class of positive self-adjoint extensions G of G_0 .

⁴⁾ This phenomenon was discovered by Langer [18].

Proposition 4.2. *The set of all positive self-adjoint extensions G of G_0 satisfying the condition*

$$JGJGf = f, \quad f \in D(G) \quad (4.6)$$

is in one-to-one correspondence with the set of all possible extensions (4.4), where \mathfrak{L}_\pm are J -orthogonal maximal positive/negative subspaces in the Krein space $(\mathfrak{H}, [\cdot, \cdot])$.

Proof. Let G be a positive self-adjoint extension of G_0 such that (4.6) holds. Then $\mathcal{C} = JG$ satisfies assertion (iii) in Theorem 2.1. Therefore, the operator \mathcal{C} determines the J -orthogonal maximal positive/negative subspaces \mathfrak{L}_\pm by the formula (2.11). These subspaces are extensions of \mathfrak{L}_\pm^0 since \mathcal{C} is an extension of \mathcal{C}_0 .

Conversely, each extension (4.4) defines an operator \mathcal{C} by formula (2.10). This operator is an extension of \mathcal{C}_0 and, hence the positive self-adjoint operator $G = J\mathcal{C}$ is an extension of $G_0 = J\mathcal{C}_0$. \square

4.2. THE CASE OF FRIEDRICHS EXTENSION

Let $D[G_0] \subset \mathfrak{H}$ be the closure of $D(G_0)$ with respect to the norm

$$\|f\|_0^2 = \|f\|^2 + (G_0f, f), \quad f \in D(G_0). \quad (4.7)$$

The operator G_F defined as follows:

$$G_F f = G_0^* f, \quad f \in D(G_F) = D[G_0] \cap D(G_0^*)$$

is called the Friedrichs extension of G_0 .

The Friedrichs extension G_F preserves the lower bound of G_0 :

$$\inf_{f \in D(G_0) \setminus \{0\}} \frac{(G_0f, f)}{(f, f)} = \inf_{f \in D(G_F) \setminus \{0\}} \frac{(G_F f, f)}{(f, f)}.$$

and it is a unique positive self-adjoint extension having the domain in $D_0[G_0]$ [3, 16].

Theorem 4.3. *The Friedrichs extension G_F satisfies the relation*

$$JG_F JG_F f = f, \quad f \in D(G_F)$$

if and only if G_F is the unique positive self-adjoint extension of G_0 .

Proof. The operator G_0 has the inverse G_0^{-1} which is defined on the dense set $JD(G_0)$ in \mathfrak{H} and the condition (4.5) can be rewritten as

$$JG_0 f = G_0^{-1} Jf, \quad f \in D(G_0). \quad (4.8)$$

By virtue of (4.8) the operator J isometrically maps the Hilbert space $(D[G_0], \|\cdot\|_0)$ onto the Hilbert space $(D[G_0^{-1}], \|\cdot\|_1)$, where $D[G_0^{-1}]$ is the completion of the domain of definition $D(G_0^{-1}) = JD(G_0)$ with respect to the norm (cf (4.7))

$$\|g\|_1^2 = \|g\|^2 + (G_0^{-1}g, g), \quad g \in JD(G_0).$$

The calculation of adjoint operators in (4.8) gives $G_0^*J = J(G_0^{-1})^*$ or $JG_0^* = (G_0^{-1})^*J$. Therefore, for any $f \in D(G_F) = D[G_0] \cap D(G_0^*)$,

$$JG_0^*f = JG_Ff = (G_0^{-1})^*Jf = (G_0^{-1})_FJf,$$

where $Jf \in D[G_0^{-1}] \cap D((G_0^{-1})^*)$ and $(G_0^{-1})_F$ is the Friedrichs extension of the positive symmetric operator G_0^{-1} . Using the well-known relation $(G_0^{-1})_F = (G_K)^{-1}$ (see, e.g. [3]), where G_K is the Krein-von Neumann extension of G_0 , we conclude that

$$JG_Ff = (G_K)^{-1}Jf, \quad f \in D(G_F).$$

Therefore, the required condition $JG_FJG_F = I$ holds if and only if the Friedrichs extension G_F of G_0 coincides with the Krein-von Neumann extension G_K . This is possible if and only if G_0 has a unique positive self-adjoint extension [16]. \square

Corollary 4.4. *The operator $\mathcal{C} = JG_F$ is the operator of \mathcal{C} -symmetry associated with (4.2) if and only if*

$$\inf_{f \in D(G_0) \setminus \{0\}} \frac{(G_0f, f)}{|(f, g)|^2} = 0 \quad (4.9)$$

for all nonzero vectors $g \in \ker(I + G_0^*)$.

Proof. It follows from Proposition 4.2 and Theorem 4.3 that the operator $\mathcal{C} = JG_F$ determines a J -orthogonal sum of maximal definite subspaces \mathfrak{L}_\pm which is the extension of (4.2) if and only if G_F is the unique positive self-adjoint extension of G_0 . The latter is equivalent to the condition (4.9) [16, Theorem 9]. \square

Let \mathcal{C} be an operator of \mathcal{C} -symmetry associated with (4.2) and let $\mathfrak{H}_\mathcal{C}$ be the corresponding Hilbert space obtained by the completion of $\mathfrak{L}_+[\dot{+}]\mathfrak{L}_-$ with respect to the inner product $(\cdot, \cdot)_\mathcal{C}$ (see (2.17)).

By construction, the original direct sum $\mathfrak{L}_+^0[\dot{+}]\mathfrak{L}_-^0$ determined by the eigenfunctions $\{f_n\}$ is a linear manifold in the new Hilbert space $(\mathfrak{H}_\mathcal{C}, (\cdot, \cdot)_\mathcal{C})$. However, in the general case, we cannot state that this linear manifold is a dense set in $\mathfrak{H}_\mathcal{C}$.

Theorem 4.5. *If the operator \mathcal{C} is determined by the Friedrichs extension G_F (i.e., if the condition (4.9) holds), then the linear span of eigenvectors $\{f_n\}$ is a dense set in $\mathfrak{H}_\mathcal{C}$*

Proof. The Friedrichs extension G_F is an example of *extremal extension* of G_0 [4]. By definition, a nonnegative self-adjoint extension G of G_0 belongs to the class $E(G_0)$ of extremal extensions if

$$\inf_{f \in D(G_0)} (G(\phi - f), (\phi - f)) = 0 \quad \text{for all } \phi \in D(G). \quad (4.10)$$

Since $\|\phi - f\|_\mathcal{C}^2 = [\mathcal{C}(\phi - f), \phi - f] = (G(\phi - f), (\phi - f))$, the condition (4.10) means that each element $\phi \in D(\mathcal{C}) = D(G)$ can be approximated by elements $f \in D(G_0) = \mathfrak{L}_+^0[\dot{+}]\mathfrak{L}_-^0$ in $\mathfrak{H}_\mathcal{C}$. Now, in order to complete the proof, it suffices to verify that the span of $\{f_n\}$ is a dense subset of $D(G_0)$ with respect to $(\cdot, \cdot)_\mathcal{C}$.

The inner product $(\cdot, \cdot)_{\mathcal{C}}$ considered on \mathfrak{L}_{\pm}^0 coincides with $\pm[\cdot, \cdot]$ and satisfies the evaluation

$$(f_{\mathfrak{L}_{\pm}^0}, f_{\mathfrak{L}_{\pm}^0})_{\mathcal{C}} = |(\mathcal{P}f_{\mathfrak{L}_{\pm}^0}, f_{\mathfrak{L}_{\pm}^0})| \leq \|\mathcal{P}f_{\mathfrak{L}_{\pm}^0}\| \|f_{\mathfrak{L}_{\pm}^0}\| = \|f_{\mathfrak{L}_{\pm}^0}\|^2 = (f_{\mathfrak{L}_{\pm}^0}, f_{\mathfrak{L}_{\pm}^0})$$

for any $f_{\mathfrak{L}_{\pm}^0} \in \mathfrak{L}_{\pm}^0$. Therefore, each element $f_{\mathfrak{L}_{\pm}^0} \in \mathfrak{L}_{\pm}^0$ can be approximated by $\text{span}\{f_n^{\pm}\}$ with respect to $(\cdot, \cdot)_{\mathcal{C}}$ since \mathfrak{L}_{\pm}^0 are closure of $\text{span}\{f_n^{\pm}\}$ with respect to (\cdot, \cdot) . This means that $\text{span}\{f_n\} = \text{span}\{f_n^+\} \cup \text{span}\{f_n^-\}$ is a dense set in $D(G_0)$ with respect to $(\cdot, \cdot)_{\mathcal{C}}$. \square

Corollary 4.6. *If the operator \mathcal{C} is determined by the Friedrichs extension G_F , then the normalized eigenvectors⁵⁾ $\{f_n\}$ forms an orthonormal basis of the Hilbert space $(\mathfrak{H}_{\mathcal{C}}, (\cdot, \cdot)_{\mathcal{C}})$.*

Proof. By virtue of (2.16) and (4.3), the normalized eigenvectors $\{f_n\}$ are an orthonormal system in the Hilbert space $(\mathfrak{H}_{\mathcal{C}}, (\cdot, \cdot)_{\mathcal{C}})$. Due to Theorem 4.5, the linear span of $\{f_n\}$ is a dense set in $\mathfrak{H}_{\mathcal{C}}$. Therefore, $\{f_n\}$ turns out to be an orthonormal basis of this Hilbert space. \square

Up to now we do not specify the domain of definition of the J -symmetric operator H . Let the domain $D(H)$ coincide with the linear span of $\{f_n\}$. Then relation (3.1) holds for any operator \mathcal{C} associated with (4.2). If the Friedrichs extension G_F determines an operator of \mathcal{C} -symmetry (i.e., if condition (4.9) holds), then the operator H turns out to be essentially self-adjoint in the Hilbert space $(\mathfrak{H}_{\mathcal{C}_F}, (\cdot, \cdot)_{\mathcal{C}_F})$ where $\mathcal{C}_F = JG_F$ and $\sigma(H)$ coincides with the closure of $\{\lambda_n\}$.

Example 4.7. Schauder basis of eigenvectors.

Let normalized eigenvectors $\{f_n\}_{n=1}^{\infty}$ of a J -symmetric operator H with real simple eigenvalues form a Schauder basis. We recall that a sequence $\{f_n\}$ is called a *Schauder basis* of a Hilbert space \mathfrak{H} if, for each $f \in \mathfrak{H}$, there exist uniquely determined scalar coefficients $\{c_n\}$ such that

$$f = \sum_{n=1}^{\infty} c_n f_n.$$

The coefficients $\{c_n\}$ can easy be specified with the use of a biorthogonal sequence⁶⁾ $\{g_n\}$, namely: $c_n = (f, g_n)$. Since the eigenvectors $\{f_n\}$ are mutually J -orthogonal and normalized, the sequence $\{g_n\}$ has the form $g_n = [f_n, f_n] J f_n$. Therefore, taking into account the separation (4.1) of eigenvectors $\{f_n\}$ by the signs of $[f_n, f_n]$, we obtain

$$f = \sum_{n=1}^{\infty} [f_n, f_n] [f, f_n] f_n = \sum_{n=1}^{\infty} [f, f_n^+] f_n^+ - \sum_{n=1}^{\infty} [f, f_n^-] f_n^-, \quad f \in \mathfrak{H}.$$

⁵⁾ That is, $\{f_n\}$ are normalized with respect to the indefinite metric $[\cdot, \cdot]$: $[f_n, f_n]^2 = 1$.

⁶⁾ Biorthonormality of sequences $\{f_n\}$ and $\{g_n\}$ means that $(f_n, g_m) = \delta_{nm}$.

If $\{f_n\}$ is a Schauder basis, then the subspaces \mathfrak{L}_{\pm}^0 defined as the closure of the linear spans of $\{f_n^{\pm}\}$, respectively are maximal positive/negative in the Krein space $(\mathfrak{H}, [\cdot, \cdot])$ [5, Statement 10.12 in Chapter 1]. Therefore, the formula (4.3) determines an unbounded operator $\mathcal{C} = \mathcal{C}_0$ with the domain of definition (4.2). For each $f \in D(\mathcal{C}) = \mathcal{D}_0$,

$$\mathcal{C}f = f_{\mathfrak{L}_+^0} - f_{\mathfrak{L}_-^0} = \sum_{n=1}^{\infty} [f_{\mathfrak{L}_+^0}, f_n^+] f_n^+ + \sum_{n=1}^{\infty} [f_{\mathfrak{L}_-^0}, f_n^-] f_n^- = \sum_{n=1}^{\infty} [f, f_n] f_n.$$

The operator $G_0 = J\mathcal{C}$ is a positive self-adjoint operator. Therefore, its Friedrichs extension coincides with G_0 .

If the domain $D(H)$ coincides with the linear span of $\{f_n\}$, then H has the property of \mathcal{C} -symmetry with the operator \mathcal{C} determined above. The operator H will be essentially self-adjoint in the new Hilbert space $(\mathfrak{H}_{\mathcal{C}}, (\cdot, \cdot)_{\mathcal{C}})$.

REFERENCES

- [1] S. Albeverio, S. Kuzhel, *\mathcal{PT} -Symmetric Operators in Quantum Mechanics: Krein Spaces Methods*, [in:] *Non-Selfadjoint Operators in Quantum Physics: Mathematical Aspects*, Fabio Bagarello, Jean-Pierre Gazeau, Franciszek H. Szafraniec, Miloslav Znojil (eds), John Wiley & Sons, Inc. 2015.
- [2] S. Albeverio, A. Motovilov, A. Shkalikov, *Bounds on variation of spectral subspaces under J -self-adjoint perturbations*, Integr. Equ. Oper. Theory **64** (2009), 455–486.
- [3] Yu. Arlinskiĭ, E. Tsekanovskii, *M. Krein's research on semi-bounded operators, its contemporary developments, and applications*, Operator Theory: Advances and Applications **190** (2009), 65–112.
- [4] Yu. M. Arlinskiĭ, S. Hassi, Z. Sebestyén, H.S.V. de Snoo, *On the class of extremal extensions of a nonnegative operator*, Recent Advances in Operator Theory and Related Topics The Bela Szokefalvi-Nagy Memorial Volume, Operator Theory: Advances and Applications **127** (2001), 41–81.
- [5] T.Ya. Azizov, I.S. Iokhvidov, *Linear Operators in Spaces with Indefinite Metric*, Wiley, Chichester, 1989.
- [6] C.M. Bender, *Introduction to \mathcal{PT} -symmetric quantum theory*, Contemp. Phys. **46** (2005) 277–292.
- [7] C.M. Bender, *Making sense of non-Hermitian Hamiltonians*, Rep. Prog. Phys. **70** (2007), 947–1018.
- [8] C.M. Bender, M. Gianfreda, *Nonuniqueness of the \mathcal{C} operator in \mathcal{PT} -symmetric quantum mechanics*, J. Phys. A. **46** (2013), 275306–275324.
- [9] C.M. Bender, S.P. Klevansky, *Nonunique \mathcal{C} operator in \mathcal{PT} quantum mechanics*, Phys. Lett. A **373** (2009), 2670–2674.
- [10] C.M. Bender, S. Kuzhel, *Unbounded \mathcal{C} -symmetries and their nonuniqueness*, J. Phys. A. **45** (2012), 444005–444019.

- [11] C.M. Bender, B. Tan, *Calculation of the hidden symmetry operator for a \mathcal{PT} -symmetric square well*, J. Phys. A. **39** (2006), 1945–1953.
- [12] B. Curgus, B. Najman, *The operator $(\operatorname{sgn}x)\frac{d^2}{dx^2}$ is similar to a selfadjoint operator in $L_2(\mathbb{R})$* , Proc. Amer. Math. Soc. **123** (1995), 1125–1128.
- [13] A. Gheondea, *Canonical forms of unbounded unitary operators in Krein space*, Publ. RIMS Kyoto Univ. **24** (1988), 205–224.
- [14] A. Grod, S. Kuzhel, V. Sudilovskaja, *On operators of transition in Krein spaces*, Opuscula Math. **31** (2011), 49–59.
- [15] V. Kostyrykin, K.A. Makarov, A.K. Motovilov, *Existence and uniqueness of solutions to the operator Riccati equation. A geometric approach*, Contemp. Math. **327** (2003), 181–198.
- [16] M.G. Krein, *Theory of self-adjoint extensions of semibounded operators and its applications I*. Math. Trans. **20** (1947), 431–495.
- [17] S. Kuzhel, *On pseudo-Hermitian operators with generalized \mathcal{C} -symmetries*, Operator Theory: Advances and Applications **190** (2009), 128–135.
- [18] H. Langer, *Maximal dual pairs of invariant subspaces of J -self-adjoint operators*, Mat. Zametki **7** (1970), 443–447 [in Russian].
- [19] H. Langer, *Spectral functions of definitizable operators in Krein spaces*, Functional Analysis Proceedings of a conference held at Dubrovnik, Yugoslavia, November 2–14, 1981, Lecture Notes in Mathematics, **948**, Springer-Verlag, Berlin, Heidelberg, New York, 1982, 1–46.
- [20] S.N. Naboko, *Conditions for similarity to unitary and self-adjoint operators*, Func. Anal. and Appl. **18** (1984) 1, 13–22.
- [21] M. Tomita, *Operators and operator algebras in Krein spaces, I Spectral Analysis in Pontrjagin spaces*, RIMS-Kokyuuroku **398** (1980), 131–158.

S. Kuzhel
kuzhel@agh.edu.pl

AGH University of Science and Technology
Faculty of Applied Mathematics
al. A. Mickiewicza 30, 30-059 Krakow, Poland

V. Sudilovskaya
veronica.sudi@gmail.com

Kyiv Vocational College
Kyiv, Ukraina

Received: February 28, 2016.

Revised: September 15, 2016.

Accepted: October 7, 2016.