

EXISTENCE RESULTS FOR A SUBLINEAR SECOND ORDER DIRICHLET BOUNDARY VALUE PROBLEM ON THE HALF-LINE

Dahmane Bouafia and Toufik Moussaoui

Communicated by Binlin Zhang

Abstract. In this paper we study the existence of nontrivial solutions for a boundary value problem on the half-line, where the nonlinear term is sublinear, by using Ekeland's variational principle and critical point theory.

Keywords: Ekeland's variational principle, critical point.

Mathematics Subject Classification: 34B40, 35A15, 35B38, 45C05, 34B24, 46T20.

1. INTRODUCTION

We consider the problem

$$\begin{cases} -(p(x)u'(x))' = \lambda q(x)f(x, u(x)), & x \in [0, +\infty), \\ u(0) = u(+\infty) = 0, \end{cases} \quad (1.1)$$

where $f : [0, +\infty) \times \mathbb{R} \rightarrow \mathbb{R}$ is a continuous function and λ is a positive parameter.

Boundary value problems with Dirichlet conditions are an area of the fast developing differential equations theory. Problems of this type arise in various fields of physics, biology, biotechnology etc. Thus, existence and multiplicity of solutions to boundary value problems on the half-line were studied by many authors. These results were obtained using upper and lower solution techniques, fixed point theory and topological degree theory, see for example, [6–9, 14] and [13, 16, 18, 19]. However, as far as we know, the study of solutions for second order boundary value problems, possibly singular on the infinite intervals via variational methods has received considerably less attention. For example, see [5] and [1, 2, 4, 12, 15]. We are interested here in the existence of solutions for a second order boundary value problem with the Sturm–Liouville operator

on the half-line. We take as a model a Dirichlet problem. Our results are obtained by applying Ekeland’s variational principle technique, and under sublinear condition of the nonlinearity $f \in C([0, +\infty) \times \mathbb{R}; \mathbb{R})$, with a positive parameter, the weighted q has a constant sign on the whole half-line, with $q > 0$ for $x \geq 0$. We give some new criteria to guarantee that the problem (1.1) has at least one classical solution. Moreover, we assume that the following conditions are satisfied:

(H1) there exist constants $a, b \in \mathbb{R}^+ \setminus \{0\}$ and $r \in (0, 1)$ such that

$$\forall x \in \mathbb{R}^+ \forall u \in \mathbb{R} : |f(x, u)| \leq a|u|^r + b,$$

(H2) $p : [0, +\infty) \rightarrow (0, +\infty)$ is continuously differentiable, $q : [0, +\infty) \rightarrow (0, +\infty)$ with $\frac{1}{p}, q \in L^1(0, +\infty)$, and

$$M_1 = \left(\int_0^{+\infty} \left(\int_x^{+\infty} \frac{ds}{p(s)} \right) dx \right)^{\frac{1}{r+1}} < +\infty,$$

$$M_2 = \left(\int_0^{+\infty} q(x) \left(\int_x^{+\infty} \frac{ds}{p(s)} \right)^{\frac{r+1}{2}} dx \right)^{\frac{1}{r+1}} < +\infty,$$

(H3) $f(x, 0) = 0$ and $\lim_{u \rightarrow 0^+} \frac{f(x, u)}{u^r} = +\infty$, uniformly for $x \in [0, +\infty)$.

Let the space $H_{0,p}^1(0, +\infty)$ defined by

$$H_{0,p}^1(0, +\infty) = \left\{ u \in AC([0, +\infty), \mathbb{R}) : \sqrt{p}u' \in L^2(0, +\infty), u(0) = u(+\infty) = 0 \right\}$$

be endowed with its natural norm

$$\|u\|_p = \left(\int_0^{+\infty} u^2(x) dx + \int_0^{+\infty} p(x)u'^2(x) dx \right)^{\frac{1}{2}}.$$

$H_{0,p}^1(0, +\infty)$ is a Hilbert space with the following inner product

$$\forall u, v \in H_{0,p}^1(0, +\infty) : (u, v) = \int_0^{+\infty} u(x)v(x) dx + \int_0^{+\infty} p(x)u'(x)v'(x) dx.$$

We define the weighted Lebesgue space

$$L_q^{r+1}(0, +\infty) = \left\{ u : (0, +\infty) \rightarrow \mathbb{R} \mid u \text{ is measurable and } \int_0^{+\infty} q(x)|u(x)|^{r+1} dx < +\infty \right\}$$

equipped with the usual norm

$$\|u\|_{r+1,q} = \left(\int_0^{+\infty} q(x)|u(x)|^{r+1} dx \right)^{\frac{1}{r+1}}.$$

Consider also the space

$$C_0[0, +\infty) = \left\{ u \in C([0, +\infty), \mathbb{R}) : \lim_{x \rightarrow +\infty} u(x) = 0 \right\}$$

endowed with the norm

$$\|u\|_\infty = \sup_{x \in [0, +\infty)} |u(x)|.$$

In our considerations we shall need the following corollary and lemmas.

Lemma 1.1 ([5]). *On $H_{0,p}^1(0, +\infty)$ the quantity $\|u\| = \left(\int_0^{+\infty} p(x)u'^2(x) dx \right)^{\frac{1}{2}}$ is a norm which is equivalent to the norm $\|u\|_p$, i.e.*

$$\forall u \in H_{0,p}^1(0, +\infty) : \|u\| \leq \|u\|_p \leq \sqrt{1 + M_1} \|u\|.$$

Lemma 1.2 ([5]). *$H_{0,p}^1(0, +\infty)$ is a reflexive Banach space.*

Lemma 1.3 ([5]). *$H_{0,p}^1(0, +\infty)$ embeds continuously in $C_0[0, +\infty)$ with*

$$\|u\|_\infty \leq \sqrt{\left\| \frac{1}{p} \right\|_{L^1}} \|u\|.$$

Corollary 1.4 ([5]). *$H_{0,p}^1(0, +\infty)$ is compactly embedded in $C_0[0, +\infty)$.*

Further we will also use the following lemmas and corollaries.

Lemma 1.5. *$C_0[0, +\infty)$ is continuously embedded in $L_q^{r+1}(0, +\infty)$.*

Proof. For any $u \in C_0[0, +\infty)$, we have

$$\|u\|_{r+1,q}^{r+1} = \int_0^{+\infty} q(x)|u(x)|^{r+1} dx \leq \sup_{x \in [0, +\infty)} |u(x)|^{r+1} \int_0^{+\infty} q(x) dx.$$

Consequently, we obtain $\|u\|_{r+1,q} \leq C \|u\|_\infty$, where $C = \|q\|_{L^1}^{\frac{1}{r+1}}$. □

Lemma 1.6. $H_{0,p}^1(0, +\infty)$ embeds continuously in $L_q^{r+1}[0, +\infty)$.

Proof. For all $u \in H_{0,p}^1(0, +\infty)$, we get

$$\begin{aligned} |u(x)|^{r+1} &= |u(+\infty) - u(x)|^{r+1} = \left| \int_x^{+\infty} u'(s) ds \right|^{r+1} = \left| \int_x^{+\infty} \sqrt{p(s)} u'(s) \frac{1}{\sqrt{p(s)}} ds \right|^{r+1} \\ &\leq \left(\int_x^{+\infty} p(s) u'^2(s) ds \right)^{\frac{r+1}{2}} \left(\int_x^{+\infty} \frac{1}{p(s)} ds \right)^{\frac{r+1}{2}}. \end{aligned}$$

Therefore, we obtain

$$\int_0^{+\infty} q(x) |u(x)|^{r+1} dx \leq \left(\int_0^{+\infty} p(s) u'^2(s) ds \right)^{\frac{r+1}{2}} \left(\int_0^{+\infty} q(x) \left(\int_x^{+\infty} \frac{1}{p(s)} ds \right)^{\frac{r+1}{2}} dx \right).$$

Thus, we have

$$\|u\|_{r+1,q} \leq M_2 \|u\|$$

which finishes the proof. □

Corollary 1.7. $H_{0,p}^1(0, +\infty)$ is compactly embedded in $L_q^{r+1}(0, +\infty)$, namely

$$H_{0,p}^1 \hookrightarrow C_0 \hookrightarrow L_q^{r+1}.$$

We are now concerned in the principal eigenvalue λ_1 of the nonlinear problem

$$\begin{cases} -(p(x)u'(x))' = \lambda q(x)|u(x)|^r, & x \geq 0, \\ u(0) = u(+\infty) = 0, \end{cases} \tag{1.2}$$

namely

$$\lambda_1 = \inf \left\{ \int_0^{+\infty} p(x) u'(x)^2 dx : u \in H_{0,p}^1 \setminus \{0\}, \int_0^{+\infty} q(x) |u(x)|^{r+1} dx = 1 \right\}.$$

Arguing as in [17, Proposition 3.2], one can easily establish the following lemma.

Lemma 1.8. The value λ_1 is positive and is achieved for some positive function $\varphi_1 \in H_{0,p}^1(0, +\infty) \setminus \{0\}$.

We need the following technique to prove our main result.

Theorem 1.9 (The weak Ekeland variational principle, [10]). *Let (E, d) be a complete metric space and let $J : E \rightarrow \mathbb{R}$ a functional that is lower semi-continuous, bounded from below. Then, for each $\varepsilon > 0$, there exists $u_\varepsilon \in E$ with*

$$J(u_\varepsilon) \leq \inf_E J + \varepsilon,$$

and whenever $w \in E$ with $w \neq u_\varepsilon$, then

$$J(u_\varepsilon) < J(w) + \varepsilon d(u_\varepsilon, w).$$

2. MAIN RESULT

To get the existence of solutions of problem (1.1), we consider the energy functional $J : H_{0,p}^1(0, +\infty) \rightarrow \mathbb{R}$ defined by

$$J(u) = \frac{1}{2} \|u\|^2 - \lambda \int_0^{+\infty} q(x)F(x, u(x))dx, \quad u \in H_{0,p}^1(0, +\infty),$$

where F denotes the primitive of f with respect to its second variable, i.e. $F(x, u) = \int_0^u f(x, s)ds$.

Proposition 2.1. *Suppose that the conditions (H1), (H2) hold. Then the functional J is Fréchet differentiable on $H_{0,p}^1(0, +\infty)$. Moreover, one has*

$$\forall v \in H_{0,p}^1(0, +\infty) : \langle J'(u), v \rangle = \int_0^{+\infty} p(x)u'(x)v'(x)dx - \lambda \int_0^{+\infty} q(x)f(x, u(x))v(x)dx.$$

Proof. First we show that J is Gâteaux-differentiable. Indeed, for all $v \in H_{0,p}^1(0, +\infty)$ and for each $t > 0$, we have

$$\begin{aligned} J(u + tv) - J(u) &= \frac{1}{2} \int_0^{+\infty} p(x)((u' + tv')(x))^2 dx - \lambda \int_0^{+\infty} q(x)F(x, (u + tv)(x))dx \\ &\quad - \frac{1}{2} \int_0^{+\infty} p(x)u'^2(x)dx + \lambda \int_0^{+\infty} q(x)F(x, u(x))dx \\ &= \frac{t^2}{2} \int_0^{+\infty} p(x)v'^2(x)dx + t \int_0^{+\infty} p(x)u'(x)v'(x)dx \\ &\quad - \lambda \int_0^{+\infty} q(x) \left[F(x, (u + tv)(x)) - F(x, u(x)) \right] dx \\ &= \frac{t^2}{2} \int_0^{+\infty} p(x)v'^2(x)dx + t \int_0^{+\infty} p(x)u'(x)v'(x)dx \\ &\quad - t\lambda \int_0^{+\infty} q(x)f(x, (u + t\theta v)(x))v(x)dx, \end{aligned}$$

where $0 < \theta < 1$.

Then

$$\begin{aligned} \frac{1}{t}[J(u + tv) - J(u)] &= \frac{t}{2} \int_0^{+\infty} p(x)v'^2(x)dx + \int_0^{+\infty} p(x)u'(x)v'(x)dx \\ &\quad - \lambda \int_0^{+\infty} q(x)f(x, (u + t\theta v)(x))v(x)dx. \end{aligned}$$

Let $t \rightarrow 0$, then, by using the Lebesgue dominated convergence theorem and (H1), we have

$$\forall v \in H_{0,p}^1(0, +\infty) : \langle J'(u), v \rangle = \int_0^{+\infty} p(x)u'(x)v'(x)dx - \lambda \int_0^{+\infty} q(x)f(x, u(x))v(x)dx.$$

Next, we show that J' is continuous. Indeed, let $(u_n) \subset H_{0,p}^1(0, +\infty)$ with $u_n \rightarrow u$ when $n \rightarrow +\infty$. Then there exists $R > 0$ such that $\|u_n\| \leq R$ for all $n \in \mathbb{N}$. Furthermore, from (H1), (H2) and Lemma 1.3 we derive that

$$\begin{aligned} q(x)|f(x, u_n(x))| &\leq aq(x)|u_n(x)|^r + bq(x) \\ &\leq a \sup_{x \in [0, +\infty)} |u_n(x)|^r q(x) + bq(x) \\ &= (a\|u_n\|_\infty^r + b)q(x) \\ &\leq \left(a \left(R \sqrt{\frac{1}{p}} \|u_n\|_{L^1} \right)^r + b \right) q(x) \in L^1(0, +\infty). \end{aligned}$$

Consequently, according to the Lebesgue dominated convergence theorem, we obtain

$$\lim_{n \rightarrow +\infty} \int_0^{+\infty} q(x)f(x, u_n(x))dx = \int_0^{+\infty} q(x)f(x, u(x))dx.$$

So, we have

$$\begin{aligned} \langle J'(u_n) - J'(u), v \rangle &= \int_0^{+\infty} p(x)u'_n(x)v'(x)dx - \lambda \int_0^{+\infty} q(x)f(x, u_n(x))v(x)dx \\ &\quad - \int_0^{+\infty} p(x)u'(x)v'(x)dx + \lambda \int_0^{+\infty} q(x)f(x, u(x))v(x)dx \\ &= \int_0^{+\infty} p(x)(u'_n(x) - u'(x))v'(x)dx \\ &\quad - \lambda \int_0^{+\infty} q(x)(f(x, u_n(x)) - f(x, u(x)))v(x)dx. \end{aligned}$$

Passing to the limit in $\langle J'(u_n) - J'(u), v \rangle$, when $n \rightarrow +\infty$, and taking into account that f is continuous, we obtain $J'(u_n) \rightarrow J'(u)$ as $n \rightarrow +\infty$. \square

Definition 2.2. We say that $u \in H_{0,p}^1(0, +\infty)$ is a weak solution of problem (1.1) if for every $v \in H_{0,p}^1(0, +\infty)$ we have

$$\langle J'(u), v \rangle = \int_0^{+\infty} p(x)u'(x)v'(x)dx - \lambda \int_0^{+\infty} q(x)f(x, u(x))v(x)dx = 0.$$

Remark 2.3. Since the nonlinear term f is continuous, then a weak solution of problem (1.1) is a classical solution.

Our main result is as follows.

Theorem 2.4. Assume (H1)–(H3) hold. Then there exists $\bar{\lambda} > 0$ such that problem (1.1) has at least one nontrivial solution u_λ for each $\lambda \in (0, \bar{\lambda})$.

Proof. From (H1) there exists $\delta_1 > 0$ such that

$$|F(x, u)| \leq a \frac{1}{r+1} |u|^{r+1} + b|u| \leq K|u|^{r+1} \quad \text{for all } |u| > \delta_1, \text{ for some } K > 0.$$

Also, from (H1), there exists $M_3 > 0$ such that

$$|F(x, u)| \leq M_3 \quad \text{for all } u \in [-\delta_1, \delta_1] \quad \text{and all } x \in (0, +\infty).$$

Therefore,

$$|F(x, u)| \leq M_3 + K|u|^{r+1} \quad \text{for all } u \in \mathbb{R} \text{ and all } x \in [0, +\infty). \tag{2.1}$$

From (2.1) and (H2) and by using the continuous embedding of $H_{0,p}^1(0, +\infty)$ in $L_q^{r+1}(0, +\infty)$ (i.e. note that $\|u\|_{r+1,q} \leq M_2\|u\|$) we get

$$\begin{aligned} J(u) &= \frac{1}{2}\|u\|^2 - \lambda \int_0^{+\infty} q(x)F(x, u(x))dx \\ &\geq \frac{1}{2}\|u\|^2 - \lambda \int_0^{+\infty} q(x) \left(M_3 + K|u|^{r+1}(x) \right) dx \\ &\geq \frac{1}{2}\|u\|^2 - \lambda M_3 \int_0^{+\infty} q(x)dx - \lambda K \int_0^{+\infty} q(x)|u|^{r+1}(x)dx \\ &= \frac{1}{2}\|u\|^2 - \lambda M_3 \int_0^{+\infty} q(x)dx - \lambda K \|u\|_{r+1,q}^{r+1} \\ &\geq \frac{1}{2}\|u\|^2 - \lambda K M_2^{r+1} \|u\|^{r+1} - \lambda M_3 \|q\|_{L^1}. \end{aligned}$$

Fix large enough $R > 0$. Then for $u \in H_{0,p}^1(0, +\infty)$ such that $\|u\| \geq R$ we have

$$J(u) \geq \frac{1}{2}\|u\|^2 - \lambda \left(KM_2^{r+1}\|u\|^{r+1} + M_3\|q\|_{L^1} \right).$$

Note that if $\lambda < \bar{\lambda} := \frac{R^2}{2(KM_2^{r+1}R^{r+1} + M_3\|q\|_{L^1})}$, this gives the existence of a real number $\rho > R$ satisfying

$$J(u) > 0 \quad \text{if } \|u\| = \rho, \quad \text{and} \quad \inf_{u \in \partial \bar{B}_\rho(0)} J(u) > 0, \tag{2.2}$$

and

$$J(u) \geq -C \quad \text{if } \|u\| \leq \rho, \quad \text{for some } C > 0.$$

Since J is a Fréchet differentiable functional, hence it is lower semicontinuous and bounded from below on $\bar{B}_\rho(0)$. We claim that

$$\inf_{u \in \bar{B}_\rho(0)} J(u) < 0. \tag{2.3}$$

Indeed, let $\varphi_1 \in H_{0,p}^1(0, +\infty)$ be the eigenfunction corresponding to the principal eigenvalue of problem (1.2) as described in Lemma 1.8. Then for any fixed λ in $(0, \bar{\lambda})$, by (H3), for any D ,

$$D > \frac{t^{1-r}\|\varphi_1\|^2}{2\lambda\|\varphi_1\|_{r+1,q}^{r+1}}, \tag{2.4}$$

there exists $0 < \epsilon_D < 1$ such that

$$f(x, u) \geq Du^r \quad \text{for } 0 < u < \epsilon_D.$$

Note that since $0 < u < 1$, therefore

$$f(x, u) \geq Du^r \quad \text{implies} \quad F(x, u) \geq Du^{r+1}. \tag{2.5}$$

Since the function φ_1 is continuous on $[0, +\infty)$ and $\varphi_1(0) = \varphi_1(+\infty) = 0$, there exists $\hat{c} > 0$ such that $\sup_{x \in [0, +\infty)} \varphi_1(x) \leq \hat{c}$. Thus, for every $0 < t < 1/\hat{c}$ (t near 0), owing to (2.4), (2.5) and Lemma 1.8 we have

$$\begin{aligned} J(t\varphi_1) &= \frac{t^2}{2}\|\varphi_1\|^2 - \lambda \int_0^{+\infty} q(x)F(x, t\varphi_1(x))dx \\ &\leq \frac{t^2}{2}\|\varphi_1\|^2 - \lambda Dt^{r+1} \int_0^{+\infty} q(x)\varphi_1^{r+1}(x)dx \\ &= \frac{t^2}{2}\|\varphi_1\|^2 - \lambda Dt^{r+1}\|\varphi_1\|_{r+1,q}^{r+1} < 0. \end{aligned}$$

So (2.3) is proved.

From (2.2) together with (2.3) we obtain

$$\inf_{u \in \overline{B}_\rho(0)} J(u) < 0 < \inf_{u \in \partial \overline{B}_\rho(0)} J(u).$$

We define a distance on $\overline{B}_\rho(0)$ as follows:

$$d(u, v) = \|u - v\| \quad \text{for } u, v \in \overline{B}_\rho(0).$$

Clearly, $\overline{B}_\rho(0)$ is a complete metric space. It is known that $J \in C^1(\overline{B}_\rho(0), \mathbb{R})$, therefore J is lower semicontinuous and bounded from below on $\overline{B}_\rho(0)$. We put

$$c_\lambda = \inf\{J(u) : u \in \overline{B}_\rho(0)\}.$$

By the Ekeland variational principle (see Theorem 1.9) in $\overline{B}_\rho(0)$, there is a minimizing sequence $(u_n) \subset \overline{B}_\rho(0)$ for all $n \in \mathbb{N} \setminus \{0\}$ such that

$$c_\lambda < J(u_n) \leq \inf_{u \in \overline{B}_\rho(0)} J(u) + \frac{1}{n} \leq c_\lambda + \frac{1}{n}, \tag{2.6}$$

$$\forall w \in \overline{B}_\rho(0) : J(u_n) \leq J(w) + \frac{1}{n} \|w - u_n\|. \tag{2.7}$$

If we put $w = u_n + th$ in (2.7) for $t > 0$, $h \in H_{0,p}^1(0, +\infty)$ and $n \in \mathbb{N} \setminus \{0\}$, then we get $J(u_n) \leq J(u_n + th) + \frac{1}{n}t\|h\|$. Thus, we have

$$\frac{1}{t}[J(u_n) - J(u_n + th)] < \frac{1}{n}\|h\|,$$

and taking into account that J is a Fréchet differentiable functional we see that

$$-\langle J'(u_n), h \rangle \leq \frac{1}{n}\|h\| \quad \text{for all } n \in \mathbb{N} \setminus \{0\},$$

Similarly, if we put $w = u_n - th$, then we get

$$\langle J'(u_n), h \rangle \leq \frac{1}{n}\|h\| \quad \text{for all } n \in \mathbb{N} \setminus \{0\}.$$

So

$$\sup_{\|h\| \leq 1} |\langle J'(u_n), h \rangle| \leq \frac{1}{n} \quad \text{for all } n \in \mathbb{N} \setminus \{0\}.$$

Therefore, we have

$$\|J'(u_n)\| \rightarrow 0 \quad \text{and} \quad J(u_n) \rightarrow c_\lambda \quad \text{as } n \rightarrow +\infty,$$

where c_λ stands for the infimum of $J(u)$ on $\overline{B}_\rho(0)$. From the above discussion, we know that (u_n) is a bounded $(P.S)_{c_\lambda}$ sequence (see [11, Definition 2.3]), and $\overline{B}_\rho(0)$ is a closed

convex set, and from Corollary 1.7, then there exist $u_\lambda \in \overline{B}_\rho(0) \subset H^1_{0,p}(0, +\infty)$ and a convergent subsequence still denoted by (u_n) , such that

$$\begin{cases} u_n \rightharpoonup u_\lambda, & \text{weakly in } H^1_{0,p}(0, +\infty), \\ u_n(x) \rightarrow u_\lambda(x), & \text{for a.e. in } (0, +\infty), \\ u_n \rightarrow u_\lambda, & \text{strongly in } L^{r+1}_q(0, +\infty). \end{cases}$$

Consequently, passing to the limit in $\langle J'(u_n), v \rangle$ as $n \rightarrow +\infty$, we deduce that

$$\int_0^{+\infty} p(x)u'_\lambda(x)v'(x)dx - \lambda \int_0^{+\infty} q(x)f(x, u_\lambda(x))v(x)dx = 0,$$

for all $v \in H^1_{0,p}(0, +\infty)$, namely $\langle J'(u_\lambda), v \rangle = 0$ for all $v \in H^1_{0,p}(0, +\infty)$.

Now it remains to show that $J(u_\lambda) = c_\lambda$. Indeed, by (2.1) and Lemma 1.3, for all $x \in [0, +\infty)$ and $n \in \mathbb{N} \setminus \{0\}$, we obtain

$$\begin{aligned} q(x)|F(x, u_n(x))| &\leq M_3q(x) + Kq(x)|u_n(x)|^{r+1} \\ &\leq M_3q(x) + Kq(x) \sup_{x \in [0, +\infty)} |u_n(x)|^{r+1} \\ &\leq M_3q(x) + Kq(x)\|u_n\|_\infty^{r+1} \\ &\leq \left(M_3 + K \left(\rho \sqrt{\left\| \frac{1}{p} \right\|_{L^1}} \right)^{r+1} \right) q(x) \in L^1(0, +\infty). \end{aligned}$$

Thus, by the Lebesgue dominated convergence theorem and (2.6), we conclude that

$$c_\lambda \leq \lim_{n \rightarrow +\infty} J(u_n) = J(u_\lambda) \leq c_\lambda.$$

Hence, c_λ is a critical value of the functional J at the point u_λ in $H^1_{0,p}(0, +\infty)$. □

REFERENCES


[1] G.A. Afrouzi, A. Hadjian, V.D. Rădulescu, *Variational analysis for Dirichlet impulsive differential equations with oscillatory nonlinearity*, Portugal. Math. (N.S.) **70**, Fasc. 3, (2013), 225–242.

[2] K. Ait-Mahiout, S. Djebali, T. Moussaoui, *Multiple solutions for a BVP on $(0, +\infty)$ via Morse theory and $H^1_{0,p}(\mathbb{R}^+)$ versus $C^1_p(\mathbb{R}^+)$ local minimizers*, Arab. J. Math. (2016), 5: 9–22.

[3] M. Badiale, E. Serra, *Semilinear Elliptic Equations for Beginners*, Springer, New York, 2011.

[4] D. Bouafia, T. Moussaoui, D. O'Regan, *Existence of solutions for a second order problem on the half-line via Ekeland's variational principle*, Discuss. Math. Differ. Incl. Control Optim. **36** (2016), 131–140.

- [5] M. Briki, S. Djebali, T. Moussaoui, *Solvability of an Impulsive Boundary Value Problem on The Half-Line Via Critical Point Theory*, Bull. Iranian Math. Soc. **43** (2017) 3, 601–615.
- [6] S. Djebali, T. Moussaoui, *A class of second order BVPs on infinite intervals*, Electron. J. Qual. Theory Differ. Equ. **4** (2006), 1–19.
- [7] S. Djebali, S. Zahar, *Bounded solutions for a derivative dependent boundary value problem on the half-line*, Dynam. Systems Appl. **19** (2010), 545–556.
- [8] S. Djebali, O. Saifi, S. Zahar, *Upper and lower solutions for BVPs on the half-line with variable coefficient and derivative depending nonlinearity*, Electron. J. Qual. Theory Differ. Equ. (2011), no. 14, 1–18.
- [9] S. Djebali, O. Saifi, S. Zahar, *Singular boundary value problems with variable coefficients on the positive half-line*, Electron. J. Differential Equations **2013** (2013), no. 73, 1–18.
- [10] I. Ekeland, *On the variational principle*, J. Math. Anal. Appl. **47** (1974), 324–353.
- [11] Y. Jabri, *The Mountain Pass Theorem, Variants, Generalizations and Some Applications*, Cambridge University Press, New York, 2003.
- [12] Y. Liu, *Existence and unboundedness of positive solutions for singular boundary value problems on half-line*, Appl. Math. Comput. **144** (2003), 543–556.
- [13] H. Lian, W. Ge, *Existence of positive solutions for Sturm–Liouville boundary value problems on the half-line*, J. Math. Anal. Appl. **321** (2006), 781–792.
- [14] H. Lian, W. Ge, *Solvability for second-order three-point boundary value problems on a half-line*, Appl. Math. Lett. **19** (2006), 1000–1006.
- [15] H. Lian, P. Wang, W. Ge, *Unbounded upper and lower solutions method for Sturm–Liouville boundary value problem on infinite intervals*, Nonlinear Anal. **70** (2009), 2627–2633.
- [16] R. Ma, *Positive solutions for second order three-point boundary value problems*, Appl. Math. Lett. **14** (2001) 1, 1–5.
- [17] K. Perera, Z. Zhang, *Nontrivial solutions of Kirchhoff-type problems via the Yang index*, J. Differential Equations **221** (2006), 246–255.
- [18] Y. Tian, W. Ge, W. Shan, *Positive solutions for three-point boundary value problem on the half-line*, Comput. Math. Appl. **53** (2007), 1029–1039.
- [19] B. Yan, D. O’Regan, R. Agarwal, *Unbounded solutions for singular boundary value problems on the semi-infinite interval: Upper and lower solutions and multiplicity*, J. Comput. Appl. Math. **197** (2006), 365–386.

Dahmane Bouafia (corresponding author)
dahmane.bouafia@univ-msila.dz
 <https://orcid.org/0000-0002-4471-9338>

University of M’sila
Department of Mathematics
M’sila, Algeria

Toufik Moussaoui
moussaoui@ens-kouba.dz

Laboratory of Fixed Point Theory and Applications
Department of Mathematics
E.N.S. Kouba, Algiers, Algeria

Received: October 26, 2019.

Revised: August 17, 2020.

Accepted: August 19, 2020.