

SINGULAR ELLIPTIC PROBLEMS WITH DIRICHLET OR MIXED DIRICHLET–NEUMANN NON-HOMOGENEOUS BOUNDARY CONDITIONS

Tomas Godoy

Communicated by Vicentiu D. Rădulescu

Abstract. Let Ω be a C^2 bounded domain in \mathbb{R}^n such that $\partial\Omega = \Gamma_1 \cup \Gamma_2$, where Γ_1 and Γ_2 are disjoint closed subsets of $\partial\Omega$, and consider the problem $-\Delta u = g(\cdot, u)$ in Ω , $u = \tau$ on Γ_1 , $\frac{\partial u}{\partial \nu} = \eta$ on Γ_2 , where $0 \leq \tau \in W^{\frac{1}{2},2}(\Gamma_1)$, $\eta \in (H^1_{0,\Gamma_1}(\Omega))'$, and $g : \Omega \times (0, \infty) \rightarrow \mathbb{R}$ is a nonnegative Carathéodory function. Under suitable assumptions on g and η we prove the existence and uniqueness of a positive weak solution of this problem. Our assumptions allow g to be singular at $s = 0$ and also at $x \in S$ for some suitable subsets $S \subset \bar{\Omega}$. The Dirichlet problem $-\Delta u = g(\cdot, u)$ in Ω , $u = \sigma$ on $\partial\Omega$ is also studied in the case when $0 \leq \sigma \in W^{\frac{1}{2},2}(\Omega)$.

Keywords: singular elliptic problems, mixed boundary conditions, weak solutions.

Mathematics Subject Classification: 35J75, 35M12, 35D30.

1. INTRODUCTION AND STATEMENT OF THE MAIN RESULTS

Let Ω be a C^2 and bounded domain in \mathbb{R}^n such that $\partial\Omega = \Gamma_1 \cup \Gamma_2$, with Γ_1 and Γ_2 disjoint closed subsets of $\partial\Omega$. Our aim in this paper is to state existence and uniqueness results for weak solutions $u \in H^1(\Omega)$ of possibly singular elliptic Dirichlet problems of the form

$$\begin{cases} -\Delta u = g(\cdot, u) & \text{in } \Omega, \\ u = \sigma & \text{on } \partial\Omega, \\ u > 0 & \text{in } \Omega, \end{cases} \quad (1.1)$$

as well as of possibly singular problems with mixed boundary conditions of the form

$$\begin{cases} -\Delta u = g(\cdot, u) & \text{in } \Omega, \\ u = \tau & \text{on } \Gamma_1, \\ \frac{\partial u}{\partial \nu} = \eta & \text{on } \Gamma_2, \\ u > 0 & \text{in } \Omega, \end{cases} \quad (1.2)$$

where $g : \Omega \times (0, \infty) \rightarrow [0, \infty)$ is a suitable nonnegative Carathéodory function $g(x, s)$ which may be singular at $s = 0$ and at $x \in S$ for some suitable subsets $S \subset \bar{\Omega}$, and, in problem (1.1), $\sigma \in H^{\frac{1}{2}}(\partial\Omega)$, whereas in problem (1.2), $\tau \in H^{\frac{1}{2}}(\Gamma_1)$ and η is a suitable function defined on Γ_2 .

Singular elliptic problems appear in the study of many nonlinear physical phenomena: thin films of viscous fluids, chemical catalysis, non-Newtonian fluids, temperature of some electrical conductors, response of a membrane cap under heavy loads, Van der Waal forces, as well as in the study of micro electro-mechanical devices (see, e.g., [8, 12, 18, 20, 21, 30, 34] and the references therein).

In [13], problem (1.1) was studied in the case $\sigma = 0$ (i.e., with homogeneous Dirichlet boundary condition) and there it was proved that if $g \in C^1(\bar{\Omega} \times (0, \infty))$ satisfies that $g(x, \cdot)$ is nonincreasing on $(0, \infty)$ for any $x \in \bar{\Omega}$ and $\lim_{s \rightarrow 0^+} g(x, s) = \infty$ uniformly on $\bar{\Omega}$, then (1.1) has a unique classical solutions $u \in C^2(\Omega) \cap C(\bar{\Omega})$.

In [21, 46], and [45], problem (1.1) was addressed when $\sigma \neq 0$ (non homogeneous Dirichlet boundary condition) obtaining, again in this case, existence and uniqueness of classical solutions when σ is regular enough.

In [11], existence and nonexistence results were obtained for classical solutions of singular bifurcation problems whose model problem is $-\Delta u = u^{-\alpha} + \lambda u^p$ in Ω , $u = 0$ on $\partial\Omega$, $u > 0$ in Ω , where $\alpha > 0$, $\lambda > 0$, and $p > 1$, and there it was proved that there exists $\lambda^* \in (0, \infty)$ such that for $\lambda < \lambda^*$ there exists at least a solution and for $\lambda > \lambda^*$ no such a solution exists. In [18] it was studied the problem with a parameter $-\Delta u = \lambda f - u^{-\alpha}$ in Ω , $u = 0$ on $\partial\Omega$, $u > 0$ in Ω , $u^{-\alpha} \in L^1(\Omega)$, where $\lambda > 0$, $0 < \alpha < 1$, and $0 \leq f \in L^1(\Omega)$. It turns out that the situation is the opposite of that in [11]: there exists $\lambda^* \in (0, \infty)$ such that for $\lambda > \lambda^*$ there exists at least a solution and for $\lambda < \lambda^*$ no such a solution exists.

In [31] it was studied the model problem

$$-\Delta u = k(x)u^{-\alpha} \text{ in } \Omega, \quad u = 0 \text{ on } \partial\Omega, \quad u > 0 \text{ in } \Omega. \quad (1.3)$$

There it was proved that if Ω is a $C^{2+\beta}$ bounded domain for some $\beta \in (0, 1)$, and $k \in C^\beta(\bar{\Omega})$ satisfies $\min_{\bar{\Omega}} k > 0$ then, for any $\alpha > 0$, problem (1.3) has a unique classical solution $u \in C^{2+\beta}(\Omega) \cap C(\bar{\Omega})$ which belongs to $C^1(\bar{\Omega})$ if $\alpha < 1$, and belongs to $H_0^1(\Omega)$ if and only if $\alpha < 3$. Moreover, if $\alpha > 1$ then $\frac{1}{c}\varphi_1^{\frac{2}{1+\alpha}} \leq u \leq c\varphi_1^{\frac{2}{1+\alpha}}$ in Ω , where c is a positive constant and φ_1 is a positive eigenfunction corresponding to the first eigenvalue for $-\Delta$ on Ω with homogeneous Dirichlet boundary condition.

After [31], several works studied problem (1.3) under weaker regularity assumptions on k and, in some of them, for more general differential operators than the Laplacian, as well as for more general nonlinearities.

In [15] it was stated the existence and uniqueness of a weak solution $u \in H_0^1(\Omega)$ of problem (1.3) in the case when k is a nonnegative and nonidentically zero function in $L^\infty(\Omega)$, and, for such a u , a global bound for ∇u was obtained. Let us mention some of them.

In [47] it was proved, among other results, that if $\alpha > 1$, $k \in L^1(\Omega)$ and $k > 0$ a.e. in Ω , then (1.3) has a weak solution $u \in H_0^1(\Omega)$ if and only if there exists $u_0 \in H_0^1(\Omega)$

such that $\int_{\Omega} k u_0^{1-\alpha} < \infty$. These results were extended in [32] to the case where the Laplacian is replaced by the p -Laplacian operator.

Singular problems for differential operators (including the p -Laplacian) more general than the Laplacian and/or with more general nonlinearities were also studied in [2, 22, 32, 37, 39, 48] and [40].

Singular problems on punctured domains were studied in [3]. The paper [9] addressed problem (1.1) in the case where $\alpha = \alpha(x)$ (variable exponent). In [5], [28, 33] and [17] it was studied the existence of solutions (either classical or weak or very weak) of (1.3) in the case where k behaves like $(\text{dist}(\cdot, \partial\Omega))^{-\beta}$ for some $\beta > 0$, and in [36] it was considered the case where k is either a nonnegative function in $L^1(\Omega)$ or a bounded Radon measure on Ω .

In [44] existence and nonexistence results were given for the problem with a parameter $-\Delta u = k(x)u^{-\alpha} + \lambda u^p$ in Ω , $u = 0$ on $\partial\Omega$, $u > 0$ in Ω in the case where $\alpha, p \in (0, 1)$, and k may change sign.

Existence results for classical solutions of Lane–Emden–Fowler equations with convection and singular potential were obtained in [19], and related problems were studied in [10, 25] and [4].

Let us mention also that in [30] it was studied the existence of positive classical solutions of the one-dimensional singular problem

$$-u''(t) = f(t)u^{-\beta}(t) + h(t) \quad \text{on } (0, 1), \quad (1.4)$$

where $\beta > 0$, f and h belong to $C(0, 1)$, $f > 0$ in $(0, 1)$, and

$$\int_0^1 t(1-t)(f(t) + |h(t)|) < \infty,$$

and with u such that one of the following boundary conditions holds:

$$u(0) = a, \quad u(1) = b, \quad (1.5)$$

$$u(0) = a, \quad u'(1) = c. \quad (1.6)$$

In [30, Theorem 1.1] it was proved that, if $a \geq 0$ and $b \geq 0$, then problem (1.4), with boundary conditions (1.5), has a unique classical solution; and in [30, Theorem 1.2] it was proved that problem (1.4), with boundary conditions (1.6), has a unique positive solution if $c > c_0 := \inf\{u'_\xi(1) : \xi > 0\}$, and has no positive solution if $c < c_0$, where, for $\xi > 0$, u_ξ is the solution, provided by [30, Theorem 1.1], of problem (1.4) with boundary conditions $u(0) = a$, $u(1) = \xi$.

The interested reader will find an updated account, concerning the topic of singular elliptic Dirichlet problems, as well as additional references, in the research books [23, 24, 41]. See also [16].

As said before, we are interested in the existence and uniqueness of weak solutions of problems (1.1) and (1.2). More specifically, our interest is to obtain a sort of n -dimensional analogues of the above quoted (Theorems 1.1 and 1.2 of [30]).

For a function $u \in H^1(\Omega)$ the value of u on $\partial\Omega$ (or on Γ_1 , or on Γ_2) will be always understood in the sense of the trace. Let us present the notion of weak solutions of Dirichlet problems we use.

Definition 1.1. Let $f : \Omega \rightarrow \mathbb{R}$ be such that $f\varphi \in L^1(\Omega)$ for any $\varphi \in H_0^1(\Omega)$, and let $\sigma : \partial\Omega \rightarrow \mathbb{R}$. We say that $u : \Omega \rightarrow \mathbb{R}$ is a weak solution of the problem

$$\begin{cases} -\Delta u = f & \text{in } \Omega, \\ u = \sigma & \text{on } \partial\Omega \end{cases} \quad (1.7)$$

if $u \in H^1(\Omega)$, $u = \sigma$ on $\partial\Omega$, and

$$\int_{\Omega} \langle \nabla u, \nabla \varphi \rangle = \int_{\Omega} f\varphi \quad \text{for any } \varphi \in H_0^1(\Omega). \quad (1.8)$$

For a function $f : \Omega \rightarrow \mathbb{R}$, we will write $f \in (H_0^1(\Omega))'$ to mean that $f\varphi \in L^1(\Omega)$ for any $\varphi \in H_0^1(\Omega)$, and that there exists a positive constant c such that $|\int_{\Omega} f\varphi| \leq c\|\varphi\|_{H_0^1(\Omega)}$ for any $\varphi \in H_0^1(\Omega)$.

Remark 1.2. If $f \in (H_0^1(\Omega))'$ and $\sigma \in H^{\frac{1}{2}}(\partial\Omega)$, then problem (1.7) has a unique weak solution $u \in H^1(\Omega)$, and there exists a positive constant c , independent of f and σ , such that

$$\|u\|_{H^1(\Omega)} \leq c \left(\|f\|_{(H_0^1(\Omega))'} + \|\sigma\|_{H^{\frac{1}{2}}(\partial\Omega)} \right),$$

for a proof of this fact see, e.g., [43, Section 8.4.1] (there it is assumed that $f \in L^2(\Omega)$, but the arguments given there works also when $f \in (H_0^1(\Omega))'$).

For $S \subset \overline{\Omega}$ we will denote by ρ_S the distance function defined by

$$\rho_S(x) := \text{dist}(x, S) \quad \text{for } x \in \Omega,$$

and, for a Lebesgue measurable subset E of Ω , $|E|$ will denote the Lebesgue measure of E .

We recall that a function $g : \Omega \times (0, \infty) \rightarrow \mathbb{R}$ is called a Carathéodory function if $g(\cdot, s)$ is Lebesgue measurable for any $s \in (0, \infty)$ and $g(x, \cdot)$ is continuous on $(0, \infty)$ for a.e. $x \in \Omega$. Our first result, concerning problem (1.1), reads as follows:

Theorem 1.3. *Let Ω be a C^2 and bounded domain in \mathbb{R}^n . Let $g : \Omega \times (0, \infty) \rightarrow [0, \infty)$ be a function satisfy the following three conditions:*

- (H1) $g : \Omega \times (0, \infty) \rightarrow \mathbb{R}$ is a nonnegative Carathéodory function such that, for each $x \in \Omega$, $g(x, \cdot)$ is nonincreasing on $(0, \infty)$.
- (H2) There exists a Lebesgue measurable subset E of Ω such that $|E| > 0$ and $g(x, s) > 0$ for any $s > 0$ and almost all $x \in E$.
- (H3) $\rho_{\partial\Omega}g(\cdot, c\rho_{\partial\Omega}) \in L^2(\Omega)$ for any $c \in (0, \infty)$.

Then for any nonnegative $\sigma \in H^{\frac{1}{2}}(\partial\Omega)$ problem (1.1) has a unique weak solution $u \in H^1(\Omega)$ and there exists a positive constant c such that $u \geq c\rho_{\partial\Omega}$ a.e. in Ω .

Let us introduce the space

$$H_{0,\Gamma_1}^1(\Omega) := \{v \in H^1(\Omega) : v = 0 \text{ on } \Gamma_1\},$$

which endowed with the inner product of $H^1(\Omega)$ is a Hilbert space. Let $(H_{0,\Gamma_1}^1(\Omega))'$ denote its topological dual. If f is a function defined on Ω , we will write $f \in (H_{0,\Gamma_1}^1(\Omega))'$ to mean that $f\varphi \in L^1(\Omega)$ and $\left| \int_{\Gamma_2} \eta\varphi \right| \leq c\|\varphi\|_{H^1(\Omega)}$ for any $\varphi \in H_{0,\Gamma_1}^1(\Omega)$, with c a positive constant independent of φ . Similarly, if η is a function defined on Γ_2 we will say that $\eta \in (H_{0,\Gamma_1}^1(\Omega))'$ to mean that $\eta\varphi \in L^1(\Gamma_2)$ and that $\left| \int_{\Gamma_2} \eta\varphi \right| \leq c\|\varphi\|_{H^1(\Omega)}$ for any $\varphi \in H_{0,\Gamma_1}^1(\Omega)$, with a positive constant c independent of φ . In both cases, the maps $\varphi \rightarrow \int_{\Omega} f\varphi$ and $\varphi \rightarrow \int_{\Gamma_2} \eta\varphi$ will still be denoted by f and η , respectively.

Weak solutions of problems with mixed nonhomogeneous Dirichlet–Neumann boundary conditions are defined as follows:

Definition 1.4. Let $f : \Omega \rightarrow \mathbb{R}$ be such that $f\varphi \in L^1(\Omega)$ for any $\varphi \in H_{0,\Gamma_1}^1(\Omega)$, let $\tau \in H^{\frac{1}{2}}(\Gamma_1)$, and let $\eta : \Gamma_2 \rightarrow \mathbb{R}$ be a measurable function such that $\eta\varphi \in L^1(\Gamma_2)$ for any $\varphi \in H_{0,\Gamma_1}^1(\Omega)$. We say that $u : \Omega \rightarrow \mathbb{R}$ is a weak solution of the problem

$$\begin{cases} -\Delta u = f & \text{in } \Omega, \\ u = \tau & \text{on } \Gamma_1, \\ \frac{\partial u}{\partial \nu} = \eta & \text{on } \Gamma_2. \end{cases} \quad (1.9)$$

if $u \in H^1(\Omega)$, $u = \tau$ on Γ_1 , and

$$\int_{\Omega} \langle \nabla u, \nabla \varphi \rangle = \int_{\Omega} f\varphi + \int_{\Gamma_2} \eta\varphi \quad \text{for any } \varphi \in H_{0,\Gamma_1}^1(\Omega). \quad (1.10)$$

Let $f : \Omega \rightarrow \mathbb{R}$ be such that $f\varphi \in L^1(\Omega)$ for any $\varphi \in H_{0,\Gamma_1}^1(\Omega)$, let $\tau : \Gamma_1 \rightarrow \mathbb{R}$, and suppose that u is a weak solution of the problem

$$\begin{cases} -\Delta u = f & \text{in } \Omega, \\ u = \tau & \text{on } \Gamma_1, \\ u = 0 & \text{on } \Gamma_2. \end{cases} \quad (1.11)$$

If $\varphi \in H_{0,\Gamma_1}^1(\Omega)$ and if φ and u are regular enough on $\overline{\Omega}$, we have

$$-\operatorname{div}(\varphi \nabla u) + \langle \nabla u, \nabla \varphi \rangle = f\varphi$$

and then, from the divergence theorem and the fact that $\varphi = 0$ on Γ_1 , we get

$$-\int_{\Gamma_2} \frac{\partial u}{\partial \nu} \varphi + \int_{\Omega} \langle \nabla u, \nabla \varphi \rangle = \int_{\Omega} f\varphi.$$

Therefore,

$$\int_{\Gamma_2} \frac{\partial u}{\partial \nu} \varphi = \int_{\Omega} \langle \nabla u, \nabla \varphi \rangle - \int_{\Omega} f\varphi.$$

This suggests the following definition.

Definition 1.5. Let $f : \Omega \rightarrow \mathbb{R}$ be such that $f \in (H_{0,\Gamma_1}^1(\Omega))'$, and let $\tau \in H^{\frac{1}{2}}(\Gamma_1)$. If $u \in H^1(\Omega)$ is the weak solution of problem (1.11), we define the (distributional) normal derivative of u on Γ_2 , as the linear functional $\frac{\partial u}{\partial \nu_{\Gamma_2}} : H_{0,\Gamma_1}^1(\Omega) \rightarrow \mathbb{R}$ defined by

$$\frac{\partial u}{\partial \nu_{\Gamma_2}}(\varphi) := \int_{\Omega} \langle \nabla u, \nabla \varphi \rangle - \int_{\Omega} f \varphi \quad \text{for any } \varphi \in H_{0,\Gamma_1}^1(\Omega). \quad (1.12)$$

For η and $\tilde{\eta}$ in $(H_{0,\Gamma_1}^1(\Omega))'$ we will write $\eta \geq \tilde{\eta}$ (respectively $\eta \leq \tilde{\eta}$) to mean that $\eta(\varphi) \geq \tilde{\eta}(\varphi)$ (resp. $\eta(\varphi) \leq \tilde{\eta}(\varphi)$) for any nonnegative $\varphi \in H_{0,\Gamma_1}^1(\Omega)$. We will write also $\eta > \tilde{\eta}$ (respectively $\eta < \tilde{\eta}$) to mean that $\eta \neq \tilde{\eta}$ and $\eta \geq \tilde{\eta}$ (resp. $\eta \leq \tilde{\eta}$).

Concerning problem (1.2) we have the following:

Theorem 1.6. Let Ω be a C^2 and bounded domain in \mathbb{R}^n such that $\partial\Omega = \Gamma_1 \cup \Gamma_2$, where Γ_1 and Γ_2 are disjoint closed sets in $\partial\Omega$. Let $g : \Omega \times (0, \infty) \rightarrow [0, \infty)$. Assume the conditions (H1)–(H2) of Theorem 1.3 and the following:

(H3') $\rho_{\Gamma_1} g(\cdot, c\rho_{\partial\Omega}) \in L^2(\Omega)$ for any $c \in (0, \infty)$.

Let τ be a nonnegative function in $H^{\frac{1}{2}}(\Gamma_1)$, let u_τ be the weak solution of the problem

$$\begin{cases} -\Delta u_\tau = g(\cdot, u_\tau) & \text{in } \Omega, \\ u_\tau = \tau & \text{on } \Gamma_1, \\ u_\tau = 0 & \text{on } \Gamma_2. \end{cases} \quad (1.13)$$

given by Theorem 1.3, and let $\eta : \Gamma_2 \rightarrow \mathbb{R}$ be such that $\eta \in (H_{0,\Gamma_1}^1(\Omega))'$. Then:

- (i) if $\eta \geq \frac{\partial u_\tau}{\partial \nu_{\Gamma_2}}$, then (1.2) has a unique weak solution $u \in H^1(\Omega)$, and there exists a positive constant c such that $u \geq c\rho_{\partial\Omega}$ in Ω ,
- (ii) if $\eta < \frac{\partial u_\tau}{\partial \nu_{\Gamma_2}}$, then (1.2) has no weak solutions.

As a consequence of Theorem 1.6 and of a weak form of the Hopf boundary lemma given in Lemma 4.4, we will get the following:

Corollary 1.7. Let $g : \Omega \times (0, \infty) \rightarrow [0, \infty)$ satisfy the conditions (H1)–(H2) of Theorem 1.3 and the condition (H3') of Theorem 1.6, let τ be a nonnegative function in $H^{\frac{1}{2}}(\Gamma_1)$ and let $\eta : \Gamma_2 \rightarrow \mathbb{R}$ be such that $\eta \in (H_{0,\Gamma_1}^1(\Omega))'$. If $\eta \geq 0$, then problem (1.2) has a unique weak solution $u \in H^1(\Omega)$, and there exists a positive constant c such that $u \geq c\rho_{\partial\Omega}$ in Ω .

The paper is organized as follows. In Section 2 we recall some general facts we need, and in Section 3 we study problem (1.1) via an approximation approach, which is adapted from [27], where the existence and uniqueness of strong solutions of (1.1) were investigated. We consider, for $\varepsilon \in (0, 1]$ and for any nonnegative $\sigma \in H^{\frac{1}{2}}(\partial\Omega)$, the problem of finding $v_\varepsilon \in H_0^1(\Omega)$ such that $-\Delta v_\varepsilon = g_\varepsilon(\cdot, v_\varepsilon + \tilde{\sigma})$ in Ω , $v_\varepsilon = 0$ on $\partial\Omega$, where $\tilde{\sigma}$ is the solution of the problem $-\Delta \tilde{\sigma} = 0$ in Ω , $\tilde{\sigma} = \sigma$ on $\partial\Omega$, and with $g_\varepsilon : \Omega \times (0, \infty) \rightarrow \mathbb{R}$ defined by $g_\varepsilon(x, s) := \min \{\varepsilon^{-1}, g(x, s + \varepsilon)\}$. By writing the above problem for v_ε as a fixed point problem and using the Schauder fixed

point theorem we prove in Lemma 3.2 the existence and uniqueness of such a v_ε , and that, in addition, the map $\varepsilon \rightarrow v_\varepsilon$ is nonincreasing. Lemma 3.3 shows that if $v(x) := \lim_{\varepsilon \rightarrow 0^+} v_\varepsilon(x)$, then $v \in H_0^1(\Omega)$,

$$\lim_{\varepsilon \rightarrow 0^+} \|v_\varepsilon - v\|_{H_0^1(\Omega)} = 0 \quad \text{and} \quad \lim_{\varepsilon \rightarrow 0^+} \|g_\varepsilon(\cdot, v_\varepsilon + \tilde{\sigma}) - g(\cdot, v + \tilde{\sigma})\|_{L^2(\Omega, \rho_{\partial\Omega}^2(x)dx)} = 0.$$

From these facts and from other additional considerations, Theorem 1.3 is proved at the end of Section 3, by showing that $u := v + \tilde{\sigma}$ is the unique solution of problem (1.1) and that it satisfies $u \geq c\rho_{\partial\Omega}$ for some constant $c > 0$.

In Section 4 we prove Theorem 1.6. The existence assertion of 1.6 is obtained by adapting, to our setting, ideas from the proof of Theorem 1.1 in [35] (which is a sub-supersolution theorem for problems of the form $-\Delta u = f(x, u)$ in Ω , $u = 0$ on $\partial\Omega$). Lemma 4.4 gives a weak form of the Hopf boundary lemma, and Corollary 1.7 is proved as a direct consequence of Theorem 1.6 and of Lemma 4.4.

2. PRELIMINARIES

Let us recall some well known facts.

Remark 2.1.

- (i) (Poincaré's inequality for functions in $H_0^1(\Omega)$, see, e.g., [38, Theorem 1.8.1])
There exists a positive constant c such that

$$\|u\|_2 \leq c \|\nabla u\|_2 \quad \text{for any } u \in H_0^1(\Omega).$$

- (ii) (Poincaré's inequality for functions in $H_{0,\Gamma_1}^1(\Omega)$, see, e.g., [43, Theorem 7.16])
There exists a positive constant c such that

$$\|u\|_2 \leq c \|\nabla u\|_2 \quad \text{for any } u \in H_{0,\Gamma_1}^1(\Omega).$$

- (iii) The inclusion $H_{0,\Gamma_1}^1(\Omega) \hookrightarrow L^2(\Omega)$ is compact. Indeed, the inclusion $H_{0,\Gamma_1}^1(\Omega) \hookrightarrow H^1(\Omega)$ is continuous and (see, e.g., [38, Theorem 1.9.15]) $H^1(\Omega)$ has compact inclusion into $L^2(\Omega)$.
- (iv) (Hardy's inequality, see, e.g., [6, p. 313], see also [38, Theorem 1.10.15])

There exists a positive constant c such that $\left\| \frac{u}{\rho_{\partial\Omega}} \right\|_2 \leq c \|\nabla u\|_2$ for any $u \in H_0^1(\Omega)$.

$H_{0,\Gamma_1}^1(\Omega)$ is a closed subspace of $H^1(\Omega)$ and thus, provided with the norm of $H^1(\Omega)$, it is a Hilbert space, and the Poincaré inequality of Remark 2.1(ii) gives that $u \rightarrow \|\nabla u\|_2$ is a norm on $H_{0,\Gamma_1}^1(\Omega)$, equivalent to the norm $\|\cdot\|_{H^1(\Omega)}$. From now on, we will consider $H_{0,\Gamma_1}^1(\Omega)$ as a Hilbert space provided with the norm $\|u\|_{H_{0,\Gamma_1}^1(\Omega)} := \|\nabla u\|_2$. Similarly, $H_0^1(\Omega)$ will be considered as a Hilbert space with the same norm.

For $\delta > 0$, let

$$\Omega_\delta := \{x \in \Omega : \rho_{\partial\Omega}(x) > \delta\} \quad \text{and} \quad A_\delta := \{x \in \Omega : \rho_{\partial\Omega}(x) \leq \delta\}.$$

Similarly, for $i = 1, 2$, we set

$$\Omega_{\Gamma_i, \delta} := \{x \in \Omega : \rho_{\Gamma_i}(x) > \delta\} \quad \text{and} \quad A_{\Gamma_i, \delta} := \{x \in \Omega : \rho_{\Gamma_i}(x) \leq \delta\}. \quad (2.1)$$

The following lemma provides an analogous of the Hardy inequality for functions in $H_{0, \Gamma_1}^1(\Omega)$.

Lemma 2.2 (Hardy's inequality for functions in $H_{0, \Gamma_1}^1(\Omega)$). *There exists a positive constant c such that*

$$\left\| \frac{u}{\rho_{\Gamma_1}} \right\|_2 \leq c \|\nabla u\|_2$$

for any $u \in H_{0, \Gamma_1}^1(\Omega)$.

Proof. Along the proof, c, c', c'' etc., will denote positive constants independent of u . Let δ_1, δ_2 be such that $0 < \delta_1 < \delta_2$ and $\Omega_{\Gamma_1, \delta_2} \neq \emptyset$. Let $\psi \in C^\infty(\overline{\Omega})$ be such that $0 \leq \psi \leq 1$ in Ω , $\psi = 1$ in A_{Γ_1, δ_1} and $\psi = 0$ in $\Omega_{\Gamma_1, \delta_2}$. Then, for $u \in H_{0, \Gamma_1}^1(\Omega)$,

$$\left\| \frac{u}{\rho_{\Gamma_1}} \right\|_2^2 = \int_{\Omega} \frac{u^2}{\rho_{\Gamma_1}^2} = \int_{A_{\Gamma_1, \delta_1}} \frac{u^2}{\rho_{\Gamma_1}^2} + \int_{\Omega \setminus A_{\Gamma_1, \delta_1}} \frac{u^2}{\rho_{\Gamma_1}^2}. \quad (2.2)$$

Now, $u\psi \in H_0^1(\Omega)$ and so, taking into account the Hardy inequality in $H_0^1(\Omega)$,

$$\int_{A_{\Gamma_1, \delta_1}} \frac{u^2}{\rho_{\Gamma_1}^2} = \int_{A_{\Gamma_1, \delta_1}} \frac{u^2 \psi^2}{\rho_{\Gamma_1}^2} \leq \int_{\Omega} \frac{u^2 \psi^2}{\rho_{\partial\Omega}^2} \leq c \int_{\Omega} |\nabla(u\psi)|^2 = c \int_{\Omega} |\psi \nabla u + u \nabla \psi|^2.$$

Thus

$$\begin{aligned} \left\| \frac{u}{\rho_{\Gamma_1}} \right\|_{L^2(A_{\Gamma_1, \delta_1})} &\leq c \|\psi \nabla u + u \nabla \psi\|_{L^2(\Omega)} \\ &\leq c \|\psi\|_{L^\infty(\Omega)} \|\nabla u\|_{L^2(\Omega)} + c \|\nabla \psi\|_{L^\infty(\Omega)} \|u\|_{L^2(\Omega)}. \end{aligned} \quad (2.3)$$

and, by the Poincaré inequality of Remark 2.1(ii), $\|u\|_{L^2(\Omega)} \leq c' \|\nabla u\|_{L^2(\Omega)}$. Thus,

$$\left\| \frac{u}{\rho_{\Gamma_1}} \right\|_{L^2(A_{\Gamma_1, \delta_1})} \leq c'' (\|\psi\|_{L^\infty(\Omega)} + \|\nabla \psi\|_{L^\infty(\Omega)}) \|\nabla u\|_{L^2(\Omega)}.$$

On the other hand,

$$\int_{\Omega \setminus A_{\Gamma_1, \delta_1}} \frac{u^2}{\rho_{\Gamma_1}^2} \leq \frac{1}{\delta_1^2} \int_{\Omega \setminus A_{\Gamma_1, \delta_1}} u^2 \leq \frac{1}{\delta_1^2} \int_{\Omega} u^2 \leq c''' \|\nabla u\|_{L^2(\Omega)}^2, \quad (2.4)$$

the last inequality by the Poincaré inequality of Remark 2.1(ii), and the lemma follows from (2.2), (2.3), and (2.4). \square

Corollary 2.3.

(i) If $f : \Omega \rightarrow \mathbb{R}$ and $f \in L^2(\Omega, \rho_{\partial\Omega}^2(x)dx)$, then $f \in (H_0^1(\Omega))'$ and

$$\|f\|_{(H_0^1(\Omega))'} \leq c \|f\|_{L^2(\Omega, \rho_{\partial\Omega}^2(x)dx)}$$

with c a positive constant independent of f .

(ii) If $f : \Omega \rightarrow \mathbb{R}$ and $f \in L^2(\Omega, \rho_{\Gamma_1}^2(x)dx)$, then $f \in (H_{0,\Gamma_1}^1(\Omega))'$ and it holds that

$$\|f\|_{(H_{0,\Gamma_1}^1(\Omega))'} \leq c \|f\|_{L^2(\Omega, \rho_{\Gamma_1}^2(x)dx)},$$

where c is a positive constant independent of f .

Proof. Suppose that $\rho_{\partial\Omega}f \in L^2(\Omega)$ and let $\varphi \in H_0^1(\Omega)$. Then, for some positive constant c independent of φ ,

$$\int_{\Omega} |f\varphi| = \int_{\Omega} \left| \rho_{\partial\Omega}f \frac{\varphi}{\rho_{\partial\Omega}} \right| \leq \|\rho_{\partial\Omega}f\|_2 \left\| \frac{\varphi}{\rho_{\partial\Omega}} \right\|_2 \leq c \|\rho_{\partial\Omega}f\|_2 \|\varphi\|_{H_0^1(\Omega)},$$

the last inequality by Remark 2.1(iii). Thus (i) holds. The proof of (ii) is similar, using Lemma 2.2 instead of Remark 2.1(iii). \square

Remark 2.4 (see, e.g., [43, Theorem 8.9]). If $0 \leq f \in (H_0^1(\Omega))'$, $0 \leq \sigma \in H^{\frac{1}{2}}(\partial\Omega)$, and if u is the weak solution of problem (1.7), then $u \geq 0$ in Ω .

Remark 2.5.

(i) (see [7, Lemma 3.2]) Suppose $0 \leq f \in L^\infty(\Omega)$, and let ζ be the solution of the problem

$$\begin{cases} -\Delta\zeta = f & \text{in } \Omega, \\ \zeta = 0 & \text{on } \partial\Omega. \end{cases} \quad (2.5)$$

Then $\zeta \geq c\rho_{\partial\Omega} \int_{\Omega} f\rho_{\partial\Omega}$ in Ω , with c a positive constant independent of f .

(ii) If $0 \leq f \in L^\infty(\Omega)$, $0 \leq \sigma \in H^{\frac{1}{2}}(\partial\Omega)$ and if $u \in H^1(\Omega)$ is a weak solution of the problem

$$\begin{cases} -\Delta u = f & \text{in } \Omega, \\ u = \sigma & \text{on } \partial\Omega, \end{cases} \quad (2.6)$$

then $u \geq c\rho_{\partial\Omega} \int_{\Omega} f\rho_{\partial\Omega}$ in Ω with c a positive constant independent of f . Indeed, let ζ be as in (i), then $u - \zeta$ satisfies, in a weak sense,

$$\begin{cases} -\Delta(u - \zeta) = f & \text{in } \Omega, \\ u - \zeta = \sigma & \text{on } \partial\Omega, \end{cases}$$

and then, by Remark 2.4, $u \geq \zeta$. Thus, by (i), $u \geq c\rho_{\partial\Omega} \int_{\Omega} f\rho_{\partial\Omega}$ in Ω , with c as in (i).

(iii) Let $f : \Omega \rightarrow \mathbb{R}$ be a nonnegative and measurable function such that $f \in (H_0^1(\Omega))'$ and $|\{x \in \Omega : f(x) > 0\}| > 0$. If $0 \leq \sigma \in H^{\frac{1}{2}}(\partial\Omega)$ and if $u \in H^1(\Omega)$ is a weak solution of the problem (2.6), then there exists a positive constant c such

that $u \geq c\rho_{\partial\Omega}$ in Ω . In fact, in such a case there exist a measurable subset $F \subset \Omega$ with $|F| > 0$ and $\lambda \in (0, \infty)$ such that $f \geq \lambda\chi_F$ in Ω . Let w be the solution of $-\Delta w = \lambda\chi_F$ in Ω , $w = 0$ on $\partial\Omega$. Then, by (i), there exists a positive constant c such that $w \geq c\rho_{\partial\Omega}$ in Ω . Also, $-\Delta(u - w) = f - \lambda\chi_F \geq 0$ in Ω and $u - w = \sigma \geq 0$ on $\partial\Omega$. Thus, by Remark 2.4, $u - w \geq 0$, and then $u \geq c\rho_{\partial\Omega}$ in Ω .

Remark 2.6. Suppose $0 \leq f \in (H_{0,\Gamma_1}^1(\Omega))'$, $0 \leq \tau \in H^{\frac{1}{2}}(\Gamma_1)$ and let $\eta : \Gamma_2 \rightarrow \mathbb{R}$ be such that $0 \leq \eta \in (H_{0,\Gamma_1}^1(\Omega))'$. If u is the weak solution of problem (1.9), then $u \geq 0$ in Ω . Indeed, since $\tau \geq 0$ we have $u^- = 0$ on Γ_1 and thus $u^- \in H_{0,\Gamma_1}^1(\Omega)$. Taking $\varphi = -u^-$ in (1.10) we get

$$-\int_{\Omega} \langle \nabla u, \nabla u^- \rangle + \int_{\Omega} f u^- + \int_{\Gamma_2} \eta u^- = 0,$$

and so

$$\int_{\Omega} |\nabla u^-|^2 = -\int_{\Omega} f u^- - \int_{\Gamma_2} \eta u^- \leq 0.$$

Thus $\int_{\Omega} |\nabla u^-|^2 = 0$. Therefore, by the Poincaré inequality of Remark 2.1(ii), $u^- = 0$ in Ω . Then $u \geq 0$ in Ω . Moreover, from Remark 2.5 (iii) used with $\sigma := u|_{\partial\Omega} \geq 0$ on $\partial\Omega$ (the restriction in the sense of the trace), it follows that, if in addition, $|\{x \in \Omega : f(x) > 0\}| > 0$, then there exists a positive constant c such that $u \geq c\rho_{\partial\Omega}$ in Ω .

3. THE CASE OF DIRICHLET BOUNDARY CONDITION

We assume, for the whole section, that $g : \Omega \times (0, \infty) \rightarrow \mathbb{R}$ satisfies the conditions (H1)–(H3) of Theorem 1.3. We first study, for $\varepsilon \in (0, 1]$ and for a nonnegative $\sigma \in H^{\frac{1}{2}}(\partial\Omega)$, the approximated problem

$$\begin{cases} -\Delta u = g_{\varepsilon}(\cdot, u) & \text{in } \Omega, \\ u = \sigma & \text{on } \partial\Omega, \end{cases} \quad (3.1)$$

where $g_{\varepsilon} : \Omega \times (0, \infty) \rightarrow \mathbb{R}$ is defined by

$$g_{\varepsilon}(x, s) := \min \{ \varepsilon^{-1}, g(x, s + \varepsilon) \}. \quad (3.2)$$

Observe that, since g satisfies (H1)–(H3), the same conditions hold for each g_{ε} . Let $\tilde{\sigma} \in H^1(\Omega)$ be the weak solution of the problem

$$\begin{cases} -\Delta \tilde{\sigma} = 0 & \text{in } \Omega, \\ \tilde{\sigma} = \sigma & \text{on } \partial\Omega. \end{cases} \quad (3.3)$$

Then, by Remark 2.4(i), $\tilde{\sigma} \geq 0$ in Ω . By writing $u = \tilde{\sigma} + v$, problem (3.1) becomes equivalent to the problem of finding a weak solution $v \in H_0^1(\Omega)$ of the problem

$$\begin{cases} -\Delta v = g_\varepsilon(\cdot, v + \tilde{\sigma}) & \text{in } \Omega, \\ v = 0 & \text{on } \partial\Omega. \end{cases} \quad (3.4)$$

Let $(-\Delta)^{-1} : L^2(\Omega) \rightarrow H_0^1(\Omega)$ be the solution operator of the homogeneous Dirichlet problem defined by $(-\Delta)^{-1}h = u$, where $u \in H_0^1(\Omega)$ is the weak solution of the problem $-\Delta u = h$ in Ω , $u = 0$ on $\partial\Omega$. We recall that $(-\Delta)^{-1} : L^2(\Omega) \rightarrow H_0^1(\Omega)$ is continuous and that, since $H_0^1(\Omega)$ has compact inclusion into $L^2(\Omega)$, $(-\Delta)^{-1} : L^2(\Omega) \rightarrow L^2(\Omega)$ is a compact operator. Let $T_\varepsilon : L^2(\Omega) \rightarrow H_0^1(\Omega)$ be defined by

$$T_\varepsilon(v) := (-\Delta)^{-1}(g_\varepsilon(\cdot, v + \tilde{\sigma})),$$

and let $C_\varepsilon := \{v \in L^2(\Omega) : 0 \leq v \leq \frac{1}{\varepsilon}(-\Delta)^{-1}(\mathbf{1})\}$. We have the following:

Lemma 3.1.

- (i) C_ε is a bounded, closed and convex subset of $L^2(\Omega)$.
- (ii) $T_\varepsilon(C_\varepsilon) \subset C_\varepsilon$.
- (iii) $T_\varepsilon : C_\varepsilon \rightarrow C_\varepsilon$ is continuous.
- (iv) $T_\varepsilon : C_\varepsilon \rightarrow C_\varepsilon$ is a compact operator.

Proof. (i) is obvious.

To show (ii) observe that if $v \in C_\varepsilon$ then $0 \leq g_\varepsilon(\cdot, v + \tilde{\sigma}) \leq \frac{1}{\varepsilon}$ a.e. in Ω and so, by Remark 2.4,

$$0 \leq (-\Delta)^{-1}(g_\varepsilon(x, v + \tilde{\sigma})) \leq \frac{1}{\varepsilon}(-\Delta)^{-1}(\mathbf{1}).$$

Thus $T_\varepsilon(v) \in C_\varepsilon$.

To prove (iii) it is enough to see that if $v \in C_\varepsilon$, and if $\{v_j\}_{j \in \mathbb{N}}$ is a sequence in C_ε that converges to v in $L^2(\Omega)$, then there exists a subsequence $\{v_{j_k}\}_{k \in \mathbb{N}}$ such that $\{T_\varepsilon(v_{j_k})\}_{k \in \mathbb{N}}$ converges to $T_\varepsilon(v)$ in $L^2(\Omega)$. Let $v \in C_\varepsilon$, and let $\{v_j\}_{j \in \mathbb{N}}$ be a sequence in C_ε which converges to v in $L^2(\Omega)$, then there exists a subsequence $\{v_{j_k}\}_{k \in \mathbb{N}}$ such that $\{v_{j_k}\}_{k \in \mathbb{N}}$ converges to v a.e. in Ω . Thus, since g_ε is a Carathéodory function, $\{g_\varepsilon(\cdot, v_{j_k} + \tilde{\sigma})\}_{k \in \mathbb{N}}$ converges to $g_\varepsilon(\cdot, v + \tilde{\sigma})$ a.e. in Ω . Then

$$\lim_{k \rightarrow \infty} |g_\varepsilon(\cdot, v_{j_k} + \tilde{\sigma}) - g_\varepsilon(\cdot, v + \tilde{\sigma})|^2 = 0$$

a.e. in Ω . Since $|g_\varepsilon(\cdot, v_{j_k} + \tilde{\sigma}) - g_\varepsilon(\cdot, v + \tilde{\sigma})|^2 \leq \frac{1}{\varepsilon^2}$, the Lebesgue dominated convergence theorem gives that $\{g_\varepsilon(\cdot, v_{j_k} + \tilde{\sigma})\}_{k \in \mathbb{N}}$ converges to $g_\varepsilon(\cdot, v + \tilde{\sigma})$ in $L^2(\Omega)$. Then $\{(-\Delta)^{-1}(g_\varepsilon(\cdot, v_{j_k} + \tilde{\sigma}))\}_{k \in \mathbb{N}}$ converges to $(-\Delta)^{-1}(g_\varepsilon(\cdot, v + \tilde{\sigma}))$ in $L^2(\Omega)$, i.e., $\{T_\varepsilon(v_{j_k})\}_{k \in \mathbb{N}}$ converges to $T_\varepsilon(v)$ in $L^2(\Omega)$. Thus (iii) holds.

To see (iv), note that $\{g_\varepsilon(\cdot, v_j + \tilde{\sigma})\}_{j \in \mathbb{N}}$ is bounded in $L^2(\Omega)$ for any sequence $\{v_j\}_{j \in \mathbb{N}}$ in C_ε , and so (iv) follows immediately from the compactness of the solution operator $(-\Delta)^{-1} : L^2(\Omega) \rightarrow L^2(\Omega)$. \square

Lemma 3.2.

(i) For $\varepsilon \in (0, 1]$, the problem

$$\begin{cases} -\Delta v_\varepsilon = g_\varepsilon(\cdot, v_\varepsilon + \tilde{\sigma}) & \text{in } \Omega, \\ v_\varepsilon = 0 & \text{on } \partial\Omega \end{cases} \quad (3.5)$$

has a unique weak solution $v_\varepsilon \in H_0^1(\Omega)$.

(ii) The map $\varepsilon \rightarrow v_\varepsilon$ is nonincreasing.

(iii) There exists a positive constant c such that $v_\varepsilon \geq c\rho_{\partial\Omega}$ for any $\varepsilon \in (0, 1]$.

(iv) $\{v_\varepsilon\}_{\varepsilon \in (0, 1]}$ is bounded in $H_0^1(\Omega)$.

Proof. From Lemma 3.1 and the Schauder fixed point theorem, T_ε has a fixed point $v_\varepsilon \in C_\varepsilon$, and so v_ε is a weak solution of problem (3.4). Suppose that $w \in H^1(\Omega)$ is another solution of (3.4). Then $v_\varepsilon - w \in H_0^1(\Omega)$ and it satisfies, in weak sense

$$\begin{cases} -\Delta(v_\varepsilon - w) = g_\varepsilon(\cdot, v_\varepsilon + \tilde{\sigma}) - g_\varepsilon(\cdot, w + \tilde{\sigma}) & \text{in } \Omega, \\ v_\varepsilon - w = 0 & \text{on } \partial\Omega. \end{cases} \quad (3.6)$$

Now, $g_\varepsilon(x, \cdot)$ is nonincreasing on $(0, \infty)$ for a.e. $x \in \Omega$, and so

$$g_\varepsilon(\cdot, v_\varepsilon + \tilde{\sigma}) - g_\varepsilon(\cdot, w + \tilde{\sigma})(v_\varepsilon - w) \leq 0 \quad \text{a.e in } \Omega.$$

Thus, taking $v_\varepsilon - w$ as a test function in (3.6), we get that $\|\nabla(v_\varepsilon - w)\|_2 = 0$, and so, by the Poincaré inequality, $v_\varepsilon = w$ in Ω . Thus (i) holds.

To prove (ii), suppose that $0 < \varepsilon < \theta \leq 1$. Then $g_\varepsilon \geq g_\theta$ on $\Omega \times (0, \infty)$. Thus, in a weak sense,

$$\begin{cases} -\Delta(v_\varepsilon) = g_\varepsilon(\cdot, v_\varepsilon + \tilde{\sigma}) \geq g_\theta(\cdot, v_\varepsilon + \tilde{\sigma}) & \text{in } \Omega, \\ v_\varepsilon = 0 & \text{on } \partial\Omega. \end{cases} \quad (3.7)$$

Also,

$$\begin{cases} -\Delta(v_\theta) = g_\theta(\cdot, v_\theta + \tilde{\sigma}) & \text{in } \Omega, \\ v_\theta = 0 & \text{on } \partial\Omega. \end{cases} \quad (3.8)$$

and so, again in a weak sense,

$$\begin{cases} -\Delta(v_\varepsilon - v_\theta) = g_\varepsilon(\cdot, v_\varepsilon + \tilde{\sigma}) - g_\theta(\cdot, v_\theta + \tilde{\sigma}) \\ \geq g_\theta(\cdot, v_\varepsilon + \tilde{\sigma}) - g_\theta(\cdot, v_\theta + \tilde{\sigma}) & \text{in } \Omega, \\ v_\varepsilon - v_\theta = 0 & \text{on } \partial\Omega. \end{cases} \quad (3.9)$$

and so, taking $-(v_\varepsilon - v_\theta)^-$ as a test function in (3.9) we get

$$\int_{\Omega} |\nabla((v_\varepsilon - v_\theta)^-)|^2 \leq - \int_{\{v_\varepsilon - v_\theta < 0\}} (g_\theta(\cdot, v_\varepsilon + \tilde{\sigma}) - g_\theta(\cdot, v_\theta + \tilde{\sigma}))(v_\varepsilon - v_\theta)^- \leq 0.$$

The last inequality because $g_\varepsilon(x, \cdot)$ is nonincreasing on $(0, \infty)$ for a.e. $x \in \Omega$. Thus $\int_\Omega |\nabla((v_\varepsilon - v_\theta)^-)|^2 = 0$, and so, by Remark 2.1(i), $(v_\varepsilon - v_\theta)^- = 0$ in Ω , and then $v_\varepsilon \geq v_\theta$ a.e. in Ω . Thus (ii) holds.

To see (iii), observe that for $\varepsilon \in (0, 1]$, by (ii), $v_\varepsilon \geq v_1$. Since $-\Delta v_1 = g_1(\cdot, v_1)$ in Ω and $0 \leq g_1(\cdot, v_1) \in L^\infty(\Omega)$, and taking into account that, by (H2), $g_1(\cdot, v_1)$ is not identically zero, Remark 2.5(i) gives that $v_1 \geq c\rho_{\partial\Omega}$ for some positive constant c . Thus $v_\varepsilon \geq c\rho_{\partial\Omega}$ and (iii) holds.

It remains to show (iv). Let c be as in (iii). We take v_ε as a test function in (3.4) to obtain

$$\begin{aligned} \|\nabla v_\varepsilon\|_2^2 &= \int_\Omega |\nabla v_\varepsilon|^2 = \int_\Omega v_\varepsilon g_\varepsilon(\cdot, v_\varepsilon + \tilde{\sigma}) \\ &\leq \int_\Omega v_\varepsilon g(\cdot, v_\varepsilon + \tilde{\sigma}) \leq \int_\Omega v_\varepsilon g(\cdot, \tilde{\sigma} + c\rho_{\partial\Omega}) \\ &= \int_\Omega \frac{v_\varepsilon}{\rho_{\partial\Omega}} \rho_{\partial\Omega} g(\cdot, \tilde{\sigma} + c\rho_{\partial\Omega}) \leq \int_\Omega \frac{v_\varepsilon}{\rho_{\partial\Omega}} \rho_{\partial\Omega} g(\cdot, c\rho_{\partial\Omega}) \end{aligned} \quad (3.10)$$

where we have used (iii), (H1), and that $g_\varepsilon \leq g$, as well as that $g(x, s)$ is nonincreasing in s . Now, by the Hölder inequality and Remark 2.1(iv), we have, for some positive constant c' independent of ε ,

$$\int_\Omega \frac{v_\varepsilon}{\rho_{\partial\Omega}} \rho_{\partial\Omega} g(\cdot, c\rho_{\partial\Omega}) \leq \left\| \frac{v_\varepsilon}{\rho_{\partial\Omega}} \right\|_2 \|\rho_{\partial\Omega} g(\cdot, c\rho_{\partial\Omega})\|_2 \leq c' \|\nabla v_\varepsilon\|_2 \|\rho_{\partial\Omega} g(\cdot, c\rho_{\partial\Omega})\|_2 \quad (3.11)$$

and, by (H3), $\|\rho_{\partial\Omega} g(\cdot, c\rho_{\partial\Omega})\|_2 < \infty$. Thus, from (3.10) and (3.11), we get

$$\|\nabla v_\varepsilon\|_2 \leq c' \|\rho_{\partial\Omega} g(\cdot, c\rho_{\partial\Omega})\|_2,$$

which ends the proof of the lemma. \square

Lemma 3.3. For $\varepsilon \in (0, 1]$, let $v_\varepsilon \in H_0^1(\Omega)$ be as given by Lemma 3.2, and let $v := \lim_{\varepsilon \rightarrow 0^+} v_\varepsilon$. Then:

- (i) $v \in H_0^1(\Omega)$ and $\lim_{\varepsilon \rightarrow 0^+} v_\varepsilon = v$ with convergence in $H_0^1(\Omega)$,
- (ii) $\lim_{\varepsilon \rightarrow 0^+} g_\varepsilon(\cdot, v_\varepsilon + \tilde{\sigma}) = g(\cdot, v + \tilde{\sigma})$ with convergence in $L^2(\Omega, \rho_{\partial\Omega}^2(x) dx)$.

Proof. Observe that $v \in H_0^1(\Omega)$. Indeed, let $\{\theta_j\}_{j \in \mathbb{N}} \subset (0, 1]$ be a sequence such that $\lim_{j \rightarrow \infty} \theta_j = 0$. By Lemma 3.2, $\{v_{\theta_j}\}_{j \in \mathbb{N}}$ is bounded in $H_0^1(\Omega)$. Thus there exist a subsequence $\{v_{\theta_{j_k}}\}_{k \in \mathbb{N}}$ and a function $w \in H_0^1(\Omega)$ such that $\{v_{\theta_{j_k}}\}_{k \in \mathbb{N}}$ converges to w strongly in $L^2(\Omega)$, and $\{\nabla v_{\theta_{j_k}}\}_{k \in \mathbb{N}}$ converges to ∇w weakly in $L^2(\Omega, \mathbb{R}^n)$. After pass to a further subsequence if necessary, we can assume also that $\{v_{\theta_{j_k}}\}_{k \in \mathbb{N}}$ converges to w a.e. in Ω . Since $v := \lim_{\varepsilon \rightarrow 0^+} v_\varepsilon$ it follows that $w = v$ and then $v \in H_0^1(\Omega)$.

To prove the lemma it is enough to see that for any sequence $\{\varepsilon_j\}_{j \in \mathbb{N}} \subset (0, 1]$ such that $\lim_{j \rightarrow \infty} \varepsilon_j = 0$ there exists a subsequence, which we still denoted by $\{\varepsilon_j\}_{j \in \mathbb{N}}$,

such that

$$\lim_{j \rightarrow \infty} \|v_{\varepsilon_j} - v\|_{H_0^1(\Omega)}^2 = 0$$

and

$$\lim_{j \rightarrow \infty} \|g_{\varepsilon_j}(\cdot, v_{\varepsilon_j} + \tilde{\sigma}) - g(\cdot, v + \tilde{\sigma})\|_{L^2(\Omega, \rho_{\partial\Omega}^2(x) dx)} = 0.$$

Now, in a weak sense,

$$\begin{cases} -\Delta(v_{\varepsilon_j} - v) = g_{\varepsilon_j}(\cdot, v_{\varepsilon_j} + \tilde{\sigma}) - g(\cdot, v + \tilde{\sigma}) & \text{in } \Omega, \\ v_{\varepsilon_j} - v = 0 & \text{on } \partial\Omega. \end{cases} \quad (3.12)$$

We take $v_{\varepsilon_j} - v$ as a test function in (3.12) and we use the Hardy inequality of Remark 2.1(iv) to obtain

$$\begin{aligned} \|v_{\varepsilon_j} - v\|_{H_0^1(\Omega)}^2 &= \int_{\Omega} |\nabla(v_{\varepsilon_j} - v)|^2 = \int_{\Omega} (g_{\varepsilon_j}(\cdot, v_{\varepsilon_j} + \tilde{\sigma}) - g(\cdot, v + \tilde{\sigma}))(v_{\varepsilon_j} - v) \\ &= \int_{\Omega} \rho_{\partial\Omega} (g_{\varepsilon_j}(\cdot, v_{\varepsilon_j} + \tilde{\sigma}) - g(\cdot, v + \tilde{\sigma})) \frac{v_{\varepsilon_j} - v}{\rho_{\partial\Omega}} \\ &\leq c \|\rho_{\partial\Omega} (g_{\varepsilon_j}(\cdot, v_{\varepsilon_j} + \tilde{\sigma}) - g(\cdot, v + \tilde{\sigma}))\|_2 \|v_{\varepsilon_j} - v\|_{H_0^1(\Omega)}. \end{aligned}$$

where c is a positive constant independent of j . Then, in order to prove the lemma, it suffices to show that

$$\lim_{j \rightarrow \infty} \|\rho_{\partial\Omega} (g_{\varepsilon_j}(\cdot, v_{\varepsilon_j} + \tilde{\sigma}) - g(\cdot, v + \tilde{\sigma}))\|_2 = 0. \quad (3.13)$$

Now,

$$\begin{aligned} &\|\rho_{\partial\Omega} (g_{\varepsilon_j}(\cdot, v_{\varepsilon_j} + \tilde{\sigma}) - g(\cdot, v + \tilde{\sigma}))\|_2^2 \\ &= \int_{\left\{g(\cdot, v_{\varepsilon_j} + \tilde{\sigma} + \varepsilon_j) \leq \frac{1}{\varepsilon_j}\right\}} \rho_{\partial\Omega}^2 (g_{\varepsilon_j}(\cdot, v_{\varepsilon_j} + \tilde{\sigma}) - g(\cdot, v + \tilde{\sigma}))^2 \\ &\quad + \int_{\left\{g(\cdot, v_{\varepsilon_j} + \tilde{\sigma} + \varepsilon_j) > \frac{1}{\varepsilon_j}\right\}} \rho_{\partial\Omega}^2 (g_{\varepsilon_j}(\cdot, v_{\varepsilon_j} + \tilde{\sigma}) - g(\cdot, v + \tilde{\sigma}))^2 \\ &= \int_{\left\{g(\cdot, v_{\varepsilon_j} + \tilde{\sigma} + \varepsilon_j) \leq \frac{1}{\varepsilon_j}\right\}} \rho_{\partial\Omega}^2 (g(\cdot, v_{\varepsilon_j} + \tilde{\sigma} + \varepsilon_j) - g(\cdot, v + \tilde{\sigma}))^2 \\ &\quad + \int_{\left\{g(\cdot, v_{\varepsilon_j} + \tilde{\sigma} + \varepsilon_j) > \frac{1}{\varepsilon_j}\right\}} \rho_{\partial\Omega}^2 \left(\frac{1}{\varepsilon} - g(\cdot, v + \tilde{\sigma})\right)^2 \end{aligned}$$

and so

$$\begin{aligned}
& \|\rho_{\partial\Omega}(g_{\varepsilon_j}(\cdot, v_{\varepsilon_j} + \tilde{\sigma}) - g(\cdot, v + \tilde{\sigma}))\|_2^2 \\
&= \int_{\Omega} \rho_{\partial\Omega}^2(g(\cdot, v_{\varepsilon_j} + \tilde{\sigma} + \varepsilon_j) - g(\cdot, v + \tilde{\sigma}))^2 \\
&\quad - \int_{\left\{g(\cdot, v_{\varepsilon_j} + \tilde{\sigma} + \varepsilon_j) > \frac{1}{\varepsilon_j}\right\}} \rho_{\partial\Omega}^2(g(\cdot, v_{\varepsilon_j} + \tilde{\sigma} + \varepsilon_j) - g(\cdot, v + \tilde{\sigma}))^2 \\
&\quad + \int_{\left\{g(\cdot, v_{\varepsilon_j} + \tilde{\sigma} + \varepsilon_j) > \frac{1}{\varepsilon_j}\right\}} \rho_{\partial\Omega}^2\left(\frac{1}{\varepsilon} - g(\cdot, v + \tilde{\sigma})\right)^2 \\
&= I_{1,j} + I_{2,j} + I_{3,j},
\end{aligned}$$

where

$$\begin{aligned}
I_{1,j} &:= \int_{\Omega} \rho_{\partial\Omega}^2(g(\cdot, v_{\varepsilon_j} + \tilde{\sigma} + \varepsilon_j) - g(\cdot, v + \tilde{\sigma}))^2, \\
I_{2,j} &:= - \int_{\left\{g(\cdot, v_{\varepsilon_j} + \tilde{\sigma} + \varepsilon_j) > \frac{1}{\varepsilon_j}\right\}} \rho_{\partial\Omega}^2(g(\cdot, v_{\varepsilon_j} + \tilde{\sigma} + \varepsilon_j) - g(\cdot, v + \tilde{\sigma}))^2, \\
I_{3,j} &:= \int_{\left\{g(\cdot, v_{\varepsilon_j} + \tilde{\sigma} + \varepsilon_j) > \frac{1}{\varepsilon_j}\right\}} \rho_{\partial\Omega}^2\left(\frac{1}{\varepsilon} - g(\cdot, v + \tilde{\sigma})\right)^2.
\end{aligned}$$

Now, since g is Carathéodory,

$$\lim_{j \rightarrow \infty} \rho_{\partial\Omega}^2(g(\cdot, v_{\varepsilon_j} + \tilde{\sigma} + \varepsilon_j) - g(\cdot, v + \tilde{\sigma}))^2 = 0$$

a.e. in Ω . Also,

$$\begin{aligned}
& \rho_{\partial\Omega}^2(g(\cdot, v_{\varepsilon_j} + \tilde{\sigma} + \varepsilon_j) - g(\cdot, v + \tilde{\sigma}))^2 \\
& \leq 2\rho_{\partial\Omega}^2 g^2(\cdot, v_{\varepsilon_j} + \tilde{\sigma} + \varepsilon_j) + 2\rho_{\partial\Omega}^2 g^2(\cdot, v + \tilde{\sigma}) \leq 4\rho_{\partial\Omega}^2 g^2(\cdot, v + \tilde{\sigma}),
\end{aligned}$$

and since $v \geq c\rho_{\partial\Omega}$ and $\tilde{\sigma} \geq 0$, (H3) gives $\rho_{\partial\Omega}^2 g^2(\cdot, v + \tilde{\sigma}) \in L^1(\Omega)$. Then, by the Lebesgue dominated convergence theorem,

$$\lim_{j \rightarrow \infty} I_{1,j} = 0.$$

Let

$$U_j := \left\{g(\cdot, v_1 + \tilde{\sigma}) > \frac{1}{\varepsilon_j}\right\}.$$

Then $U_{j+1} \subset U_j$ for any j , $U_1 \subset \Omega$, and $\bigcap_{j=1}^{\infty} U_j = \{g(\cdot, v_1 + \tilde{\sigma}) = \infty\}$. Since $\rho_{\partial\Omega}^2 g^2(\cdot, v + \tilde{\sigma}) \in L^1(\Omega)$ it follows that $\left| \bigcap_{j=1}^{\infty} U_j \right| = 0$. Then $\lim_{j \rightarrow \infty} |U_j| = 0$, and thus

$$\lim_{j \rightarrow \infty} \int_{U_j} \rho_{\partial\Omega}^2 g^2(\cdot, v + \tilde{\sigma}) = 0. \quad (3.14)$$

Now,

$$\rho_{\partial\Omega}^2 (g(\cdot, v_{\varepsilon_j} + \tilde{\sigma} + \varepsilon_j) - g(\cdot, v + \tilde{\sigma}))^2 \leq 2\rho_{\partial\Omega}^2 g^2(\cdot, v + \tilde{\sigma})$$

and so

$$\begin{aligned} |I_{2,j}| &\leq \int_{\left\{g(\cdot, v_{\varepsilon_j} + \tilde{\sigma} + \varepsilon_j) > \frac{1}{\varepsilon_j}\right\}} \rho_{\partial\Omega}^2 (g(\cdot, v_{\varepsilon_j} + \tilde{\sigma} + \varepsilon_j) - g(\cdot, v + \tilde{\sigma}))^2 \\ &\leq \int_{U_j} 2\rho_{\partial\Omega}^2 g^2(\cdot, v + \tilde{\sigma}). \end{aligned}$$

Then, by (3.14),

$$\lim_{j \rightarrow \infty} I_{2,j} = 0.$$

Finally,

$$\rho_{\partial\Omega}^2 \left(\frac{1}{\varepsilon} - g(\cdot, v + \tilde{\sigma}) \right)^2 \leq 2\rho_{\partial\Omega}^2 \frac{1}{\varepsilon^2} + 2\rho_{\partial\Omega}^2 g^2(\cdot, v + \tilde{\sigma}),$$

and then

$$\begin{aligned} |I_{3,j}| &\leq \int_{\left\{g(\cdot, v_{\varepsilon_j} + \tilde{\sigma} + \varepsilon_j) > \frac{1}{\varepsilon_j}\right\}} \left(2\rho_{\partial\Omega}^2 \frac{1}{\varepsilon^2} + 2\rho_{\partial\Omega}^2 g^2(\cdot, v + \tilde{\sigma}) \right) \\ &\leq 2 \int_{\left\{g(\cdot, v_{\varepsilon_j} + \tilde{\sigma} + \varepsilon_j) > \frac{1}{\varepsilon_j}\right\}} \rho_{\partial\Omega}^2 g^2(\cdot, v_{\varepsilon_j} + \tilde{\sigma} + \varepsilon_j) \\ &\quad + 2 \int_{\left\{g(\cdot, v_{\varepsilon_j} + \tilde{\sigma} + \varepsilon_j) > \frac{1}{\varepsilon_j}\right\}} \rho_{\partial\Omega}^2 g^2(\cdot, v + \tilde{\sigma}) \\ &\leq 4 \int_{U_j} \rho_{\partial\Omega}^2 g^2(\cdot, v + \tilde{\sigma}), \end{aligned}$$

and thus, by (3.14),

$$\lim_{j \rightarrow \infty} I_{3,j} = 0,$$

which concludes the proof of the lemma. \square

Proof of Theorem 1.3. For $\varepsilon \in (0, 1]$, let $v_\varepsilon \in H_0^1(\Omega)$ be as given by Lemma 3.2 and let $v = \lim_{\varepsilon \rightarrow 0^+} v_\varepsilon$. Let $u_\varepsilon := \tilde{\sigma} + v_\varepsilon$ and let $u := \lim_{\varepsilon \rightarrow 0^+} u_\varepsilon = \tilde{\sigma} + v$. By Lemma 3.2, v_ε is a weak solution of the problem

$$\begin{cases} -\Delta v_\varepsilon = g_\varepsilon(\cdot, v_\varepsilon + \tilde{\sigma}) & \text{in } \Omega, \\ v_\varepsilon = 0 & \text{on } \partial\Omega, \end{cases}$$

and thus $\int_\Omega \langle \nabla v_\varepsilon, \nabla \varphi \rangle = \int_\Omega g_\varepsilon(\cdot, v_\varepsilon + \tilde{\sigma}) \varphi$ for any $\varphi \in H_0^1(\Omega)$, and so (since $\int_\Omega \langle \nabla \tilde{\sigma}, \nabla \varphi \rangle = 0$ for any $\varepsilon \in (0, 1]$ and $\varphi \in H_0^1(\Omega)$)

$$\int_\Omega \langle \nabla u_\varepsilon, \nabla \varphi \rangle = \int_\Omega g_\varepsilon(\cdot, u_\varepsilon) \varphi \quad \text{for any } \varphi \in H_0^1(\Omega). \quad (3.15)$$

Let $\varphi \in H_0^1(\Omega)$. From Lemma 3.3 it follows that $u \in H^1(\Omega)$ and that $\lim_{\varepsilon \rightarrow 0^+} u_\varepsilon = u$ with convergence in $H_0^1(\Omega)$. Then $\lim_{\varepsilon \rightarrow 0^+} \int_\Omega \langle \nabla u_\varepsilon, \nabla \varphi \rangle = \int_\Omega \langle \nabla u, \nabla \varphi \rangle$. Again by Lemma 3.3, $\lim_{\varepsilon \rightarrow 0^+} g_\varepsilon(\cdot, u_\varepsilon) = g(\cdot, u)$ with convergence in $L^2(\Omega, \rho_{\partial\Omega}^2(x) dx)$ and thus $\lim_{\varepsilon \rightarrow 0^+} \int_\Omega g_\varepsilon(\cdot, u_\varepsilon) \varphi = \int_\Omega g(\cdot, u) \varphi$. Then, from (3.15),

$$\int_\Omega \langle \nabla u, \nabla \varphi \rangle = \int_\Omega g(\cdot, u) \varphi.$$

Thus u is a weak solution of problem (1.1). Also, Lemma 3.2 gives that $v_\varepsilon \geq c\rho_{\partial\Omega}$ for some positive constant c independent of ε , and then $u \geq c\rho_{\partial\Omega}$ in Ω .

If w is another weak solution of (1.1), then $u - w \in H_0^1(\Omega)$ and

$$\int_\Omega \langle \nabla(u - w), \nabla \varphi \rangle = \int_\Omega (g(\cdot, u) - g(\cdot, w)) \varphi$$

for any $\varphi \in H_0^1(\Omega)$. We take $\varphi = u - w$ and, since $g(x, s)$ is nonincreasing in s , we get

$$\int_{N\Omega} |\nabla(u - w)|^2 = \int_\Omega (g(\cdot, u) - g(\cdot, w))(u - w) \leq 0.$$

Thus $\int_\Omega |\nabla(u - w)|^2 = 0$ which, by the Poincaré inequality, gives $u = w$. \square

4. THE CASE OF MIXED DIRICHLET-NEUMAN BOUNDARY CONDITIONS

Our aim in this section is to prove Theorems 1.6 and 1.7. We assume, from now on, that $g : \Omega \times (0, \infty) \rightarrow \mathbb{R}$ satisfies the conditions (H1) and (H2) of Theorem 1.3 as well as the condition (H3') of Theorem 1.6. Since the condition (H3') implies the condition (H3) of Theorem 1.3 (because $\rho_{\partial\Omega} \leq \rho_{\Gamma_1}$), all the results of the previous section for the Dirichlet problems still hold under our new assumptions.

Remark 4.1.

(i) If $f \in L^2(\Omega, \rho_{\Gamma_1}^2(x)dx)$, $\tau \in H^{\frac{1}{2}}(\Gamma_1)$ and $\eta \in (H_{0,\Gamma_1}^1(\Omega))'$ (notice that we are not assuming that η is a function defined in Γ_2), then the problem of finding $u \in H^1(\Omega)$ such that

$$\begin{cases} \int_{\Omega} \langle \nabla u, \nabla \varphi \rangle = \int_{\Omega} f \varphi + \eta(\varphi) \text{ for any } \varphi \in H_{0,\Gamma_1}^1(\Omega), \\ u = \tau \text{ on } \Gamma_1 \end{cases} \quad (4.1)$$

has a unique solution, and it satisfies

$$\|u\|_{H^1(\Omega)} \leq c(\|f\|_{(H_0^1(\Omega))'} + \|\tau\|_{H^{\frac{1}{2}}(\Gamma_1)} + \|\eta\|_{(H_{0,\Gamma_1}^1(\Omega))'}). \quad (4.2)$$

for some positive constant c independent of f , τ and η . Indeed, let $\sigma \in H^{\frac{1}{2}}(\partial\Omega)$ be defined by $\sigma = \tau$ on Γ_1 and $\sigma = 0$ on Γ_2 , and let $\xi \in H^1(\Omega)$ be such that $\xi = \sigma$ on $\partial\Omega$. By writing $u = z + \xi$, the problem of finding u becomes equivalent to the problem of finding $z \in H_{0,\Gamma_1}^1(\Omega)$ such that

$$\int_{\Omega} \langle \nabla z, \nabla \varphi \rangle = \int_{\Omega} f \varphi - \int_{\Omega} \langle \nabla \xi, \nabla \varphi \rangle + \eta(\varphi) \quad \text{for any } \varphi \in H_{0,\Gamma_1}^1(\Omega), \quad (4.3)$$

i.e., such that

$$B(z, \varphi) = L(\varphi) \text{ for any } \varphi \in H_{0,\Gamma_1}^1(\Omega),$$

where, for $w \in H_{0,\Gamma_1}^1(\Omega)$ and $\varphi \in H_{0,\Gamma_1}^1(\Omega)$,

$$B(w, \varphi) := \int_{\Omega} \langle \nabla w, \nabla \varphi \rangle \quad \text{and} \quad L(\varphi) := \int_{\Omega} f \varphi - \int_{\Omega} \langle \nabla \xi, \nabla \varphi \rangle + \eta(\varphi).$$

Since B is a continuous and coercive bilinear form on $H_{0,\Gamma_1}^1(\Omega) \times H_{0,\Gamma_1}^1(\Omega)$ and $L \in (H_{0,\Gamma_1}^1(\Omega))'$, the Lax Milgram theorem gives the existence and uniqueness of the solution $z \in H_{0,\Gamma_1}^1(\Omega)$ of (4.3), and that it satisfies $\|z\|_{H_{0,\Gamma_1}^1(\Omega)} \leq c' \|L\|_{(H_{0,\Gamma_1}^1(\Omega))'}$ for some positive constant c' independent of f , τ , and η . Then problem (4.1) has a unique solution $u \in H^1(\Omega)$ given by $u := z + \xi$. And, since

$$\|L\|_{(H_{0,\Gamma_1}^1(\Omega))'} \leq \|f\|_{(H_0^1(\Omega))'} + \|\xi\|_{H^1(\Omega)} + \|\eta\|_{(H_{0,\Gamma_1}^1(\Omega))'}$$

and (see [43, Section 7.9.3, formula (7.48)])

$$\|\tau\|_{H^{\frac{1}{2}}(\Gamma_1)} = \|\sigma\|_{H^{\frac{1}{2}}(\Gamma_1)} = \inf \left\{ \|w\|_{H^1(\Omega)} : w \in H^1(\Omega) \text{ and } w = \sigma \text{ on } \partial\Omega \right\},$$

we get (4.2).

(ii) From (i) it follows that if $f \in L^2(\Omega, \rho_{\Gamma_1}^2(x)dx)$, $\tau \in H^{\frac{1}{2}}(\Gamma_1)$ and if $\eta : \Gamma_2 \rightarrow \mathbb{R}$ belongs to $(H_{0,\Gamma_1}^1(\Omega))'$, then the problem

$$\begin{cases} -\Delta u = f & \text{in } \Omega, \\ u = \tau & \text{on } \Gamma_1, \\ \frac{\partial u}{\partial \nu} = \eta & \end{cases} \quad (4.4)$$

has a unique weak solution $u \in H^1(\Omega)$.

Definition 4.2. For $\tau \in H^{\frac{1}{2}}(\Gamma_1)$, $\eta \in (H_{0,\Gamma_1}^1(\Omega))'$, let

$$S_{\tau,\eta} : L^2(\Omega, \rho_{\Gamma_1}^2(x)dx) \rightarrow H^1(\Omega)$$

be the solution operator of the problem

$$\begin{cases} \int_{\Omega} \langle \nabla u, \nabla \varphi \rangle = \int_{\Omega} h\varphi + \eta(\varphi) & \text{for any } \varphi \in H_{0,\Gamma_1}^1(\Omega), \\ u = \tau \text{ on } \Gamma_1, \end{cases} \quad (4.5)$$

defined by $S_{\tau,\eta}(h) = u$, where u is the weak solution of (4.5). If no confusion arises, we will write S instead of $S_{\tau,\eta}$.

Lemma 4.3. Let $\tau \in H^{\frac{1}{2}}(\Gamma_1)$, $\eta \in (H_{0,\Gamma_1}^1(\Omega))'$. Then:

- (i) $S : L^2(\Omega, \rho_{\Gamma_1}^2(x)dx) \rightarrow H^1(\Omega)$ is continuous,
- (ii) $S : L^2(\Omega, \rho_{\Gamma_1}^2(x)dx) \rightarrow L^2(\Omega)$ is a continuous and compact operator,
- (iii) if h_1 and h_2 belong to $L^2(\Omega, \rho_{\Gamma_1}^2(x)dx)$ and $h_1 \leq h_2$ then $S(h_1) \leq S(h_2)$,
- (iv) if, in addition, $\tau \geq 0$ and $\eta \geq 0$ then $S(h) \geq 0$ for any nonnegative $h \in L^2(\Omega, \rho_{\Gamma_1}^2(x)dx)$.

Proof. If h_1 and h_2 belong to $L^2(\Omega, \rho_{\Gamma_1}^2(x)dx)$ and if $u_1 = S(h_1)$ and $u_2 = S(h_2)$ then $u_1 - u_2$ satisfies

$$\begin{cases} \int_{\Omega} \langle \nabla(u_1 - u_2), \nabla \varphi \rangle = \int_{\Omega} (h_1 - h_2)\varphi & \text{for any } \varphi \in H_{0,\Gamma_1}^1(\Omega), \\ u_1 - u_2 = 0 \text{ on } \Gamma_1, \end{cases} \quad (4.6)$$

and so, by (4.2),

$$\|u_1 - u_2\|_{H^1(\Omega)} \leq c \|h_1 - h_2\|_{(H^1(\Omega))'} \leq c' \|h_1 - h_2\|_{L^2(\Omega, \rho_{\Gamma_1}^2(x)dx)}$$

with c and c' positive constants independent of h_1 and h_2 . Then (i) holds, and (ii) follows from (i) and from the fact that the inclusion $H^1(\Omega) \hookrightarrow L^2(\Omega)$ is continuous and compact.

To prove (iii) observe that if $u_1 = S(h_1)$ and $u_2 = S(h_2)$, then, from (4.6) used with $\varphi = (u_1 - u_2)^+$, we get $\int_{\Omega} |\nabla((u_1 - u_2)^+)|^2 \leq 0$ and so

$$\int_{\Omega} |\nabla((u_1 - u_2)^+)|^2 = 0.$$

Then, by the Poincaré inequality of Remark 2.1(ii), $(u_1 - u_2)^+ = 0$ and thus $u_1 \leq u_2$. To see (iv) suppose $\eta \geq 0$ and $0 \leq h \in L^2(\Omega, \rho_{\Gamma_1}^2(x)dx)$. Let $u = S(h)$. Then

$$\int_{\Omega} \langle \nabla u, \nabla \varphi \rangle = \int_{\Omega} h\varphi + \eta(\varphi) \quad \text{for any } \varphi \in H_{0,\Gamma_1}^1(\Omega).$$

We take $\varphi = -u^-$ to obtain

$$\int_{\Omega} |\nabla u^-|^2 = - \int_{\Omega} \langle \nabla u, \nabla u^- \rangle = - \int_{\Omega} h u^- - \int_{\Gamma_2} \eta u^- \leq 0.$$

Then $\int_{\Omega} |\nabla u^-|^2 = 0$ and so, by Remark 2.1(ii), $u^- = 0$. \square

Proof of Theorem 1.6. Let $u_{\tau} \in H^1(\Omega)$ be the solution of the problem

$$\begin{cases} -\Delta u_{\tau} = g(\cdot, u_{\tau}) & \text{in } \Omega, \\ u_{\tau} = \tau & \text{on } \Gamma_1, \\ u_{\tau} = 0 & \text{on } \Gamma_2 \end{cases} \quad (4.7)$$

given by Theorem 1.3. Let $\eta : \Gamma_2 \rightarrow \mathbb{R}$ be such that $\eta \in (H_{0,\Gamma_1}^1(\Omega))'$ and $\eta \geq \frac{\partial u_{\tau}}{\partial \nu}_{\Gamma_2}$ in $(H_{0,\Gamma_1}^1(\Omega))'$, let $\Phi \in H^1(\Omega)$ be the solution of the problem

$$\begin{cases} \int_{\Omega} \langle \nabla \Phi, \nabla \varphi \rangle = \left(\eta - \frac{\partial u_{\tau}}{\partial \nu}_{\Gamma_2} \right) (\varphi) \text{ for any } \varphi \in H_{0,\Gamma_1}^1(\Omega), \\ \Phi = 0 \text{ on } \Gamma_1 \end{cases} \quad (4.8)$$

(by Remark 4.1(i), there exists such a unique Φ), and let $z = \Phi + u_{\tau}$. Since $\eta - \frac{\partial u_{\tau}}{\partial \nu}_{\Gamma_2} \geq 0$, Lemma 4.3(iv) gives that $\Phi \geq 0$, thus $u_{\tau} \leq z$. Note that for any nonnegative $\varphi \in H_{0,\Gamma_1}^1(\Omega)$,

$$\int_{\Omega} \langle \nabla u_{\tau}, \nabla \varphi \rangle = \int_{\Omega} g(\cdot, u_{\tau}) \varphi + \frac{\partial u_{\tau}}{\partial \nu}_{\Gamma_2} (\varphi) \leq \int_{\Omega} g(\cdot, u_{\tau}) \varphi + \eta(\varphi),$$

and so, for any nonnegative $\varphi \in H_{0,\Gamma_1}^1(\Omega)$,

$$\begin{aligned} \int_{\Omega} \langle \nabla z, \nabla \varphi \rangle &= \int_{\Omega} \langle \nabla u_{\tau}, \nabla \varphi \rangle + \int_{\Omega} \langle \nabla \Phi, \nabla \varphi \rangle \\ &= \int_{\Omega} g(\cdot, u_{\tau}) \varphi + \frac{\partial u_{\tau}}{\partial \nu}_{\Gamma_2} (\varphi) + \left(\eta - \frac{\partial u_{\tau}}{\partial \nu}_{\Gamma_2} \right) (\varphi) \\ &\geq \int_{\Omega} g(\cdot, \Phi + u_{\tau}) \varphi + \eta(\varphi) = \int_{\Omega} g(\cdot, z) \varphi + \eta(\varphi), \end{aligned}$$

where we have used that $\frac{\partial \Phi}{\partial \nu}_{\Gamma_2} (\varphi) = \int_{\Omega} \langle \nabla \Phi, \nabla \varphi \rangle$ and that $g = g(x, s)$ is nonincreasing in s .

Then for any nonnegative $\varphi \in H_{0,\Gamma_1}^1(\Omega)$,

$$\int_{\Omega} \langle \nabla u_{\tau}, \nabla \varphi \rangle \leq \int_{\Omega} g(\cdot, u_{\tau}) \varphi + \eta(\varphi) \quad (4.9)$$

and

$$\int_{\Omega} \langle \nabla z, \nabla \varphi \rangle \geq \int_{\Omega} g(\cdot, z) \varphi + \eta(\varphi). \quad (4.10)$$

To prove the existence assertion of the theorem we will show that problem (1.2) has a solution u^* such that $u_{\tau} \leq u^* \leq z$.

As in the proof of [35, Theorem 1.1] we define $\bar{g} : \Omega \times \mathbb{R} \rightarrow \mathbb{R}$ by

$$\bar{g}(x, s) := \begin{cases} g(x, u_{\tau}(x)) & \text{if } s \leq u_{\tau}(x), \\ g(x, s) & \text{if } u_{\tau}(x) < s < z(x), \\ g(x, z(x)) & \text{if } s \geq z(x). \end{cases}$$

It is easy to check that \bar{g} is a nonnegative Carathéodory function on $\Omega \times \mathbb{R}$ (because g is a Carathéodory function on $\Omega \times (0, \infty)$ and u_{τ}, z are measurable functions) and that $\bar{g}(x, s)$ is nonincreasing in s . Moreover, if $E \subset \Omega$ is the set given by the condition (H2) then $\bar{g}(x, s) > 0$ for any $x \in E$ and $s > 0$. Also, since $u_{\tau} \leq z$ and, taking into account that, by Theorem 1.3, $u_{\tau} \geq c\rho_{\partial\Omega}$ for some $c \in (0, \infty)$ and that $\bar{g}(x, s)$ is nonnegative and nonincreasing in s , we obtain that

$$0 \leq \bar{g}(\cdot, s) \leq \bar{g}(\cdot, u_{\tau}) = g(\cdot, u_{\tau}) \leq g(\cdot, c\rho_{\partial\Omega}) \quad (4.11)$$

for any $s \in \mathbb{R}$, and so, for any $v \in L^2(\Omega)$, $0 \leq \rho_{\Gamma_1} \bar{g}(\cdot, v) \leq \rho_{\Gamma_1} g(\cdot, c\rho_{\partial\Omega})$, and, by (H3'), $\rho_{\Gamma_1} g(\cdot, c\rho_{\partial\Omega}) \in L^2(\Omega)$. Therefore, taking into account the Hardy inequality of Lemma 2.2 we have, for any $\varphi \in H_{0,\Gamma_1}^1(\Omega)$,

$$\int_{\Omega} |\bar{g}(\cdot, v) \varphi| = \int_{\Omega} \rho_{\Gamma_1} \bar{g}(\cdot, v) \left| \frac{\varphi}{\rho_{\Gamma_1}} \right| \leq \|\rho_{\Gamma_1} \bar{g}(\cdot, v)\|_2 \left\| \frac{\varphi}{\rho_{\Gamma_1}} \right\|_2 \leq c' \|\varphi\|_{H^1(\Omega)}$$

with c' a positive constant independent of v and φ . Thus $\bar{g}(\cdot, v) \in (H_{0,\Gamma_1}^1(\Omega))'$ and

$$\|\bar{g}(\cdot, v)\|_{(H_{0,\Gamma_1}^1(\Omega))'} \leq c' \quad (4.12)$$

with c' independent of v . Following the lines of the proof of [35, Theorem 1.1] we consider the operator $T : L^2(\Omega) \rightarrow L^2(\Omega)$ defined by

$$T(v) := S(\bar{g}(\cdot, v)).$$

with S given by Definition 4.2.

We prove that:

- (1) T is continuous,
- (2) T is a compact operator,
- (3) there exists $R > 0$ such that $T(L^2(\Omega)) \subset \bar{B}$, where \bar{B} is the closed ball in $L^2(\Omega)$ centered at 0 and with radius R .

To prove (1) we proceed similarly to the proof of Lemma 3.1(iii). It is enough to see that if $v \in L^2(\Omega)$, and if $\{v_j\}_{j \in \mathbb{N}}$ is a sequence in $L^2(\Omega)$ that converges to v in $L^2(\Omega)$, then there exists a subsequence $\{v_{j_k}\}_{k \in \mathbb{N}}$ such that $\{T(v_{j_k})\}_{k \in \mathbb{N}}$ converges to $T(v)$ in $L^2(\Omega)$. Let $v \in L^2(\Omega)$, and let $\{v_j\}_{j \in \mathbb{N}}$ be a sequence in $L^2(\Omega)$ which converges to v in $L^2(\Omega)$, then there exists a subsequence $\{v_{j_k}\}_{k \in \mathbb{N}}$ such that $\{v_{j_k}\}_{k \in \mathbb{N}}$ converges to v a.e. in Ω . Thus, since \bar{g} is a Carathéodory function, $\{\bar{g}(\cdot, v_{j_k})\}_{k \in \mathbb{N}}$ converges to $\bar{g}(\cdot, v)$ a.e. in Ω . Then $\lim_{k \rightarrow \infty} |\bar{g}(\cdot, v_{j_k}) - \bar{g}(\cdot, v)|^2 = 0$ a.e. in Ω . By (4.11), $|\bar{g}(\cdot, v_{j_k}) - \bar{g}(\cdot, v)|^2 \leq 4g^2(\cdot, c\rho_{\partial\Omega})$, and, by (H3') we have $\int_{\Omega} \rho_{\Gamma_1}^2 g^2(\cdot, c\rho_{\partial\Omega}) < \infty$. Then, by the Lebesgue dominated convergence theorem, $\{\bar{g}(\cdot, v_{j_k})\}_{k \in \mathbb{N}}$ converges to $\bar{g}(\cdot, v)$ in $L^2(\Omega, \rho_{\Gamma_1}^2(x)dx)$. Then, by Lemma 4.3(ii), $\{S(\bar{g}(\cdot, v_{j_k}))\}_{k \in \mathbb{N}}$ converges to $S(\bar{g}(\cdot, v))$ in $L^2(\Omega)$, i.e., $\{T(v_{j_k})\}_{k \in \mathbb{N}}$ converges to $T(v)$ in $L^2(\Omega)$. Thus (1) holds.

To see (2) note that, by (4.11), $\{\bar{g}(\cdot, v_j)\}_{j \in \mathbb{N}}$ is bounded in $L^2(\Omega, \rho_{\Gamma_1}^2(x)dx)$ for any sequence $\{v_j\}_{j \in \mathbb{N}}$ in $L^2(\Omega)$, and that $S : L^2(\Omega, \rho_{\Gamma_1}^2(x)dx) \rightarrow L^2(\Omega)$ is compact.

To see (3) observe that, by (4.2) and (4.12), we have, for any $v \in L^2(\Omega)$,

$$\begin{aligned} \|T(v)\|_2 &= \|S(\bar{g}(\cdot, v))\|_2 \\ &\leq c(\|\bar{g}(\cdot, v)\|_{(H_0^1(\Omega))'} + \|\tau\|_{H^{\frac{1}{2}}(\Gamma_1)} + \|\eta\|_{(H_{0,\Gamma_1}^1(\Omega))'}) \\ &\leq c(c' + \|\tau\|_{H^{\frac{1}{2}}(\Gamma_1)} + \|\eta\|_{(H_{0,\Gamma_1}^1(\Omega))'}) \end{aligned}$$

with c and c' positive constants independent of v .

Now, as in [35, Theorem 1.1], from (1), (2), (3) and the Schauder fixed point theorem, there exists $u^* \in L^2(\Omega)$ such that $T(u^*) = u^*$, i.e., such that

$$\begin{cases} -\Delta u^* = \bar{g}(\cdot, u^*) & \text{in } \Omega, \\ u^* = \tau & \text{on } \Gamma_1, \\ \frac{\partial u^*}{\partial \nu} = \eta & \text{on } \Gamma_2. \end{cases} \quad (4.13)$$

To complete the proof of the existence assertion of the theorem it suffices to see that $u_\tau \leq u^* \leq z$ (because in such a case $\bar{g}(\cdot, u^*) = g(\cdot, u^*)$ and, by Theorem 1.3, $u_\tau \geq c\rho_{\partial\Omega}$ for some positive constant c). From (4.10), (4.13), and since $\bar{g}(\cdot, z) = g(\cdot, z)$ we have,

for any nonnegative $\varphi \in H_{0,\Gamma_1}^1(\Omega)$,

$$\begin{aligned} \int_{\Omega} \langle \nabla(z - u^*), \nabla \varphi \rangle &= \int_{\Omega} \langle \nabla z, \nabla \varphi \rangle - \int_{\Omega} \langle \nabla u^*, \nabla \varphi \rangle \\ &\geq \int_{\Omega} g(\cdot, z) \varphi + \eta(\varphi) - \int_{\Omega} \bar{g}(\cdot, u^*) - \eta(\varphi) \\ &= \int_{\Omega} (\bar{g}(\cdot, z) - \bar{g}(\cdot, u^*)) \varphi \end{aligned}$$

which, by taking $\varphi = (z - u^*)^-$ gives

$$\int_{\Omega} |\nabla((z - u^*)^-)|^2 \leq - \int_{\Omega} (\bar{g}(\cdot, z) - \bar{g}(\cdot, u^*)) (z - u^*)^- \leq 0,$$

the last inequality because $\bar{g}(x, s)$ is nonincreasing in s . Then, by Remark 2.1(ii), $(z - u^*)^- = 0$ and so $u^* \leq z$.

Similarly, from (4.7), (4.13) and since $\bar{g}(\cdot, u_\tau) = g(\cdot, u_\tau)$, we have, for any nonnegative $\varphi \in H_{0,\Gamma_1}^1(\Omega)$,

$$\begin{aligned} \int_{\Omega} \langle \nabla(u^* - u_\tau), \nabla \varphi \rangle &= \int_{\Omega} (\bar{g}(\cdot, u^*) - \bar{g}(\cdot, u_\tau)) \varphi + \eta(\varphi) - \frac{\partial u_\tau}{\partial \nu} \Big|_{\Gamma_2}(\varphi) \\ &\leq \int_{\Omega} (\bar{g}(\cdot, u^*) - \bar{g}(\cdot, u_\tau)) \varphi, \end{aligned} \tag{4.14}$$

the last inequality by our assumption that $\eta \geq \frac{\partial u_\tau}{\partial \nu} \Big|_{\Gamma_2}$. Observe that $u^* - u_\tau \in H_{0,\Gamma_1}^1(\Omega)$ and that, since $\bar{g}(\cdot, s)$ is nonincreasing in s ,

$$(\bar{g}(\cdot, u^*) - \bar{g}(\cdot, u_\tau))(u^* - u_\tau)^- \geq 0.$$

Thus, taking $\varphi = -(u^* - u_\tau)^-$ in (4.14) we obtain $\int_{\Omega} |\nabla((u^* - u_\tau)^-)|^2 = 0$, which implies $(u^* - u_\tau)^- = 0$ and so $u_\tau \leq u^*$.

Suppose that $w \in H^1(\Omega)$ is another solution of (1.2). Then $u^* - w \in H_{0,\Gamma_1}^1(\Omega)$ and, in a weak sense,

$$\begin{cases} -\Delta(u^* - w) = g(\cdot, u^*) - g(\cdot, w) & \text{in } \Omega, \\ u^* - w = 0 & \text{on } \Gamma_1, \\ \frac{\partial(u^* - w)}{\partial \nu} = 0 & \text{on } \Gamma_2, \end{cases} \tag{4.15}$$

that is,

$$\int_{\Omega} \langle \nabla(u^* - w), \nabla \varphi \rangle = \int_{\Omega} (g(\cdot, u^*) - g(\cdot, w)) \varphi \text{ for any } \varphi \in H_{0,\Gamma_1}^1(\Omega), \tag{4.16}$$

Now, $g(x, \cdot)$ is nonincreasing on $(0, \infty)$ for a.e. $x \in \Omega$, and so

$$g(\cdot, u^*) - g(\cdot, w)(u^* - w) \leq 0 \text{ a.e. in } \Omega.$$

Thus, taking $\varphi = u^* - w$ in (4.16), we get

$$\int_{\Omega} |\nabla(u^* - w)|^2 = \int_{\Omega} (g(\cdot, u^*) - g(\cdot, w))(u^* - w) \leq 0$$

and so $\|\nabla(u^* - w)\|_2 = 0$. Then, by Remark 2.1(ii), $u^* = w$ in Ω . This concludes the proof of the part (i) of the theorem.

To see (ii), suppose that $\eta < \frac{\partial u_{\tau}}{\partial \nu}_{\Gamma_2}$ and that u is a weak solution of problem (1.2). Then, for any nonnegative $\varphi \in H^1_{0,\Gamma_1}(\Omega)$,

$$\int_{\Omega} \langle \nabla u, \nabla \varphi \rangle = \int_{\Omega} g(\cdot, u)\varphi + \eta(\varphi) \leq \int_{\Omega} g(\cdot, u)\varphi + \frac{\partial u_{\tau}}{\partial \nu}_{\Gamma_2}(\varphi),$$

and

$$\int_{\Omega} \langle \nabla u_{\tau}, \nabla \varphi \rangle = \int_{\Omega} g(\cdot, u_{\tau})\varphi + \frac{\partial u_{\tau}}{\partial \nu}_{\Gamma_2}(\varphi).$$

Thus, for any nonnegative $\varphi \in H^1_{0,\Gamma_1}(\Omega)$,

$$\int_{\Omega} \langle \nabla(u - u_{\tau}), \nabla \varphi \rangle \leq \int_{\Omega} (g(\cdot, u) - g(\cdot, u_{\tau}))\varphi.$$

Now we take $\varphi = (u - u_{\tau})^+$ to obtain that

$$\int_{\Omega} |\nabla((u - u_{\tau})^+)|^2 \leq \int_{\Omega} (g(\cdot, u) - g(\cdot, u_{\tau}))(u - u_{\tau})^+ \leq 0,$$

the last inequality because $g(x, s)$ is nonincreasing in s . Thus $(u - u_{\tau})^+ = 0$ and so $u \leq u_{\tau}$. Since u is nonnegative and $u_{\tau} = 0$ on Γ_2 we conclude that $u = 0$ on Γ_2 . Then u is a solution of problem (1.13) and, by Theorem 1.3, this problem has a unique solution. Then $u = u_{\tau}$, and so $\eta = \frac{\partial u_{\tau}}{\partial \nu}_{\Gamma_2}$, which is a contradiction. Therefore no such a solution u exists. \square

Lemma 4.4. *If $0 \leq f \in L^2(\Omega, d_{\Gamma_1}^2(x)dx)$, $0 \leq \tau \in H^{\frac{1}{2}}(\Gamma_1)$, and if $u \in H^1(\Omega)$ is the weak solution of the problem*

$$\begin{cases} -\Delta u = f & \text{in } \Omega, \\ u = \tau & \text{on } \Gamma_1, \\ u = 0 & \text{on } \Gamma_2, \end{cases}$$

then $\frac{\partial u}{\partial \nu}_{\Gamma_2} \leq 0$.

Proof. Let $\Psi : \partial\Omega \rightarrow \mathbb{R}$ be defined by $\Psi = \tau$ on Γ_1 and $\Psi = 0$ on Γ_2 . Then $0 \leq \Psi \in H^{\frac{1}{2}}(\partial\Omega)$ and thus there exists $\tilde{\Psi} \in H^1(\Omega)$ such that $\tilde{\Psi} = \Psi$ on $\partial\Omega$. By replacing $\tilde{\Psi}$ by $\tilde{\Psi}^+$ if necessary, we can assume that $\tilde{\Psi} \geq 0$ in Ω . Now, Ω is a bounded domain with C^2 boundary, and then $C^\infty(\bar{\Omega})$ is dense in $W^{1,2}(\Omega)$ (see [1, Theorem 3.18]). Then there exists a sequence $\{\tilde{\Psi}_j\}_{j \in \mathbb{N}} \subset C^\infty(\bar{\Omega})$ such that $\{\tilde{\Psi}_j\}_{j \in \mathbb{N}}$ converges to $\tilde{\Psi}$ in $H^1(\Omega)$. An inspection of the proof of [1, Theorem 3.18] shows that, since $\tilde{\Psi}$ is nonnegative, the functions $\tilde{\Psi}_j$ can be chosen nonnegative. For $\gamma > 0$, let $\Omega_{\Gamma_2, \gamma}$ and $A_{\Gamma_2, \gamma}$ be defined as in (2.1). Let δ be a positive number such that $\Gamma_1 \cap A_{\Gamma_2, 4\delta} = \emptyset$, and let $\phi \in C^\infty(\bar{\Omega})$ be such that $0 \leq \phi \leq 1$, $\phi = 0$ in $A_{\Gamma_2, \delta}$ and $\phi = 1$ in $\Omega_{\Gamma_2, 2\delta}$. Then $0 \leq \phi\tilde{\Psi}_j \in C^\infty(\bar{\Omega})$, $\phi\tilde{\Psi}_j = 0$ on Γ_2 , and $\{\phi\tilde{\Psi}_j\}_{j \in \mathbb{N}}$ converges to $\phi\tilde{\Psi}$ in $H^1(\Omega)$. For $j \in \mathbb{N}$, let $\Psi_j := \phi\tilde{\Psi}_j|_{\partial\Omega}$ and let $f_j : \Omega \rightarrow \mathbb{R}$ be defined by $f_j(x) := \min\{j, f(x)\}$. Then $\Psi_j = 0$ on Γ_2 , $\{\Psi_j|_{\Gamma_1}\}_{j \in \mathbb{N}}$ converges to τ in $H^{\frac{1}{2}}(\Gamma_1)$ and $\{f_j\}_{j \in \mathbb{N}}$ converges to f in $L^2(\Omega, d_{\Gamma_1}^2(x)dx)$. In particular, $\{f_j\}_{j \in \mathbb{N}}$ converges to f in $(H_{0, \Gamma_1}^1(\Omega))'$. Now, $f_j \in L^\infty(\Omega)$ and Ψ_j is the restriction to $\partial\Omega$ of a function in $C^\infty(\bar{\Omega})$. Then (see, e.g., [29, Theorem 2.4.2.5], see also [26, Theorem 9.15]), the problem

$$\begin{cases} -\Delta u_j = f_j & \text{in } \Omega, \\ u_j = \Psi_j & \text{on } \partial\Omega \end{cases} \quad (4.17)$$

has a unique strong solution $u_j \in \bigcap_{1 < p < \infty} W^{2,p}(\Omega) \subset C^1(\bar{\Omega})$. Since $f_j \geq 0$ and $\Psi_j \geq 0$ we have $u_j \geq 0$. Also, $u_j = 0$ on Γ_2 , and then the Hopf boundary lemma, as stated in [42, Theorem 1.1], gives that $\frac{\partial u_j}{\partial \nu}(x) < 0$ for any $x \in \Gamma_2$. On the other hand, $\{f_j\}_{j \in \mathbb{N}}$ converges to f in $(H_{0, \Gamma_1}^1(\Omega))'$ and $\{\Psi_j\}_{j \in \mathbb{N}}$ converges to Ψ in $H^{\frac{1}{2}}(\partial\Omega)$, then $\{u_j\}_{j \in \mathbb{N}}$ converges to u in $H^1(\Omega)$. Let φ be an arbitrary nonnegative function in $H_{0, \Gamma_1}^1(\Omega)$. From (4.17), we have $-\operatorname{div}(\varphi \nabla u_j) + \langle \nabla u_j, \nabla \varphi \rangle = f_j \varphi$ in Ω , and so, by the divergence theorem (as stated, for example, in [14, Lemma A.1]),

$$-\int_{\Gamma_2} \varphi \frac{\partial u_j}{\partial \nu} + \int_{\Omega} \langle \nabla u_j, \nabla \varphi \rangle = \int_{\Omega} f_j \varphi.$$

Then $\int_{\Omega} \langle \nabla u_j, \nabla \varphi \rangle - \int_{\Omega} f_j \varphi \geq 0$ and thus, taking into account that $\{\nabla u_j\}_{j \in \mathbb{N}}$ converges to ∇u in $L^2(\Omega, \mathbb{R}^n)$ and that $\{f_j\}_{j \in \mathbb{N}}$ converges to f in $(H_{0, \Gamma_1}^1(\Omega))'$, we get that $\int_{\Omega} \langle \nabla u, \nabla \varphi \rangle - \int_{\Omega} f \varphi \geq 0$. Then $\frac{\partial u}{\partial \nu}|_{\Gamma_2} \leq 0$. \square

Proof of Corollary 1.7. Let u_τ be the solution (given by Theorem 1.3) of problem (4.7). By Lemma 4.4, we have $\frac{\partial u}{\partial \nu}|_{\Gamma_2} \leq 0$. Then the corollary follows immediately from Theorem 1.6. \square

REFERENCES


- [1] R.A. Adams, *Sobolev Spaces*, Academic Press, New York, San Francisco, London, 1975.
- [2] R.P. Agarwal, K. Perera, D. O'Regan, *A variational approach to singular quasilinear problems with sign changing nonlinearities*, Appl. Anal. **85** (2006), no. 10, 1201–1206.
- [3] I. Bachar, H. Măagli, V. Rădulescu, *Singular solutions of a nonlinear elliptic equation in a punctured domain*, Electron. J. Qual. Theory Differ. Equ. **2017**, Paper no. 94.
- [4] B. Bougherara, J. Giacomoni, *Existence of mild solutions for a singular parabolic equation and stabilization*, Adv. Nonlinear Anal. **4** (2015), no. 2, 123–134.
- [5] B. Bougherara, J. Giacomoni, J. Hernández, *Existence and regularity of weak solutions for singular elliptic problems*, Proceedings of the 2014 Madrid Conference on Applied Mathematics in honor of Alfonso Casal, 19–30, Electron. J. Differ. Equ. Conf. **22**, Texas State Univ., San Marcos, TX, 2015.
- [6] H. Brezis, *Functional Analysis, Sobolev Spaces and Partial Differential Equations*, Universitext, Springer, New York, 2011.
- [7] H. Brezis, X. Cabre, *Some simple nonlinear PDE's without solutions*, Boll. Unione Mat. Ital. Sez. B Artic. Ric. Mat. **1** (1998), no. 2, 223–262.
- [8] A. Callegari, A. Nachman, *A nonlinear singular boundary value problem in the theory of pseudoplastic fluids*, SIAM J. Appl. Math. **38** (1980), no. 2, 275–281.
- [9] Y. Chu, Y. Gao, W. Gao, *Existence of solutions to a class of semilinear elliptic problem with nonlinear singular terms and variable exponent*, J. Funct. Spaces (2016), Art. ID 9794739.
- [10] F. Cîrstea, M. Ghergu, V. Rădulescu, *Combined effects of asymptotically linear and singular nonlinearities in bifurcation problems of Lane–Emden–Fowler type*, J. Math. Pures Appl. **84** (2005), no. 4, 493–508.
- [11] M.M. Coclite, G. Palmieri, *On a singular nonlinear Dirichlet problem*, Comm. Partial Differential Equations **14** (1989), no. 10, 1315–1327.
- [12] D.S. Cohen, H.B. Keller, *Some positive problems suggested by nonlinear heat generators*, J. Math. Mech. **16** (1967), 1361–1376.
- [13] M.G. Crandall, P.H. Rabinowitz, L. Tartar, *On a Dirichlet problem with a singular nonlinearity*, Comm. Partial Differential Equations **2** (1977), no. 2, 193–222.
- [14] M. Cuesta, P. Takáč, *A strong comparison principle for positive solutions of degenerate elliptic equations*, Differential Integral Equations **13** (2000), no. 4–6, 721–746.
- [15] M.A. del Pino, *A global estimate for the gradient in a singular elliptic boundary value problem*, Proc. Roy. Soc. Edinburgh Sect. A **122** (1992), no. 3–4, 341–352.
- [16] J.I. Diaz, J. Hernández, *Positive and free boundary solutions to singular nonlinear elliptic problems with absorption: an overview and open problems*, Electron. J. Differ. Equ. Conf. **21** (2014), 31–44.
- [17] J.I. Diaz, J. Hernandez, J.M. Rakotoson, *On very weak positive solutions to some semilinear elliptic problems with simultaneous singular nonlinear and spatial dependence terms*, Milan J. Math. **79** (2011), Article no. 233.

- [18] J.I. Diaz, J.M. Morel, L. Oswald, *An elliptic equation with singular nonlinearity*, Comm. Partial Differential Equations **12** (1987), no. 12, 1333–1344.
- [19] L. Dupaigne, M. Ghergu, V. Rădulescu, *Lane–Emden–Fowler equations with convection and singular potential*, J. Math. Pures Appl. **87** (2007), no. 6, 563–581.
- [20] P. Feng, *On the structure of positive solutions to an elliptic problem arising in thin film equations*, J. Math. Anal. Appl. **370** (2010), no. 2, 573–583.
- [21] W. Fulks, J.S. Maybee, *A singular nonlinear equation*, Osaka J. Math. **12** (1960), 1–19.
- [22] L. Gasiński, N.S. Papageorgiou, *Nonlinear elliptic equations with singular terms and combined nonlinearities*, Ann. Henri Poincaré **13** (2012), no. 3, 481–512.
- [23] M. Ghergu, V.D. Rădulescu, *Singular Elliptic Problems: Bifurcation and Asymptotic Analysis*, Oxford Lecture Series in Mathematics and its Applications, vol. 37, The Clarendon Press, Oxford University Press, Oxford, 2008.
- [24] M. Ghergu, V.D. Rădulescu, *Nonlinear PDEs. Mathematical Models in Biology, Chemistry and Population Genetics*, Springer-Verlag, Berlin, Heidelberg, New York, 2012.
- [25] M. Ghergu, V.D. Rădulescu, *Bifurcation and asymptotics for the Lane–Emden–Fowler equation*, C. R. Math. Acad. Sci. Paris **337** (2003), no. 4, 259–264.
- [26] D. Gilbarg, N.S. Trudinger, *Elliptic Partial Differential Equations of Second Order*, Springer-Verlag, Berlin, Heidelberg, New York, 2001.
- [27] T. Godoy, *Strong solutions for singular Dirichlet elliptic problems*, Electron. J. Qual. Theory Differ. Equ. **2022**, Paper no. 40, 1–20.
- [28] T. Godoy, A. Guerin, *Positive weak solutions of elliptic Dirichlet problems with singularities in both the dependent and the independent variables*, Electron. J. Qual. Theory Differ. Equ. **2019**, Paper no. 54, 1–17.
- [29] P. Grisvard, *Elliptic Problems in Nonsmooth Domains*, Pitman Advanced Publishing Program, Marshfield, Massachusetts, 1985.
- [30] J. Janus, J. Myjak, *A generalizad Emden–Fowler equation with a negative exponent*, Nonlinear Anal. **23** (1994), no. 8, 953–970.
- [31] A.C. Lazer, P.J. McKenna, *On a singular nonlinear elliptic boundary-value problem*, Proc. Amer. Math. Soc. **111** (1991), no. 3, 721–730.
- [32] Q. Li, W. Gao, *Existence of weak solutions to a class of singular elliptic equations*, Mediterr. J. Math. **13** (2016), no. 6, 4917–4927.
- [33] H. Mâagli, *Asymptotic behavior of positive solutions of a semilinear Dirichlet problem*, Nonlinear Anal. **74** (2011), no. 9, 2941–2947.
- [34] L. Ma, J.C. Wei, *Properties of positive solutions to an elliptic equation with negative exponent*, J. Funct. Anal. **254** (2008), no. 4, 1058–1087.
- [35] M. Montenegro, A.C. Ponce, *The sub-supersolution method for weak solutions*, Proc. Amer. Math. Soc. **136** (2008), no. 7, 2429–2438.
- [36] F. Oliva, F. Petitta, *Finite and infinite energy solutions of singular elliptic problems: existence and uniqueness*, J. Differential Equations **264** (2018), no. 1, 311–340.

- [37] N.S. Papageorgiou, G. Smyrlis, *Nonlinear elliptic equations with singular reaction*, Osaka J. Math. **53** (2016), no. 2, 489–514.
- [38] N.S. Papageorgiou, V.D. Rădulescu, D.D. Repovš, *Nonlinear Analysis, Theory and Methods*, Springer Monographs in Mathematics, Springer, Cham, 2019.
- [39] K. Perera, E.A. Silva, *Multiple positive solutions of singular p -Laplacian problems via variational methods*, [in:] Differential & Difference Equations and Applications, Hindawi Publ. Corp., New York, 2006, pp. 915–924.
- [40] K. Perera, E.A. Silva, *On singular p -Laplacian problems*, Differential Integral Equations **20** (2007), no. 1, 105–120.
- [41] V.D. Rădulescu, *Singular phenomena in nonlinear elliptic problems. From blow-up boundary solutions to equations with singular nonlinearities*, [in:] Handbook of Differential Equations: Stationary Partial Differential Equations, vol. 4, Elsevier/North-Holland, Amsterdam, 2007, pp. 483–591.
- [42] J. Sabina de Lis, *Hopf maximum principle revisited*, Electron. J. Differential Equations **2015**, no. 115, 9 pp.
- [43] S. Salsa, *Partial Differential Equations in Action – From Modelling to Theory*, Springer, Milan, Berlin, Heidelberg, New York, 2008.
- [44] J. Shi, M. Yao, *On a singular nonlinear semilinear elliptic problem*, Proc. Roy. Soc. Edinburgh Sect. A **128** (1998), no. 6, 1389–1401.
- [45] J. Shi, M. Yao, *Positive solutions for elliptic equations with singular nonlinearity*, Electron. J. Qual. Theory Differ. Equ. **2005**, no. 04, 11 pp.
- [46] C.A. Stuart, *Existence and approximation of solutions of nonlinear elliptic problems*, Mathematics Report, Battelle Advanced Studies Center, Geneva, Switzerland, vol. 86, 1976.
- [47] S. Yijing, Z. Duanzhi, *The role of the power 3 for elliptic equations with negative exponents*, Calc. Var. Partial Differential Equations **49** (2014), no. 3–4, 909–922.
- [48] Z. Zhang, *The asymptotic behaviour of the unique solution for the singular Lane–Emden–Fowler equation*, J. Math. Anal. Appl. **312** (2005), no. 1, 33–43.

Tomas Godoy

godoy@famaf.unc.edu.ar

 <https://orcid.org/0000-0002-8804-9137>

Universidad Nacional de Córdoba

Facultad de Matemática, Astronomía, Física y Computación

Av. Medina Allende s/n, Ciudad Universitaria, 5000 Córdoba, Argentina

Received: August 7, 2022.

Revised: November 21, 2022.

Accepted: November 25, 2022.