

Mustapha Chellali

## BOUNDS ON THE 2-DOMINATION NUMBER IN CACTUS GRAPHS

**Abstract.** A 2-dominating set of a graph  $G$  is a set  $D$  of vertices of  $G$  such that every vertex not in  $S$  is dominated at least twice. The minimum cardinality of a 2-dominating set of  $G$  is the 2-domination number  $\gamma_2(G)$ . We show that if  $G$  is a nontrivial connected cactus graph with  $k(G)$  even cycles ( $k(G) \geq 0$ ), then  $\gamma_2(G) \geq \gamma_t(G) - k(G)$ , and if  $G$  is a graph of order  $n$  with at most one cycle, then  $\gamma_2(G) \geq (n + \ell - s)/2$  improving Fink and Jacobson's lower bound for trees with  $\ell > s$ , where  $\gamma_t(G)$ ,  $\ell$  and  $s$  are the total domination number, the number of leaves and support vertices of  $G$ , respectively. We also show that if  $T$  is a tree of order  $n \geq 3$ , then  $\gamma_2(T) \leq \beta(T) + s - 1$ , where  $\beta(T)$  is the independence number of  $T$ .

**Keywords:** 2-domination number, total domination number, independence number, cactus graphs, trees.

**Mathematics Subject Classification:** 05C69.

### 1. INTRODUCTION

Let  $G = (V(G), E(G))$  be a graph with the vertex set  $V(G)$  and edge set  $E(G)$ . The *degree* of a vertex  $v$ ,  $\deg_G(v)$ , is the number of vertices adjacent to  $v$ . A vertex of degree one is called a *leaf* and its neighbor is called a *support vertex*. If  $u$  is a support vertex, then  $L_u$  will denote the set of leaves attached at  $u$ . A graph  $G$  is called a *cactus graph* if each edge of  $G$  is contained in at most one cycle. A cactus graph having one cycle is called a *unicycle graph* and a connected cactus graph with no cycles is called a *tree*. A tree  $T$  is a *double star* if it contains exactly two vertices that are not leaves. A double star with  $p$  and  $q$  leaves attached at each support vertex, respectively, is denoted by  $S_{p,q}$ . For a graph  $G$  we denote by  $n(G)$ ,  $\ell(G)$  and  $s(G)$  the number of vertices, leaves and support vertices of  $G$ , respectively (we use  $n$ ,  $\ell$  and  $s$  if there is no ambiguity).

We are interested in a variation of domination in graphs, called 2-domination. A subset  $D$  of  $V(G)$  is a *2-dominating set* if every vertex not in  $S$  is adjacent to at least 2 vertices of  $D$ . The *2-domination number*  $\gamma_2(G)$  is the minimum cardinality of a 2-dominating set of  $G$ . Note that every graph  $G$  has a 2-dominating set since  $V(G)$  is such a set. The concept of 2-domination was introduced by Fink and Jacobson [5, 6], and studied for example in [1, 2]. The *independence number*  $\beta(G)$  of a graph  $G$  is the maximum cardinality among the independent sets of vertices of  $G$  and the *total domination number*  $\gamma_t(G)$  of a graph  $G$  is the minimum cardinality of a dominating set whose induced subgraph contains no isolated vertices. For more details on domination and its variations see the books of Haynes, Hedetniemi and Slater [8, 9].

In this paper we show that if  $G$  is a nontrivial connected cactus graph with  $k(G)$  even cycles ( $k(G) \geq 0$ ), then  $\gamma_2(G) \geq \gamma_t(G) - k(G)$ , and if  $G$  is a graph of order  $n$  with at most one cycle, then  $\gamma_2(G) \geq (n + \ell - s)/2$ . Finally, we show that every tree  $T$  of order at least three satisfies  $\gamma_2(T) \leq \beta(T) + s - 1$ .

## 2. LOWER BOUNDS

Before presenting our main results, we make a couple of straightforward observations.

**Observation 1.** *Every 2-dominating set of a graph  $G$  contains every leaf.*

**Observation 2.** *Let  $T$  be a tree obtained from a nontrivial tree  $T'$  by adding a star  $K_{1,p}$  with the center vertex  $v$  attached by an edge  $vw$  at a vertex  $w$  of  $T'$ . Then:*

- 1)  $\gamma_2(T') \leq \gamma_2(T) - |L_v|$ , with equality if  $p \geq 2$  or  $w$  is a leaf in  $T'$ .
- 2)  $\beta(T') = \beta(T) - |L_v|$ .

*Proof.* 1) Let  $D$  be a  $\gamma_2(T)$ -set. By Observation 1,  $D$  contains  $L_v$  and without loss of generality  $v \notin D$  (else replace  $v$  by  $w$  in  $D$ ). Thus  $D - L_v$  is a 2-dominating set of  $T'$  and hence  $\gamma_2(T') \leq \gamma_2(T) - |L_v|$ . Now if  $p \geq 2$ , that is  $v$  is adjacent to at least two leaves, then every  $\gamma_2(T')$ -set can be extended to a 2-dominating set of  $T$  by adding  $L_v$ . So  $\gamma_2(T) \leq \gamma_2(T') + |L_v|$ , which leads the equality. Assume now that  $p = 1$  and  $w$  is a leaf of  $T'$ . By Observation 1,  $w$  is in every  $\gamma_2(T')$ -set  $D'$ , so  $D'$  is extended to a 2-dominating set of  $T$  by adding the leaf neighbor of  $v$ . Therefore,  $\gamma_2(T) \leq \gamma_2(T') + 1$ , implying the equality  $\gamma_2(T') = \gamma_2(T) - 1$ .

2) Obvious. □

In [7], Haynes *et al.* showed that the 2-domination number is bounded from below by the total domination number for every nontrivial tree.

**Theorem 3 (Haynes *et al.* [7]).** *For every nontrivial tree,  $\gamma_2(T) \geq \gamma_t(T)$ .*

Below, we extend this result onto cactus graphs. The total domination and 2-domination numbers of a cycle are easy to compute.

**Observation 4.** For a cycle  $C_n$  on  $n \geq 3$  vertices:


- i)  $\gamma_t(C_n) = n/2$  if  $n \equiv 0 \pmod{4}$  and  $\gamma_t(C_n) = \lfloor n/2 \rfloor + 1$  otherwise,
- ii)  $\gamma_2(C_n) = \lceil n/2 \rceil$ .

**Theorem 5.** If  $G$  is a nontrivial connected cactus graph with  $k(G)$  even cycles ( $k(G) \geq 0$ ), then  $\gamma_2(G) \geq \gamma_t(G) - k(G)$ , and this bound is sharp.

*Proof.* If  $G$  is a tree, then  $k(G) = 0$  and by Theorem 3 the result is valid. If  $G$  is a cycle  $C_n$ , then by Observation 4, the result holds. Thus we assume that  $G$  is neither a tree nor a cycle  $C_n$ . Among all connected cactus graphs with  $k(G)$  even cycles that do not satisfy the result, let  $G$  be one which contains as few vertices and edges as possible. Let  $S$  be a  $\gamma_2(G)$ -set. Assume first that there are two adjacent vertices  $x, y$  on some cycle such that  $x, y$  are both in  $S$  or both not in  $S$ . Consider the spanning graph  $G'$  obtained by removing the edge  $xy$ . Then  $S$  is a 2-dominating set of  $G'$  and hence  $\gamma_2(G) = |S| \geq \gamma_2(G')$ . Also  $\gamma_t(G') \geq \gamma_t(G)$ , since every total dominating set of  $G'$  is a total dominating set of  $G$ . Now  $G'$  satisfies the result and so  $\gamma_2(G') \geq \gamma_t(G') - k(G')$ . Since  $k(G) \geq k(G')$ , it follows that  $\gamma_2(G) \geq \gamma_2(G') \geq \gamma_t(G') - k(G') \geq \gamma_t(G) - k(G)$ , a contradiction.

Thus we assume that all vertices on the cycles are contained alternately in  $S$ . This implies that  $G$  contains no odd cycle. Let  $u, v$  be two adjacent vertices on an even cycle such that  $u \in S$  and  $v \notin S$ . Let  $G''$  be the spanning graph of  $G$  obtained by removing the edge  $uv$ . Then  $S \cup \{v\}$  is a 2-dominating set of  $G''$  and so  $\gamma_2(G'') \leq |S| + 1$ . There also is  $\gamma_t(G'') \geq \gamma_t(G)$  and  $k(G) = k(G'') + 1$ . Now since  $G''$  satisfies  $\gamma_2(G'') \geq \gamma_t(G'') - k(G'')$ , we obtain  $\gamma_2(G) + 1 \geq \gamma_2(G'') \geq \gamma_t(G'') - k(G'') \geq \gamma_t(G) - k(G) + 1$ . Therefore,  $\gamma_2(G) \geq \gamma_t(G) - k(G)$ , a contradiction.

That this bound is sharp may be seen by considering the cactus graph  $G_q$  ( $q \geq 1$ ) formed from  $q$  path  $P_5$ , each one with the center vertex  $v_i$  where  $1 \leq i \leq q$  and  $q$  cycle  $C_6$  by adding edges between all center vertices so that the subgraph induced by the center vertices is a path  $P_q$ . Then we identify a vertex of a cycle  $C_6$  with one leaf of each path  $P_5$ . See Figure 1 for an example of  $G_3$ . For  $G_q$ ,  $\gamma_2(G_q) = 5q$ ,  $\gamma_t(G_q) = 6q$  and  $k(G_q) = q$ . □



**Fig. 1.** The graph  $G_3$

In [5], Fink and Jacobson have established a lower bound on the 2-domination number for every tree in term of its order.

**Theorem 6** ([5]). *If  $T$  is a tree of order  $n$ , then  $\gamma_2(T) \geq (n + 1)/2$ .*

Next we give a lower bound for the 2-domination number in trees that improves Fink and Jacobson's one if  $\ell > s$ .

**Lemma 7.** *If  $T$  is a tree of order  $n$  with  $\ell$  leaves and  $s$  support vertices, then  $\gamma_2(T) \geq (n + \ell - s)/2$ , and this bound is sharp.*

*Proof.* We proceed by induction on the order of  $T$ . If  $\text{diam}(T) \in \{0, 1\}$ , then the result is valid. If  $\text{diam}(T) = 2$ , then  $T$  is a star  $K_{1,p}$  ( $p \geq 2$ ), where  $\gamma_2(T) = p$  and  $(n + \ell - s)/2 = p$ , so the result is valid. If  $\text{diam}(T) = 3$ , then  $T$  is a vdouble star  $S_{p,q}$ , where  $\gamma_2(T) = p + q$  if  $\min\{p, q\} \geq 2$  and  $\gamma_2(T) = 2 + \max\{p, q\}$  otherwise. Thus again the result is valid. Assume that for every tree  $T'$  of order  $n'$  with  $n > n'$ , there is  $\gamma_2(T') \geq (n' + \ell' - s')/2$ .

Let  $T$  be a tree of order  $n$ . Root  $T$  at a vertex  $r$  of maximum eccentricity  $\text{diam}(T) \geq 4$ . Let  $v$  be a support vertex of maximum distance from  $r$  and  $u$  the parent of  $v$  in the rooted tree.

Let  $T' = T - (L_v \cup \{v\})$ . Then  $n' = n - (|L_v| + 1)$  and  $T'$  is nontrivial. We consider two cases.

**Case 1.**  $\deg_T(v) \geq 3$ . By Observation 2, since  $|L_v| \geq 2$ , then  $\gamma_2(T) = \gamma_2(T') + |L_v|$ . If  $u$  is not a leaf in  $T'$ , then  $\ell' = \ell - |L_v|$  and  $s' = s - 1$ . Applying the inductive hypothesis to  $T'$ ,

$$\gamma_2(T) - |L_v| = \gamma_2(T') \geq (n' + \ell' - s')/2 = (n + \ell - s)/2 - |L_v|,$$

hence  $\gamma_2(T) \geq (n + \ell - s)/2$ .

If  $u$  is a leaf in  $T'$ , then  $\ell' = \ell - |L_v| + 1$  and  $s' \leq s$ . Applying the inductive hypothesis to  $T'$ ,

$$\gamma_2(T) - |L_v| = \gamma_2(T') \geq (n' + \ell' - s')/2 \geq (n + \ell - s)/2 - |L_v|,$$

hence  $\gamma_2(T) \geq (n + \ell - s)/2$ .

**Case 2.**  $\deg_T(v) = 2$ , that is  $|L_v| = 1$ . If  $u$  is not a leaf in  $T'$ , then  $\ell' = \ell - 1$  and  $s' = s - 1$ . Again by Observation 2,  $\gamma_2(T) - 1 \geq \gamma_2(T')$ . Applying the inductive hypothesis to  $T'$ , we obtain the desired result. Now if  $u$  is a leaf in  $T'$ , then by Observation 2,  $\gamma_2(T) - 1 = \gamma_2(T')$ . Also  $\ell' = \ell$  and  $s' \leq s$ . Applying the inductive hypothesis to  $T'$ , the result follows.

That this bound is sharp may be seen in a tree  $T$  where every vertex  $T$  is either a leaf or a support vertex adjacent to at least two leaves. Clearly,  $n = \ell + s$  and  $\gamma_2(T) = \ell = (n + \ell - s)/2$ .  $\square$

Notice that in [1], Blidia *et al.* showed that every nontrivial tree  $T$  satisfies  $\gamma_2(T) \leq (n + \ell)/2$ . So Lemma 7 gives in some sense a best framing for the 2-domination number in trees.

**Theorem 8.** *If  $G$  is a graph of order  $n$  with at most one cycle,  $\ell$  leaves and  $s$  support vertices, then  $\gamma_2(G) \geq (n + \ell - s)/2$ , and this bound is sharp.*

*Proof.* If all the components of  $G$  are trees, then by Lemma 7 the result holds. If  $G$  is a cycle  $C_n$  then  $\ell = s = 0$  and by Observation 4,  $\gamma_2(C_n) = \lceil n/2 \rceil$ , implying that the result is valid. Thus  $G$  contains a component  $H$  that is a unicycle graph with a cycle  $C$  where at least one vertex of  $C$  has degree at least three. It suffices to prove the theorem for the subgraph  $H$ . Let  $S$  be a  $\gamma_2(H)$ -set and assume that  $H$  is the smallest connected unicycle graph that does not satisfy the theorem.

Suppose that  $H$  contains a support vertex, say  $v \notin C$ . We further assume that  $v$  is at maximum distance from  $C$ . Then  $L_v \subset S$  and without loss of generality  $v \notin S$  (else replace  $v$  by its neighbor, say  $w$ , in the unique path from  $v$  to  $C$ ). Let  $H' = H - (L_v \cup \{v\})$ . Then  $H'$  is a connected unicycle graph with  $n(H') = n(H) - (|L_v| + 1)$  and  $S - L_v$  is a 2-dominating set of  $H'$ . Hence  $\gamma_2(H) - |L_v| \geq \gamma_2(H')$  and since  $H'$  is smaller than  $H$ , it satisfies the theorem. If  $\deg_H(w) \geq 3$  then  $\ell(H') = \ell(H) - |L_v|$  and  $s(H') = s(H) - 1$ .

It follows that

$$\begin{aligned} \gamma_2(H) - |L_v| &\geq \gamma_2(H') \geq (n(H') + \ell(H') - s(H'))/2 = \\ &= (n(H) - (|L_v| + 1) + \ell(H) - |L_v| - s(H) + 1) / 2 \end{aligned}$$

and, therefore,  $\gamma_2(H) \geq (n(H) + \ell(H) - s(H))/2$ , contradicting our assumption.

Now if  $\deg_H(w) = 2$ , then  $\ell(H') = \ell(H) - |L_v| + 1$  and  $s(H') \leq s(H)$ . It follows that

$$\begin{aligned} \gamma_2(H) - |L_v| &\geq \gamma_2(H') \geq (n(H') + \ell(H') - s(H'))/2 \geq \\ &\geq (n(H) - (|L_v| + 1) + \ell(H) - |L_v| + 1 - s(H))/2 \end{aligned}$$

and, therefore,  $\gamma_2(H) \geq (n(H) + \ell(H) - s(H))/2$ , a contradiction.

It remains to examine the case where every support vertex of  $H$  is on the cycle  $C$ . Let  $u$  be a support vertex on  $C$  such that  $u \in S$ . Let  $H'$  be the graph obtained from  $H$  by removing all leaves adjacent to  $u$ . Then  $S - L_u$  is a 2-dominating set of  $H'$ ,  $\ell' = \ell - |L_u|$  and  $s' = s - 1$ . Thus

$$\begin{aligned} \gamma_2(H) - |L_u| &\geq \gamma_2(H') \geq (n(H') + \ell(H') - s(H'))/2 = \\ &= (n(H) - |L_u| + \ell(H) - |L_u| - s(H) + 1) / 2 \end{aligned}$$

and, therefore,  $\gamma_2(H) > (n(H) + \ell(H) - s(H))/2$ , a contradiction.

Thus we assume that every support vertex on  $C$  is not in  $S$ . If  $C$  is a triangle, that is  $C = C_3$  then it is a simple task to check the result depending on whether  $C$  contains one, two or three support vertices. Thus we assume that the length of  $C$  is at least four. Let  $x$  be a support vertex and  $y, z$  its two neighbors on  $C$ . Let  $H'$  be the graph obtained from  $H$  by removing  $x$  and its leaves and by adding a new edge  $yz$ . Then  $S - L_x$  is a 2-dominating set of  $H'$ ,  $n(H') = n(H) - (|L_x| + 1)$ ,  $\ell(H') = \ell(H) - |L_x|$  and  $s(H') = s(H) - 1$ .

It follows that

$$\begin{aligned}\gamma_2(H) - |L_x| &\geq \gamma_2(H') \geq (n(H') + \ell(H') - s(H'))/2 = \\ &= (n(H) - |L_x| - 1 + \ell(H) - |L_x| - s(H) + 1)/2\end{aligned}$$

and, therefore,  $\gamma_2(H) \geq (n(H) + \ell(H) - s(H))/2$ , a contradiction.

The unicycle graph  $G$  formed by a cycle  $C$  where each vertex on  $C$  is adjacent to at least two leaves shows that the lower bound of Theorem 8 is attained.  $\square$

Note that the lower bound in Theorem 8 is not valid for cactus graphs with at least two cycles. To see this, consider the graph  $G_k$  formed by  $k \geq 2$  cycles  $C_4$  by identifying a vertex from each cycle into one vertex. Then  $n(G_k) = 3k + 1$ ,  $\ell = s = 0$  and  $\gamma_2(G) = k + 1 < (n(G_k) + \ell - s)/2 = (3k + 1)/2$ .

### 3. UPPER BOUND

It is shown in [1] that the 2-domination number is bounded from below by the independence number for every tree  $T$ . In this section we establish an upper bound for the 2-domination number in terms of the independence number and the number of support vertices, which gives a good framing for the 2-domination number in trees.

**Theorem 1.** *If  $T$  is a tree of order at least three with  $s$  support vertices, then  $\gamma_2(T) \leq \beta(T) + s - 1$  and this bound is sharp.*

*Proof.* We proceed by induction on the number of vertices of  $T$ . If  $\text{diam}(T) = 2$  then  $T$  is a star  $K_{1,p}$  ( $p \geq 2$ ) where  $\gamma_2(T) = \beta(T) = p$  and  $s = 1$ , so the result holds. If  $\text{diam}(T) = 3$  then  $T$  is a double star  $S_{p,q}$  with  $q \geq p$  where  $\gamma_2(T) = p + q$  if  $p \geq 2$  and  $\gamma_2(T) = q + 2$  otherwise,  $\beta(T) = p + q$  and  $s = 2$ . Thus the result is valid. Assume that for every tree  $T'$  of order  $n'$  with  $n > n' \geq 3$ , there is  $\gamma_2(T') \leq \beta(T') + s' - 1$ .

Let  $T$  be a tree of order  $n$ . Root  $T$  at a vertex  $r$  of maximum eccentricity  $\text{diam}(T) \geq 4$ . Let  $v$  be a support vertex of maximum distance from  $r$  and  $u$  the parent of  $v$  in the rooted tree.

Let  $T' = T - (\{v\} \cup L_v)$ . Since  $\text{diam}(T) \geq 4$ , the order of  $T'$  is at least three. We consider two cases.

**Case 1.**  $\deg_T(v) \geq 3$ . By Observation 2,  $\gamma_2(T) - |L_v| = \gamma_2(T')$ ,  $\beta(T) - |L_v| = \beta(T')$  and  $s' \leq s$ . Applying our induction to  $T'$ , we obtain:

$$\gamma_2(T) - |L_v| = \gamma_2(T') \leq \beta(T') + s' - 1 \leq \beta(T) - |L_v| + s - 1.$$

Hence  $\gamma_2(T) \leq \beta(T) + s - 1$ .

**Case 2.**  $\deg_T(v) = 2$ . Then  $v$  is adjacent to exactly one leaf, say  $v'$ , so  $|L_v| = 1$ . We again consider two cases.

**Case 2.1.**  $\deg_T(u) = 2$ . Then  $s' \leq s$ , and by Observation 2,  $\gamma_2(T) - 1 = \gamma_2(T')$  and  $\beta(T) - 1 = \beta(T')$ . Applying the inductive hypothesis to  $T'$ , we obtain the desired inequality.

**Case 2.2.**  $\deg_T(u) \geq 3$ . Then  $s' = s - 1$  and by Observation 2,  $\beta(T) - 1 = \beta(T')$ . Also  $\gamma_2(T) \leq \gamma_2(T') + 2$ , since every  $\gamma_2(T')$ -set can be extended to a 2-dominating set of  $T$  by adding  $\{v, v'\}$ . By induction on  $T'$

$$\gamma_2(T) \leq \gamma_2(T') + 2 \leq \beta(T') + s' + 1 = (\beta(T) - 1) + (s - 1) + 1,$$

hence  $\gamma_2(T) \leq \beta(T) + s - 1$ .

The upper bound is sharp for the path  $P_n$  of even order  $n \geq 4$ . □

In [4], Favaron proved that every tree  $T$  of order  $n$  with  $\ell$  leaves satisfies  $\beta(T) \geq (n + \ell)/3$ . Using Lemma 7 and Theorem 1, we obtain the following corollary for the independence number, which in some sense improves Favaron's one [4] for trees.

**Corollary 2.** *If  $T$  is a tree of order at least 3 with  $\ell$  leaves and  $s$  support vertices, then  $\beta(T) \geq (n + \ell - 3s + 2)/2$ .*

## REFERENCES

- [1] Blidia M., Chellali M., Favaron O., *Independence and 2-domination in trees*. Australasian J. Comb. **33** (2005), 317–327.
- [2] Blidia M., Chellali M., Volkmann L., *Bounds of the 2-domination number of graphs* Utilitas Math., to appear.
- [3] Favaron O., *On a conjecture of Fink and Jacobson concerning  $k$ -domination and  $k$ -dependence*. J. Combinat. Theory Ser. B **39** (1985), 101–102.
- [4] Favaron O. *A bound on the independent domination number of a tree*, Int. J. Graph Theory **1(1)** (1992), 19–27.
- [5] Fink J. F., Jacobson M. S.,  *$n$ -domination in graphs*, in: Alavi Y. and Schwenk A. J. (eds), Graph Theory with Applications to Algorithms and Computer Science, New York, Wiley, 1985, 283–300.
- [6] Fink J. F., Jacobson M. S., *On  $n$ -domination,  $n$ -dependence and forbidden subgraphs*. Graph Theory with Applications to Algorithms and Computer Science. New York, Wiley, 1985, 301–312.
- [7] Haynes T. W., Hedetniemi S. T., Henning M. A., Slater P. J.,  *$H$ -forming sets in graphs*, Discrete Mathematics **262** (2003), 159–169.
- [8] Haynes T. W., Hedetniemi S. T., Slater P. J., *Fundamentals of Domination in Graphs*, Marcel Dekker, New York, 1998.
- [9] Haynes T. W., Hedetniemi S. T., Slater P. J. (eds), *Domination in Graphs: Advanced Topics*, Marcel Dekker, New York, 1998.

Mustapha Chellali  
m.chellali@yahoo.com

University of Blida  
Department of Mathematics  
B.P. 270, Blida, Algeria

*Received: October 31, 2005.*