

Jarosław Mikołajski, Ewa Schmeidel

COMPARISON OF PROPERTIES OF SOLUTIONS
OF DIFFERENTIAL EQUATIONS
AND RECURRENCE EQUATIONS
WITH THE SAME CHARACTERISTIC EQUATION
(ON EXAMPLE OF THIRD ORDER LINEAR EQUATIONS
WITH CONSTANT COEFFICIENTS)

Abstract. Third order linear homogeneous differential and recurrence equations with constant coefficients are considered. We take the both equations with the same characteristic equation. We show that these equations (differential and recurrence) can have solutions with different properties concerning oscillation and boundedness. Especially the numbers of suitable types of solutions taken out from fundamental sets are presented. We give conditions under which the asymptotic properties considered are the same for the both equations.

Keywords: differential equation, recurrence, linear, third order, oscillatory solution, bounded solution.

Mathematics Subject Classification: 34A05, 34C10, 34C11, 39A10.

1. INTRODUCTION

Concurrently with the development of modern computational methods, discrete equations, so-called recurrences, have acquired a great importance. They often replace differential equations in mathematical modelling. For example, the logistic equation describing population increase is written in the form of a differential equation $y' = y(b - ay)$ (Verhulst–Pearl equation) or in the form of a recurrence equation $u_{n+1} = u_n(b - au_n)$ (Pielou equation) (see [4] and [13]).

It appears that application of a discrete mathematical model instead of the continuous one can lead to qualitatively different solutions, particularly with respect to oscillation or boundedness. Different quantities of solutions can be seen even in the case of linear equations with constant coefficients. Oscillation of such third order

equations is investigated in [11] and boundedness in [10]. In this paper we present the numbers of suitable types of solutions taken out from fundamental sets. We also give conditions under which the asymptotic properties considered are the same for two equations: differential and recurrence ones.

We consider a third order linear homogeneous equation with constant coefficients written in the continuous and discrete form, namely the differential equation

$$\frac{d^3 f}{dt^3} + a \frac{d^2 f}{dt^2} + b \frac{df}{dt} + cf = 0 \quad (1)$$

and the recurrence equation

$$u_{n+3} + au_{n+2} + bu_{n+1} + cu_n = 0, \quad (2)$$

where $f : R \rightarrow R$, $u : N \rightarrow R$, R denotes the set of real numbers, N the set of positive integers and a, b, c are real constants.

A function $f : R \rightarrow R$ (a sequence $u : N \rightarrow R$) is called trivial if there exists a $t_0 \in R$ (an $n_0 \in N$) such that $f(t) = 0$ for every $t > t_0$ ($u_n = 0$ for every $n > n_0$). Otherwise, a function f (a sequence u) is said to be nontrivial. Every nontrivial real function f (sequence u) satisfying equation (1) ((2)) is called a solution of this equation.

A solution of equation (1) ((2)) is called nonoscillatory if there exists a $t_0 \in R$ (an $n_0 \in N$) such that it is a positive function (sequence) on the set $[t_0, \infty)$ ($\{n \in N : n \geq n_0\}$) or a negative function (sequence) on this set. Otherwise, a solution is said to be oscillatory.

A solution of equation (1) ((2)) is called bounded if it is a bounded function (sequence) on the set $[0, \infty)$ (N). Otherwise a solution is said to be unbounded.

Equation (1) or (2) is called oscillatory (nonoscillatory, bounded, unbounded) if each of its solutions is oscillatory (nonoscillatory, bounded, unbounded).

Fundamentals of differential equations theory can be found in many monographs, e.g. Stiepanow [19]. The background of recurrences theory is given by Agarwal [1], Elaydi [4], Kelley and Peterson [6] and, in Polish, by Koźniewska [8]. Oscillation of solutions of third order difference equations was considered for example by Kobza [7], Migda, Schmeidel and Drozdowicz [9], Popenda and Schmeidel [14], Saker [15], Smith [17], Thandapani and Mahalingam [20] and boundedness by Andruch-Sobiło and Migda [2], Dośla and Kobza [3], Graef and Thandapani [5], Smith [16], Smith and Taylor [18].

Putting $f(t) = \exp(rt)$ in equation (1) or $u_n = r^n$ in equation (2), the same characteristic equation

$$r^3 + ar^2 + br + c = 0 \quad (3)$$

is obtained in the continuous as well as in the discrete case. Solutions of equation (3) are described in [12].

Let us denote

$$q = c - \frac{1}{3}ab + \frac{2}{27}a^3, \quad (4)$$

$$\Delta = c^2 + \frac{4}{27}b^3 - \frac{2}{3}abc - \frac{1}{27}a^2b^2 + \frac{4}{27}a^3c. \tag{5}$$

We will use the following theorem, which can be found in [11].

Theorem 1.

1° If $\Delta > 0$, then (3) has one real root r_1 and two complex conjugate roots r_2, r_3 with non-vanishing imaginary parts:

$$\begin{aligned} r_1 &= \sqrt[3]{\frac{-q-\sqrt{\Delta}}{2}} + \sqrt[3]{\frac{-q+\sqrt{\Delta}}{2}} - \frac{a}{3}, \\ r_2 &= -\frac{1}{2} \left(\sqrt[3]{\frac{-q-\sqrt{\Delta}}{2}} + \sqrt[3]{\frac{-q+\sqrt{\Delta}}{2}} \right) - \frac{a}{3} + i\frac{\sqrt{3}}{2} \left(\sqrt[3]{\frac{-q-\sqrt{\Delta}}{2}} - \sqrt[3]{\frac{-q+\sqrt{\Delta}}{2}} \right), \\ r_3 &= -\frac{1}{2} \left(\sqrt[3]{\frac{-q-\sqrt{\Delta}}{2}} + \sqrt[3]{\frac{-q+\sqrt{\Delta}}{2}} \right) - \frac{a}{3} - i\frac{\sqrt{3}}{2} \left(\sqrt[3]{\frac{-q-\sqrt{\Delta}}{2}} - \sqrt[3]{\frac{-q+\sqrt{\Delta}}{2}} \right). \end{aligned} \tag{6}$$

2° If $\Delta = 0$, then (3) has three real roots given by (7), but one is multiple:

$$r_1 = -2\sqrt[3]{\frac{q}{2}} - \frac{a}{3}, \quad r_2 = r_3 = \sqrt[3]{\frac{q}{2}} - \frac{a}{3}. \tag{7}$$

3° If $\Delta < 0$, then (3) has three different real roots:

$$r_{k+1} = \sqrt[6]{16(q^2 - \Delta)} \cos \frac{\arccos \frac{-q}{\sqrt{q^2 - \Delta}} + 2k\pi}{3} - \frac{a}{3}, \quad k = 0, 1, 2. \tag{8}$$

Remark 1. The following equality

$$q^2 - \Delta = \frac{4}{27} \left(\frac{1}{27}a^6 - \frac{1}{3}a^4b + a^2b^2 - b^3 \right)$$

holds.

Remark 2. If $\Delta = 0$ and $q \neq 0$, then the roots of (3) can also be written in form (8).

Remark 3. The case $\Delta = 0$ and $q = 0$ holds only for $r_1 = r_2 = r_3$.

2. PARTITION OF SOLUTIONS IN FUNDAMENTAL SETS

Assume $c \neq 0$, which is required for (2) to be a third order equation. Then roots of equation (3) are not equal to zero. In the case $\Delta > 0$, the roots r_2 and r_3 can be written in the form

$$\begin{aligned} r_2 &= |r_2| (\cos \varphi + i \sin \varphi), \\ r_3 &= |r_2| (\cos \varphi - i \sin \varphi), \end{aligned}$$

Table 1

Case	Equation (1)	Equation (2)	Remarks
$\Delta > 0$	$e^{r_1 t}, e^{\operatorname{Re} r_2 t} \cos(\operatorname{Im} r_2 t), e^{\operatorname{Re} r_2 t} \sin(\operatorname{Im} r_2 t)$	$r_1^n, r_2^n \cos n\varphi, r_2^n \sin n\varphi$	r_1, r_2 defined by (6), $r_2 = r_2 (\cos \varphi + i \sin \varphi)$
$\Delta = 0, r_1 \neq r_2$	$e^{r_1 t}, e^{r_2 t}, te^{r_2 t}$	r_1^n, r_2^n, nr_2^n	r_1, r_2 defined by (7)
$\Delta = 0, r_1 = r_2$	$e^{r_1 t}, te^{r_1 t}, t^2 e^{r_1 t}$	$r_1^n, nr_1^n, n^2 r_1^n$	r_1, r_2 defined by (7)
$\Delta < 0$	$e^{r_1 t}, e^{r_2 t}, e^{r_3 t}$	r_1^n, r_2^n, r_3^n	r_1, r_2, r_3 defined by (8)

where φ is an argument of r_2 . We will consider fundamental sets of solutions indicated in Table 1.

Let $\Delta < 0$. Then there exist $k, l, m \in \{1, 2, 3\}$ such that $r_k < r_l < r_m$. Set $R_1 = r_k, R_2 = r_l, R_3 = r_m$.

Let $\Delta = 0$. Set $R_1 = \min\{r_1, r_2\}, R_2 = r_2, R_3 = \max\{r_1, r_2\}$. Thus $R_1 \leq R_2 \leq R_3$.

The numbers of oscillatory solutions taken out from the fundamental sets (Tab. 1) are presented in Table 2 and bounded solutions in Table 3 (see the attached interleaf).

Table 2

Case	Equation (1)	Equation (2)
$\Delta > 0, r_1 < 0$	2	3
$\Delta > 0, r_1 > 0$	2	2
$\Delta \leq 0, R_1, R_2, R_3 < 0$	0	3
$\Delta \leq 0, R_1, R_2 < 0, R_3 > 0$	0	2
$\Delta \leq 0, R_1 < 0, R_2, R_3 > 0$	0	1
$\Delta \leq 0, R_1, R_2, R_3 > 0$	0	0

3. MAIN RESULTS

Remark 4. Equation (1) is never oscillatory. Thus this equation can be nonoscillatory or it can have oscillatory and nonoscillatory solutions. However, equation (2) is oscillatory or nonoscillatory or it can have both oscillatory and nonoscillatory solutions.

Let us

$$d = \sqrt[3]{\frac{-q - \sqrt{\Delta}}{2}} + \sqrt[3]{\frac{-q + \sqrt{\Delta}}{2}} \quad \text{for } \Delta \geq 0,$$

$$D = \sqrt{d^2 + \frac{1}{3}ad + \frac{1}{9}a^2 - \frac{3}{2}\sqrt[3]{2(q^2 - \Delta)}},$$

$$p_k = \cos \frac{\arccos \frac{-q}{\sqrt{q^2 - \Delta}} + 2k\pi}{3} \quad \text{for } \Delta < 0 \quad (k = 0, 1, 2).$$

Theorem 2. Equations (1) and (2) are nonoscillatory if and only if one of the following cases holds:

1° $\Delta \leq 0, \Delta^2 + q^2 > 0, p_k > \frac{a}{3\sqrt[6]{16(q^2 - \Delta)}}$ for $k = 0, 1, 2,$

2° $\Delta = q = 0, \quad a < 0.$

Proof. In Case (1°), by Theorem 1 and Remark 2, from (8) there follows

$$r_k = \sqrt[6]{16(q^2 - \Delta)}p_{k-1} - \frac{a}{3} > 0$$

and, obviously, $R_k > 0$ for $k = 1, 2, 3.$ In Case (ii), by Remark 3 and (7) we obtain $r_1 = r_2 = r_3 > 0.$ By Table 2, the proof is completed. □

Theorem 3. Equations (1) and (2) are bounded if and only if one of the following cases holds:

1° $\Delta > 0, \max(-\frac{2}{3}a, \frac{1}{3}a - 1) \leq d < \frac{1}{3}a, \quad D \leq 1,$

2° $\Delta \leq 0, \Delta^2 + q^2 > 0, \frac{a-3}{3\sqrt[6]{16(q^2 - \Delta)}} \leq p_k < \frac{a}{3\sqrt[6]{16(q^2 - \Delta)}}$ for $k = 0, 1, 2$ (the equality $p_k = \frac{a-3}{3\sqrt[6]{16(q^2 - \Delta)}}$ is valid for at most one k),

3° $\Delta = q = 0, \quad 0 < a < 3.$

Proof. In Case (1°), by Theorem 1, from (6) there follows $r_1 = d - \frac{1}{3}a < 0$ and $r_1 \geq -1.$ Moreover,

$$\text{Re } r_2 = -\frac{1}{2}d - \frac{1}{3}a \leq -\frac{1}{2}(-\frac{2}{3}a) - \frac{1}{3}a = 0$$

and

$$|r_2| = \sqrt{\frac{1}{4}d^2 + \frac{1}{3}ad + \frac{1}{9}a^2 + \frac{3}{4} \left(d^2 - 4\sqrt[3]{\frac{q^2 - \Delta}{4}} \right)} =$$

$$= \sqrt{d^2 + \frac{1}{3}ad + \frac{1}{9}a^2 - 3\sqrt[3]{\frac{q^2 - \Delta}{4}}} = D \leq 1.$$

In Case (2°), by Theorem 1 and Remark 2, from (8) we obtain $-1 \leq r_k < 0$ for $k = 1, 2, 3$ and $r_k = -1$ is valid for at most one $k.$ The same inequalities hold for R_k ($k = 1, 2, 3$). In Case (3°), by Remark 3 and (7), there is $-1 < r_1 = r_2 = r_3 < 0.$ By Table 3, the proof is completed. □

Theorem 4. *Equations (1) and (2) are unbounded if and only if one of the following cases holds:*

$$1^\circ \Delta > 0, \frac{1}{3}a + 1 < d < -\frac{2}{3}a, D > 1,$$

$$2^\circ \Delta \leq 0, \Delta^2 + q^2 > 0, p_k > \frac{a+3}{\sqrt[3]{16(q^2-\Delta)}} \text{ for } k = 0, 1, 2,$$

$$3^\circ \Delta = q = 0, a < -3.$$

Proof. Analogously as in proof of Theorem 3, we obtain:

in Case (1°): $r_1 > 1, \operatorname{Re} r_2 > 0, |r_2| > 1$;

in Case (2°): $R_1, R_2, R_3 > 1$ for $k = 1, 2, 3$;

in Case (3°): $r_1 = r_2 = r_3 > 1$.

By Table 3, this completes the proof. \square

Remark 5. *Case (1°) of Theorem 3 can only hold for $a > 0$ and Case (1°) of Theorem 4 only for $a < -1$.*

Indeed,

$$\text{if } -\frac{2}{3}a < \frac{1}{3}a, \text{ then } a > 0;$$

$$\text{if } \frac{1}{3}a + 1 < -\frac{2}{3}a, \text{ then } a < -1.$$

REFERENCES

- [1] Agarwal R. P., *Difference Equations and Inequalities. Theory, Methods and Applications*, Marcel Dekker, New York, 2000 (2nd ed.).
- [2] Andruch-Sobiło A., Migda M., *Bounded solutions of third order nonlinear difference equations*, Rocky Mountain Journal of Mathematics 36 (2006) 1, 23–24.
- [3] Došla Z., Kobza A., *Global asymptotic properties of third order difference equations*, Computers and Mathematics with Applications 48 (2004) 1–2, 191–200.
- [4] Elaydi S. N., *An Introduction to Difference Equations*, Springer, 1996.
- [5] Graef J. R., Thandapani E., *Oscillatory and asymptotic behavior of solutions of third order delay difference equations*, Funkcialaj Ekvacioj 42 (1999) 3, 355–369.
- [6] Kelley W. G., Peterson A. C., *Difference Equations, an Introduction with Applications*, Academic Press, 1991.
- [7] Kobza A., *Property A for third order difference equations*, Studies of the University of Zilina, Mathematical Series 17 (2003), 109–14.
- [8] Koźniewska I., *Równania rekurencyjne*, PWN, Warszawa, 1972 (in Polish).
- [9] Migda M., Schmeidel E., Drozdowicz A., *Nonoscillation results for some third order nonlinear difference equation*, Folia Facultatis Scientiarum Naturalium Universitatis Masarykianae Brunensis Mathematica 13 (2003), 185–192.

- [10] Mikołajski J., Schmeidel E., *Comparison of boundedness of solutions of third order linear differential and recurrence equations with constant coefficients* (submitted).
- [11] Mikołajski J., Schmeidel J., *Porównanie rozwiązań równań różniczkowych i równań rekurencyjnych o tym samym równaniu charakterystycznym* (in Polish, submitted).
- [12] Mostowski A., Stark A., *Elementy algebry wyższej*, PWN, Warszawa, 1968 (in Polish).
- [13] Palczewski A., *Równania różniczkowe zwyczajne*, WNT, Warszawa, 1999 (in Polish).
- [14] Popena J., Schmeidel E., *Nonoscillatory solutions of third order difference equations*, *Portugaliae Mathematica* 49 (1992), 233–239.
- [15] Saker S.H., *Oscillation of third-order difference equations*, *Portugaliae Mathematica* 61 (2004) 3, 249–257.
- [16] Smith B., *Oscillatory and asymptotic behavior in certain third order difference equations*, *Rocky Mountain Journal of Mathematics* 17 (1987) 3, 597–606.
- [17] Smith B., *Oscillation and no oscillation theorems for third order quasi-adjoint difference equations*, *Portugaliae Mathematica* 45 (1988) 3, 229–243.
- [18] Smith B., Taylor W.E., *Oscillatory and asymptotic behavior of certain fourth order difference equations*, *Rocky Mountain Journal of Mathematics* 16 (1986) 2, 403–406.
- [19] Stiepanow W.W., *Równania różniczkowe*, PWN, Warszawa, 1964 (in Polish).
- [20] Thandapani E., Mahalingam K., *Oscillatory properties of third order neutral delay difference equations*, *Demonstratio Mathematica* 35 (2002), 325–337.

Jarosław Mikołajski

jmikolaj@math.put.poznan.pl

Poznań University of Technology

Institute of Mathematics, Faculty of Electrical Engineering

ul. Piotrowo 3a, 60-965 Poznań, Poland

Ewa Schmeidel

ewa.schmeidel@put.poznan.pl

Poznań University of Technology

Institute of Mathematics, Faculty of Electrical Engineering

ul. Piotrowo 3a, 60-965 Poznań, Poland

Received: August 29, 2005.