

BOUNDARY VALUE PROBLEMS FOR n -TH ORDER DIFFERENTIAL INCLUSIONS WITH FOUR-POINT INTEGRAL BOUNDARY CONDITIONS

Bashir Ahmad and Sotiris K. Ntouyas

Abstract. In this paper, we discuss the existence of solutions for a four-point integral boundary value problem of n -th order differential inclusions involving convex and non-convex multivalued maps. The existence results are obtained by applying the nonlinear alternative of Leray Schauder type and some suitable theorems of fixed point theory.

Keywords: differential inclusions, four-point integral boundary conditions, existence, non-linear alternative of Leray Schauder type, fixed point theorems.

Mathematics Subject Classification: 34A60, 34B10, 34B15.

1. INTRODUCTION

In this paper, we consider the following n -th order differential inclusion with four-point integral boundary conditions

$$\begin{cases} x^{(n)}(t) \in F(t, x(t)), & 0 < t < 1, \\ x(0) = \alpha \int_{\xi}^{\xi} x(s) ds, & x'(0) = 0, \quad x''(0) = 0, \dots, x^{(n-2)}(0) = 0, \\ x(1) = \beta \int_{\eta}^{\eta} x(s) ds, & 0 < \xi < \eta < 1, \end{cases} \quad (1.1)$$

where $F : [0, T] \times \mathbb{R} \rightarrow \mathcal{P}(\mathbb{R})$ is a multivalued map, $\mathcal{P}(\mathbb{R})$ is the family of all subsets of \mathbb{R} and $\alpha, \beta \in \mathbb{R}$.

Multi-point boundary conditions arise in a variety of problems of applied mathematics and physics. Nonlocal multi-point problems constitute an important class of boundary value problems and have been addressed by many authors, for instance, see [1, 5–7, 10, 17, 19–21, 23, 26, 27, 30, 31].

Integral boundary conditions have various applications in applied fields such as blood flow problems, chemical engineering, thermoelasticity, underground water flow,

population dynamics, etc. For a detailed description of the integral boundary conditions, we refer the reader to the papers [2, 11] and references therein.

Differential inclusions arise in the mathematical modelling of certain problems in economics, optimal control, stochastic analysis, etc. and are widely studied by many authors, see [3, 4, 8, 9, 14, 28, 29] and the references therein.

The aim of our paper is to establish some existence results for the problem (1.1), when the right hand side is convex as well as nonconvex valued. The first result relies on the nonlinear alternative of Leray-Schauder type. In the second result, we shall combine the nonlinear alternative of Leray-Schauder type for single-valued maps with a selection theorem due to Bressan and Colombo for lower semicontinuous multivalued maps with nonempty closed and decomposable values, while in the third result, we shall use the fixed point theorem for contraction multivalued maps due to Covitz and Nadler. The methods used are standard, however their exposition in the framework of problem (1.1) is new.

The paper is organized as follows: in Section 2 we recall some preliminary facts that we need in the sequel and in Section 3 we prove our main results.

2. PRELIMINARIES

Let us recall some basic definitions on multi-valued maps [16, 22].

For a normed space $(X, \|\cdot\|)$, let $P_{cl}(X) = \{Y \in \mathcal{P}(X) : Y \text{ is closed}\}$, $P_b(X) = \{Y \in \mathcal{P}(X) : Y \text{ is bounded}\}$, $P_{cp}(X) = \{Y \in \mathcal{P}(X) : Y \text{ is compact}\}$, and $P_{cp,c}(X) = \{Y \in \mathcal{P}(X) : Y \text{ is compact and convex}\}$. A multi-valued map $G : X \rightarrow \mathcal{P}(X)$ is convex (closed) valued if $G(x)$ is convex (closed) for all $x \in X$. The map G is bounded on bounded sets if $G(\mathbb{B}) = \bigcup_{x \in \mathbb{B}} G(x)$ is bounded in X for all $\mathbb{B} \in P_b(X)$ (i.e., $\sup_{x \in \mathbb{B}} \{\sup\{|y| : y \in G(x)\}\} < \infty$). G is called upper semi-continuous (u.s.c.) on X if for each $x_0 \in X$, the set $G(x_0)$ is a nonempty closed subset of X , and if for each open set N of X containing $G(x_0)$, there exists an open neighborhood \mathcal{N}_0 of x_0 such that $G(\mathcal{N}_0) \subseteq N$. G is said to be completely continuous if $G(\mathbb{B})$ is relatively compact for every $\mathbb{B} \in P_b(X)$. If the multi-valued map G is completely continuous with nonempty compact values, then G is u.s.c. if and only if G has a closed graph, i.e., $x_n \rightarrow x_*$, $y_n \rightarrow y_*$, $y_n \in G(x_n)$ imply $y_* \in G(x_*)$. G has a fixed point if there is $x \in X$ such that $x \in G(x)$. The fixed point set of the multivalued operator G will be denoted by $FixG$. A multivalued map $G : [0; 1] \rightarrow P_{cl}(\mathbb{R})$ is said to be measurable if for every $y \in \mathbb{R}$, the function

$$t \longmapsto d(y, G(t)) = \inf\{|y - z| : z \in G(t)\}$$

is measurable.

Let $C([0, 1], \mathbb{R})$ denote a Banach space of continuous functions from $[0, 1]$ into \mathbb{R} with the norm $\|x\|_\infty = \sup_{t \in [0, 1]} |x(t)|$. Let $L^1([0, 1], \mathbb{R})$ be the Banach space of measurable functions $x : [0, 1] \rightarrow \mathbb{R}$ which are Lebesgue integrable and normed by $\|x\|_{L^1} = \int_0^1 |x(t)| dt$.

Definition 2.1. A multivalued map $F : [0, 1] \times \mathbb{R} \rightarrow \mathcal{P}(\mathbb{R})$ is said to be Carathéodory if:

- (i) $t \mapsto F(t, x)$ is measurable for each $x \in \mathbb{R}$,
- (ii) $x \mapsto F(t, x)$ is upper semicontinuous for almost all $t \in [0, 1]$.

Further, a Carathéodory function F is called L^1 -Carathéodory if

- (iii) for each $\alpha > 0$, there exists $\varphi_\alpha \in L^1([0, 1], \mathbb{R}^+)$ such that

$$\|F(t, x)\| = \sup\{|v| : v \in F(t, x)\} \leq \varphi_\alpha(t)$$

for all $\|x\|_\infty \leq \alpha$ and for a.e. $t \in [0, 1]$.

For each $y \in C([0, 1], \mathbb{R})$, define the set of selections of F by

$$S_{F,y} := \{v \in L^1([0, 1], \mathbb{R}) : v(t) \in F(t, y(t)) \text{ for a.e. } t \in [0, 1]\}.$$

Let X be a nonempty closed subset of a Banach space E and $G : X \rightarrow \mathcal{P}(E)$ be a multivalued operator with nonempty closed values. G is lower semi-continuous (l.s.c.) if the set $\{y \in X : G(y) \cap B \neq \emptyset\}$ is open for any open set B in E . Let A be a subset of $[0, 1] \times \mathbb{R}$. A is $\mathcal{L} \otimes \mathcal{B}$ measurable if A belongs to the σ -algebra generated by all sets of the form $\mathcal{J} \times \mathcal{D}$, where \mathcal{J} is Lebesgue measurable in $[0, 1]$ and \mathcal{D} is Borel measurable in \mathbb{R} . A subset \mathcal{A} of $L^1([0, 1], \mathbb{R})$ is decomposable if for all $x, y \in \mathcal{A}$ and measurable $\mathcal{J} \subset [0, 1] = J$, the function $x\chi_{\mathcal{J}} + y\chi_{J-\mathcal{J}} \in \mathcal{A}$, where $\chi_{\mathcal{J}}$ stands for the characteristic function of \mathcal{J} .

Definition 2.2. Let Y be a separable metric space and let $N : Y \rightarrow \mathcal{P}(L^1([0, 1], \mathbb{R}))$ be a multivalued operator. We say N has a property (BC) if N is lower semi-continuous (l.s.c.) and has nonempty closed and decomposable values.

Let $F : [0, 1] \times \mathbb{R} \rightarrow \mathcal{P}(\mathbb{R})$ be a multivalued map with nonempty compact values. Define a multivalued operator $\mathcal{F} : C([0, 1] \times \mathbb{R}) \rightarrow \mathcal{P}(L^1([0, 1], \mathbb{R}))$ associated with F as

$$\mathcal{F}(x) = \{w \in L^1([0, 1], \mathbb{R}) : w(t) \in F(t, x(t)) \text{ for a.e. } t \in [0, 1]\},$$

which is called the Nemytskii operator associated with F .

Definition 2.3. Let $F : [0, 1] \times \mathbb{R} \rightarrow \mathcal{P}(\mathbb{R})$ be a multivalued function with nonempty compact values. We say F is of lower semi-continuous type (l.s.c. type) if its associated Nemytskii operator \mathcal{F} is lower semi-continuous and has nonempty closed and decomposable values.

Let (X, d) be a metric space induced from the normed space $(X; \|\cdot\|)$. Consider $H_d : \mathcal{P}(X) \times \mathcal{P}(X) \rightarrow \mathbb{R} \cup \{\infty\}$ given by

$$H_d(A, B) = \max \left\{ \sup_{a \in A} d(a, B), \sup_{b \in B} d(A, b) \right\},$$

where $d(A, b) = \inf_{a \in A} d(a; b)$ and $d(a, B) = \inf_{b \in B} d(a; b)$. Then $(P_{b,cl}(X), H_d)$ is a metric space and $(P_{cl}(X), H_d)$ is a generalized metric space (see [24]).

Definition 2.4. A multivalued operator $N : X \rightarrow P_{cl}(X)$ is called:

(a) γ -Lipschitz if and only if there exists $\gamma > 0$ such that

$$H_d(N(x), N(y)) \leq \gamma d(x, y) \text{ for each } x, y \in X,$$

(b) a contraction if and only if it is γ -Lipschitz with $\gamma < 1$.

The following lemmas will be used in the sequel.

Lemma 2.5 ([25]). *Let X be a Banach space. Let $F : [0, T] \times \mathbb{R} \rightarrow \mathcal{P}_{cp,c}(X)$ be an L^1 -Carathéodory multivalued map and let Θ be a linear continuous mapping from $L^1([0, 1], X)$ to $C([0, 1], X)$. Then the operator*

$$\Theta \circ S_F : C([0, 1], X) \rightarrow \mathcal{P}_{cp,c}(C([0, 1], X)), \quad x \mapsto (\Theta \circ S_F)(x) = \Theta(S_{F,x})$$

is a closed graph operator in $C([0, 1], X) \times C([0, 1], X)$.

Lemma 2.6 (Nonlinear alternative for Kakutani maps [18]). *Let E be a Banach space, C a closed convex subset of E , U an open subset of C and $0 \in U$. Suppose that $F : \bar{U} \rightarrow \mathcal{P}_{cp,c}(C)$ is a upper semicontinuous compact map; where $\mathcal{P}_{cp,c}(C)$ denotes the family of nonempty, compact convex subsets of C . Then either:*

- (i) F has a fixed point in \bar{U} , or
- (ii) there is $u \in \partial U$ and $\lambda \in (0, 1)$ with $u \in \lambda F(u)$.

Lemma 2.7 ([12]). *Let Y be a separable metric space and let $N : Y \rightarrow \mathcal{P}(L^1([0, 1], \mathbb{R}))$ be a multivalued operator satisfying the property (BC). Then N has a continuous selection, that is, there exists a continuous function (single-valued) $g : Y \rightarrow L^1([0, 1], \mathbb{R})$ such that $g(x) \in N(x)$ for every $x \in Y$.*

Lemma 2.8 ([15]). *Let (X, d) be a complete metric space. If $N : X \rightarrow P_{cl}(X)$ is a contraction, then $FixN \neq \emptyset$.*

In order to define the solution of (1.1), we consider the following lemma.

Lemma 2.9. *For a given $y \in C[0, 1]$, the unique solution of the boundary value problem*

$$\begin{cases} x^{(n)}(t) = y(t), & 0 < t < 1, \\ x(0) = \alpha \int_{\xi}^{\xi} x(s) ds, \quad x'(0) = 0, \quad x''(0) = 0, \dots, \quad x^{(n-2)}(0) = 0, \\ x(1) = \beta \int_{\eta}^1 x(s) ds, & 0 < \xi < \eta < 1, \end{cases} \tag{2.1}$$

is given by

$$\begin{aligned}
 x(t) &= \int_0^t \frac{(t-s)^{n-1}}{(n-1)!} y(s) ds + \\
 &+ \frac{1}{n\Delta} \left[\alpha(n - \beta(1 - \eta^n)) \int_0^\xi \left(\int_0^s \frac{(s-m)^{n-1}}{(n-1)!} y(m) dm \right) ds + \right. \\
 &+ \alpha\beta\xi^n \int_\eta^1 \left(\int_0^s \frac{(s-m)^{n-1}}{(n-1)!} y(m) dm \right) ds - \alpha\xi^n \int_0^1 \frac{(1-s)^{n-1}}{(n-1)!} y(s) ds \left. \right] + \\
 &+ \frac{t^{n-1}}{\Delta} \left[-\alpha(1 - \beta(1 - \eta)) \int_0^\xi \left(\int_0^s \frac{(s-m)^{n-1}}{(n-1)!} y(m) dm \right) ds + \right. \\
 &+ \beta(1 - \alpha\xi) \int_\eta^1 \left(\int_0^s \frac{(s-m)^{n-1}}{(n-1)!} y(m) dm \right) ds - \\
 &\left. - (1 - \alpha\xi) \int_0^1 \frac{(1-s)^{n-1}}{(n-1)!} y(s) ds \right], \tag{2.2}
 \end{aligned}$$

where

$$\Delta = \frac{\alpha\xi^n}{n} \left(1 - \beta(1 - \eta) \right) + (1 - \alpha\xi) \left(1 - \frac{\beta(1 - \eta^n)}{n} \right) \neq 0. \tag{2.3}$$

Proof. We know that the general solution of the equation $x^{(n)}(t) = y(t)$ can be written as

$$x(t) = c_1 + c_2 t + c_3 t^2 + \dots + c_n t^{n-1} + \int_0^t \frac{(t-s)^{n-1}}{(n-1)!} y(s) ds, \tag{2.4}$$

where $c_1, c_2, \dots, c_n \in \mathbb{R}$ are arbitrary constants. Applying the boundary conditions for the problem (2.1), we find that $c_2 = 0, \dots, c_{n-1} = 0$,

$$\begin{aligned}
 c_1 &= \frac{\alpha}{\Delta} \left(1 - \frac{\beta(1 - \eta^n)}{n} \right) \int_0^\xi \left(\int_0^s \frac{(s-m)^{n-1}}{(n-1)!} y(m) dm \right) ds + \\
 &+ \frac{\alpha\xi^n}{n\Delta} \left\{ \beta \int_\eta^1 \left(\int_0^s \frac{(s-m)^{n-1}}{(n-1)!} y(m) dm \right) ds - \int_0^1 \frac{(1-s)^{n-1}}{(n-1)!} y(s) ds \right\}
 \end{aligned}$$

and

$$\begin{aligned}
 c_n &= -\frac{\alpha}{\Delta} \left(1 - \beta(1 - \eta)\right) \int_0^\xi \left(\int_0^s \frac{(s - m)^{n-1}}{(n - 1)!} y(m) dm \right) ds + \\
 &\quad + \frac{(1 - \alpha\xi)}{\Delta} \left\{ \beta \int_\eta^1 \left(\int_0^s \frac{(s - m)^{n-1}}{(n - 1)!} y(m) dm \right) ds - \int_0^1 \frac{(1 - s)^{n-1}}{(n - 1)!} y(s) ds \right\},
 \end{aligned}$$

where Δ is given by (2.3). Substituting the values of c_1, \dots, c_n in (2.4), we get (2.2). □

Definition 2.10. A function $x \in C^n([0, 1], \mathbb{R})$ is a solution of the problem (1.1) if there exists a function $f \in L^1([0, 1], \mathbb{R})$ such that $f(t) \in F(t, x(t))$ a.e. on $[0, 1]$ and

$$\begin{aligned}
 x(t) &= \int_0^t \frac{(t - s)^{n-1}}{(n - 1)!} f(s) ds + \\
 &\quad + \frac{1}{n\Delta} \left[\alpha \left(n - \beta(1 - \eta^n) \right) \int_0^\xi \left(\int_0^s \frac{(s - m)^{n-1}}{(n - 1)!} f(m) dm \right) ds + \right. \\
 &\quad + \alpha\beta\xi^n \int_\eta^1 \left(\int_0^s \frac{(s - m)^{n-1}}{(n - 1)!} f(m) dm \right) ds - \alpha\xi^n \int_0^1 \frac{(1 - s)^{n-1}}{(n - 1)!} f(s) ds \left. \right] + \\
 &\quad + \frac{t^{n-1}}{\Delta} \left[-\alpha \left(1 - \beta(1 - \eta) \right) \int_0^\xi \left(\int_0^s \frac{(s - m)^{n-1}}{(n - 1)!} f(m) dm \right) ds + \right. \\
 &\quad + \beta(1 - \alpha\xi) \int_\eta^1 \left(\int_0^s \frac{(s - m)^{n-1}}{(n - 1)!} f(m) dm \right) ds + \\
 &\quad \left. - (1 - \alpha\xi) \int_0^1 \frac{(1 - s)^{n-1}}{(n - 1)!} f(s) ds \right].
 \end{aligned}$$

3. MAIN RESULTS

Theorem 3.1. *Assume that:*

- (H₁) $F : [0, 1] \times \mathbb{R} \rightarrow \mathcal{P}(\mathbb{R})$ is Carathéodory and has convex values,
- (H₂) there exists a continuous nondecreasing function $\psi : [0, \infty) \rightarrow (0, \infty)$ and a function $p \in L^1([0, 1], \mathbb{R}^+)$ such that

$$\|F(t, x)\|_{\mathcal{P}} := \sup\{|y| : y \in F(t, x)\} \leq p(t)\psi(\|x\|_\infty) \quad \text{for each } (t, x) \in [0, 1] \times \mathbb{R},$$

(H₃) there exists a number $M > 0$ such that

$$\frac{M}{\frac{1}{(n-1)!} \left(1 + \frac{\delta_1 + \delta_2}{|\Delta|}\right) \psi(\|x\|_\infty) \|p\|_{L^1}} > 1, \tag{3.1}$$

where

$$\delta_1 = \frac{|\alpha|\xi^n}{n} \left(|n - \beta(1 - \eta^n)| + |\beta| + 1\right), \tag{3.2}$$

and

$$\delta_2 = |\alpha(1 - \beta((1 - \eta)))|\xi^n + (|\beta| + 1)|1 - \alpha\xi|. \tag{3.3}$$

Then the boundary value problem (1.1) has at least one solution on $[0, 1]$.

Proof. Define an operator $\Omega : C([0, 1], \mathbb{R}) \rightarrow \mathcal{P}(C([0, 1], \mathbb{R}))$ by

$$\Omega(x) = \left\{ h \in C([0, 1], \mathbb{R}) : \begin{aligned} h(t) = & \int_0^t \frac{(t-s)^{n-1}}{(n-1)!} f(s) ds + \\ & + \frac{1}{n\Delta} \left[\alpha(n - \beta(1 - \eta^n)) \int_0^\xi \left(\int_0^s \frac{(s-m)^{n-1}}{(n-1)!} f(m) dm \right) ds + \right. \\ & + \alpha\beta\xi^n \int_\eta^1 \left(\int_0^s \frac{(s-m)^{n-1}}{(n-1)!} f(m) dm \right) ds - \\ & - \alpha\xi^n \int_0^1 \frac{(1-s)^{n-1}}{(n-1)!} f(s) ds \left. \right] + \\ & + \frac{t^{n-1}}{\Delta} \left[-\alpha(1 - \beta(1 - \eta)) \int_0^\xi \left(\int_0^s \frac{(s-m)^{n-1}}{(n-1)!} f(m) dm \right) ds + \right. \\ & + \beta(1 - \alpha\xi) \int_\eta^1 \left(\int_0^s \frac{(s-m)^{n-1}}{(n-1)!} f(m) dm \right) ds - \\ & \left. - (1 - \alpha\xi) \int_0^1 \frac{(1-s)^{n-1}}{(n-1)!} f(s) ds \right] \end{aligned} \right\}$$

for $f \in S_{F,x}$. We will show that Ω satisfies the assumptions of the nonlinear alternative of Leray-Schauder type. The proof consists of several steps. As a first step, we show that Ω is convex for each $x \in C([0, 1], \mathbb{R})$. For that, let $h_1, h_2 \in \Omega$. Then there exists $f_1, f_2 \in S_{F,x}$ such that for each $t \in [0, 1]$, we have

$$\begin{aligned}
h_i(t) &= \int_0^t \frac{(t-s)^{n-1}}{(n-1)!} f_i(s) ds + \\
&+ \frac{1}{n\Delta} \left[\alpha(n - \beta(1 - \eta^n)) \int_0^\xi \left(\int_0^s \frac{(s-m)^{n-1}}{(n-1)!} f_i(m) dm \right) ds + \right. \\
&+ \alpha\beta\xi^n \int_\eta^1 \left(\int_0^s \frac{(s-m)^{n-1}}{(n-1)!} f_i(m) dm \right) ds - \alpha\xi^n \int_0^1 \frac{(1-s)^{n-1}}{(n-1)!} f_i(s) ds \left. \right] + \\
&+ \frac{t^{n-1}}{\Delta} \left[-\alpha(1 - \beta(1 - \eta)) \int_0^\xi \left(\int_0^s \frac{(s-m)^{n-1}}{(n-1)!} f_i(m) dm \right) ds + \right. \\
&+ \beta(1 - \alpha\xi) \int_\eta^1 \left(\int_0^s \frac{(s-m)^{n-1}}{(n-1)!} f_i(m) dm \right) ds - \\
&\left. - (1 - \alpha\xi) \int_0^1 \frac{(1-s)^{n-1}}{(n-1)!} f_i(s) ds \right], \quad i = 1, 2.
\end{aligned}$$

Let $0 \leq \omega \leq 1$. Then, for each $t \in [0, 1]$, we have

$$\begin{aligned}
[\omega h_1 + (1 - \omega)h_2](t) &= \int_0^t \frac{(t-s)^{n-1}}{(n-1)!} [\omega f_1(s) + (1 - \omega)f_2(s)] ds + \\
&+ \frac{1}{n\Delta} \left[\alpha(n - \beta(1 - \eta^n)) \int_0^\xi \left(\int_0^s \frac{(s-m)^{n-1}}{(n-1)!} [\omega f_1(m) + (1 - \omega)f_2(m)] dm \right) ds + \right. \\
&+ \alpha\beta\xi^n \int_\eta^1 \left(\int_0^s \frac{(s-m)^{n-1}}{(n-1)!} [\omega f_1(m) + (1 - \omega)f_2(m)] dm \right) ds - \\
&\left. - \alpha\xi^n \int_0^1 \frac{(1-s)^{n-1}}{(n-1)!} [\omega f_1(s) + (1 - \omega)f_2(s)] ds \right] + \\
&+ \frac{t^{n-1}}{\Delta} \left[-\alpha(1 - \beta(1 - \eta)) \int_0^\xi \left(\int_0^s \frac{(s-m)^{n-1}}{(n-1)!} [\omega f_1(m) + (1 - \omega)f_2(m)] dm \right) ds + \right. \\
&+ \beta(1 - \alpha\xi) \int_\eta^1 \left(\int_0^s \frac{(s-m)^{n-1}}{(n-1)!} [\omega f_1(s) + (1 - \omega)f_2(s)] dm \right) ds - \\
&\left. - (1 - \alpha\xi) \int_0^1 \frac{(1-s)^{n-1}}{(n-1)!} [\omega f_1(s) + (1 - \omega)f_2(s)] ds \right].
\end{aligned}$$

Since $S_{F,x}$ is convex (F has convex values), therefore it follows that $\omega h_1 + (1 - \omega)h_2 \in \Omega(x)$.

Next, we show that Ω maps bounded sets into bounded sets in $C([0, 1], \mathbb{R})$. For a positive number r , let $B_r = \{x \in C([0, 1], \mathbb{R}) : \|x\|_\infty \leq r\}$ be a bounded set in $C([0, 1], \mathbb{R})$. Then, for each $h \in \Omega(x)$, $x \in B_r$, there exists $f \in S_{F,x}$ such that

$$\begin{aligned} h(t) = & \int_0^t \frac{(t-s)^{n-1}}{(n-1)!} f(s) ds + \frac{1}{n\Delta} \left[\alpha (n - \beta(1 - \eta^n)) \int_0^\xi \left(\int_0^s \frac{(s-m)^{n-1}}{(n-1)!} f(m) dm \right) ds + \right. \\ & + \alpha \beta \xi^n \int_\eta^1 \left(\int_0^s \frac{(s-m)^{n-1}}{(n-1)!} f(m) dm \right) ds - \alpha \xi^n \int_0^1 \frac{(1-s)^{n-1}}{(n-1)!} f(s) ds \Big] + \\ & + \frac{t^{n-1}}{\Delta} \left[-\alpha (1 - \beta(1 - \eta)) \int_0^\xi \left(\int_0^s \frac{(s-m)^{n-1}}{(n-1)!} f(m) dm \right) ds + \right. \\ & \left. + \beta(1 - \alpha \xi) \int_\eta^1 \left(\int_0^s \frac{(s-m)^{n-1}}{(n-1)!} f(m) dm \right) ds - (1 - \alpha \xi) \int_0^1 \frac{(1-s)^{n-1}}{(n-1)!} f(s) ds \right], \end{aligned}$$

and

$$\begin{aligned} |h(t)| \leq & \left| \frac{\alpha}{n\Delta} [(n - \beta(1 - \eta^n)) - n(1 - \beta(1 - \eta))t^{n-1}] \right| \times \\ & \times \int_0^\xi \left(\int_0^s \frac{(s-m)^{n-1}}{(n-1)!} |f(m)| dm \right) ds + \\ & + \left| \frac{\beta}{n\Delta} [\alpha \xi^n + n(1 - \alpha \xi)t^{n-1}] \right| \int_\eta^1 \left(\int_0^s \frac{(s-m)^{n-1}}{(n-1)!} |f(m)| dm \right) ds + \\ & + \left| \frac{1}{n\Delta} [\alpha \xi^n + n(1 - \alpha \xi)t^{n-1}] \right| \int_0^1 \frac{(1-s)^{n-1}}{(n-1)!} |f(s)| ds + \\ & + \int_0^t \frac{(t-s)^{n-1}}{(n-1)!} |f(s)| ds \leq \\ & \leq \left\{ \left| \frac{\alpha}{n\Delta} [(n - \beta(1 - \eta^n)) - n(1 - \beta(1 - \eta))t^{n-1}] \right| \frac{\xi^n}{(n-1)!} + \right. \\ & + \left| \frac{\beta}{n\Delta} [\alpha \xi^n + n(1 - \alpha \xi)t^{n-1}] \right| \frac{1}{(n-1)!} + \\ & \left. + \left| \frac{1}{n\Delta} [\alpha \xi^n + n(1 - \alpha \xi)t^{n-1}] \right| \frac{1}{(n-1)!} + \frac{1}{(n-1)!} \right\} \int_0^1 p(s) \psi(\|x\|_\infty) ds \leq \\ & \leq \frac{1}{(n-1)!} \left(1 + \frac{\delta_1 + \delta_2}{|\Delta|} \right) \psi(\|x\|_\infty) \int_0^1 p(s) ds \leq \\ & \leq \frac{1}{(n-1)!} \left(1 + \frac{\delta_1 + \delta_2}{|\Delta|} \right) \psi(r) \int_0^1 p(s) ds. \end{aligned}$$

where Δ , δ_1 , and δ_2 are given by (2.3), (3.2) and (3.3) respectively. Thus,

$$\|h\|_\infty \leq \frac{1}{(n-1)!} \left(1 + \frac{\delta_1 + \delta_2}{|\Delta|}\right) \psi(r) \int_0^1 p(s) ds.$$

Now we show that Ω maps bounded sets into equicontinuous sets of $C([0, 1], \mathbb{R})$. Let $t', t'' \in [0, 1]$ with $t' < t''$ and $x \in B_r$, where B_r is a bounded set of $C([0, 1], \mathbb{R})$. For each $h \in \Omega(x)$, we obtain

$$\begin{aligned} & |h(t'') - h(t')| = \\ & = \left| \int_0^{t''} \frac{(t'' - s)^{n-1}}{(n-1)!} f(s) ds - \int_0^{t'} \frac{(t' - s)^{n-1}}{(n-1)!} f(s) ds + \right. \\ & \quad + \frac{|(t'')^{n-1} - (t')^{n-1}|}{\Delta} \left[-\alpha(1 - \beta(1 - \eta)) \int_0^\xi \left(\int_0^s \frac{(s-m)^{n-1}}{(n-1)!} f(m) dm \right) ds + \right. \\ & \quad + \beta(1 - \alpha\xi) \int_\eta^1 \left(\int_0^s \frac{(s-m)^{n-1}}{(n-1)!} f(m) dm \right) ds - \\ & \quad \left. - (1 - \alpha\xi) \int_0^1 \frac{(1-s)^{n-1}}{(n-1)!} f(s) ds \right] \Big| \leq \\ & \leq \left| \int_0^{t'} \frac{(t' - s)^{n-1} - (t'' - s)^{n-1}}{(n-1)!} f(s) ds - \int_{t'}^{t''} \frac{(t'' - s)^{n-1}}{(n-1)!} f(s) ds \right| + \\ & \quad + \left| \frac{|(t'')^{n-1} - (t')^{n-1}|}{\Delta} \left[-\alpha(1 - \beta(1 - \eta)) \int_0^\xi \left(\int_0^s \frac{(s-m)^{n-1}}{(n-1)!} f(m) dm \right) ds + \right. \right. \\ & \quad + \beta(1 - \alpha\xi) \int_\eta^1 \left(\int_0^s \frac{(s-m)^{n-1}}{(n-1)!} f(m) dm \right) ds - \\ & \quad \left. - (1 - \alpha\xi) \int_0^1 \frac{(1-s)^{n-1}}{(n-1)!} f(s) ds \right] \Big|. \end{aligned}$$

Obviously the right hand side of the above inequality tends to zero independently of $x \in B_r$ as $t'' - t' \rightarrow 0$. As Ω satisfies the above three assumptions, therefore it follows by the Ascoli-Arzelá theorem that $\Omega : C([0, 1], \mathbb{R}) \rightarrow \mathcal{P}(C([0, 1], \mathbb{R}))$ is completely continuous.

In our next step, we show that Ω has a closed graph. Let $x_n \rightarrow x_*$, $h_n \in \Omega(x_n)$ and $h_n \rightarrow h_*$. Then we need to show that $h_* \in \Omega(x_*)$. Associated with $h_n \in \Omega(x_n)$, there exists $f_n \in S_{F, x_n}$ such that for each $t \in [0, 1]$,

$$\begin{aligned}
h_n(t) = & \int_0^t \frac{(t-s)^{n-1}}{(n-1)!} f_n(s) ds + \\
& + \frac{1}{n\Delta} \left[\alpha(n - \beta(1 - \eta^n)) \int_0^\xi \left(\int_0^s \frac{(s-m)^{n-1}}{(n-1)!} f_n(m) dm \right) ds + \right. \\
& + \alpha\beta\xi^n \int_\eta^1 \left(\int_0^s \frac{(s-m)^{n-1}}{(n-1)!} f_n(t)(m) dm \right) ds - \alpha\xi^n \int_0^1 \frac{(1-s)^{n-1}}{(n-1)!} f_n(s) ds \left. \right] + \\
& + \frac{t^{n-1}}{\Delta} \left[-\alpha(1 - \beta(1 - \eta)) \int_0^\xi \left(\int_0^s \frac{(s-m)^{n-1}}{(n-1)!} f_n(m) dm \right) ds + \right. \\
& + \beta(1 - \alpha\xi) \int_\eta^1 \left(\int_0^s \frac{(s-m)^{n-1}}{(n-1)!} f_n(m) dm \right) ds - \\
& \left. - (1 - \alpha\xi) \int_0^1 \frac{(1-s)^{n-1}}{(n-1)!} f_n(s) ds \right].
\end{aligned}$$

Thus we have to show that there exists $f_* \in S_{F,x_*}$ such that for each $t \in [0, 1]$,

$$\begin{aligned}
h_*(t) = & \int_0^t \frac{(t-s)^{n-1}}{(n-1)!} f_*(s) ds + \\
& + \frac{1}{n\Delta} \left[\alpha(n - \beta(1 - \eta^n)) \int_0^\xi \left(\int_0^s \frac{(s-m)^{n-1}}{(n-1)!} f_*(m) dm \right) ds + \right. \\
& + \alpha\beta\xi^n \int_\eta^1 \left(\int_0^s \frac{(s-m)^{n-1}}{(n-1)!} f_*(m) dm \right) ds - \alpha\xi^n \int_0^1 \frac{(1-s)^{n-1}}{(n-1)!} f_*(s) ds \left. \right] + \\
& + \frac{t^{n-1}}{\Delta} \left[-\alpha(1 - \beta(1 - \eta)) \int_0^\xi \left(\int_0^s \frac{(s-m)^{n-1}}{(n-1)!} f_*(m) dm \right) ds + \right. \\
& + \beta(1 - \alpha\xi) \int_\eta^1 \left(\int_0^s \frac{(s-m)^{n-1}}{(n-1)!} f_*(m) dm \right) ds - \\
& \left. - (1 - \alpha\xi) \int_0^1 \frac{(1-s)^{n-1}}{(n-1)!} f_*(s) ds \right].
\end{aligned}$$

Let us consider the continuous linear operator $\Theta : L^1([0, 1], \mathbb{R}) \rightarrow C([0, 1], \mathbb{R})$ given by

$$\begin{aligned}
 f \mapsto \Theta(f)(t) = & \int_0^t \frac{(t-s)^{n-1}}{(n-1)!} f(s) ds + \\
 & + \frac{1}{n\Delta} \left[\alpha(n - \beta(1 - \eta^n)) \int_0^\xi \left(\int_0^s \frac{(s-m)^{n-1}}{(n-1)!} f(m) dm \right) ds + \right. \\
 & + \alpha\beta\xi^n \int_\eta^1 \left(\int_0^s \frac{(s-m)^{n-1}}{(n-1)!} f(m) dm \right) ds - \alpha\xi^n \int_0^1 \frac{(1-s)^{n-1}}{(n-1)!} f(s) ds \left. \right] + \\
 & + \frac{t^{n-1}}{\Delta} \left[-\alpha(1 - \beta(1 - \eta)) \int_0^\xi \left(\int_0^s \frac{(s-m)^{n-1}}{(n-1)!} f(m) dm \right) ds + \right. \\
 & + \beta(1 - \alpha\xi) \int_\eta^1 \left(\int_0^s \frac{(s-m)^{n-1}}{(n-1)!} f(m) dm \right) ds - \\
 & \left. - (1 - \alpha\xi) \int_0^1 \frac{(1-s)^{n-1}}{(n-1)!} f(s) ds \right].
 \end{aligned}$$

Observe that

$$\begin{aligned}
 \|h_n(t) - h_*(t)\| = & \left\| \int_0^t \frac{(t-s)^{n-1}}{(n-1)!} (f_n(s) - f_*(s)) ds + \right. \\
 & + \frac{1}{n\Delta} \left[\alpha(n - \beta(1 - \eta^n)) \int_0^\xi \left(\int_0^s \frac{(s-m)^{n-1}}{(n-1)!} (f_n(m) - f_*(m)) dm \right) ds + \right. \\
 & + \alpha\beta\xi^n \int_\eta^1 \left(\int_0^s \frac{(s-m)^{n-1}}{(n-1)!} (f_n(m) - f_*(m)) dm \right) ds - \\
 & \left. - \alpha\xi^n \int_0^1 \frac{(1-s)^{n-1}}{(n-1)!} (f_n(s) - f_*(s)) ds \right] + \\
 & + \frac{t^{n-1}}{\Delta} \left[-\alpha(1 - \beta(1 - \eta)) \int_0^\xi \left(\int_0^s \frac{(s-m)^{n-1}}{(n-1)!} (f_n(m) - f_*(m)) dm \right) ds + \right. \\
 & + \beta(1 - \alpha\xi) \int_\eta^1 \left(\int_0^s \frac{(s-m)^{n-1}}{(n-1)!} (f_n(m) - f_*(m)) dm \right) ds - \\
 & \left. - (1 - \alpha\xi) \int_0^1 \frac{(1-s)^{n-1}}{(n-1)!} (f_n(s) - f_*(s)) ds \right] \left\| \rightarrow 0 \text{ as } n \rightarrow \infty.
 \end{aligned}$$

Thus, it follows by Lemma 2.5 that $\Theta \circ S_F$ is a closed graph operator. Further, we have $h_n(t) \in \Theta(S_{F,x_n})$. Since $x_n \rightarrow x_*$, therefore, we have

$$\begin{aligned} h_*(t) &= \int_0^t \frac{(t-s)^{n-1}}{(n-1)!} f_*(s) ds + \\ &+ \frac{1}{n\Delta} \left[\alpha(n - \beta(1 - \eta^n)) \int_0^\xi \left(\int_0^s \frac{(s-m)^{n-1}}{(n-1)!} f_*(m) dm \right) ds + \right. \\ &+ \alpha\beta\xi^n \int_\eta^1 \left(\int_0^s \frac{(s-m)^{n-1}}{(n-1)!} f_*(m) dm \right) ds - \alpha\xi^n \int_0^1 \frac{(1-s)^{n-1}}{(n-1)!} f_*(s) ds \left. \right] + \\ &+ \frac{t^{n-1}}{\Delta} \left[-\alpha(1 - \beta(1 - \eta)) \int_0^\xi \left(\int_0^s \frac{(s-m)^{n-1}}{(n-1)!} f_*(m) dm \right) ds + \right. \\ &+ \beta(1 - \alpha\xi) \int_\eta^1 \left(\int_0^s \frac{(s-m)^{n-1}}{(n-1)!} f_*(m) dm \right) ds - \\ &\left. - (1 - \alpha\xi) \int_0^1 \frac{(1-s)^{n-1}}{(n-1)!} f_*(s) ds \right] \end{aligned}$$

for some $f_* \in S_{F,x_*}$.

Finally, we discuss a priori bounds on solutions. Let x be a solution of (1.1). Then there exists $f \in L^1([0, 1], \mathbb{R})$ with $f \in S_{F,x}$ such that, for $t \in [0, 1]$, we have

$$\begin{aligned} x(t) &= \int_0^t \frac{(t-s)^{n-1}}{(n-1)!} f(s) ds + \\ &+ \frac{1}{n\Delta} \left[\alpha(n - \beta(1 - \eta^n)) \int_0^\xi \left(\int_0^s \frac{(s-m)^{n-1}}{(n-1)!} f(m) dm \right) ds + \right. \\ &+ \alpha\beta\xi^n \int_\eta^1 \left(\int_0^s \frac{(s-m)^{n-1}}{(n-1)!} f(m) dm \right) ds - \alpha\xi^n \int_0^1 \frac{(1-s)^{n-1}}{(n-1)!} f(s) ds \left. \right] + \\ &+ \frac{t^{n-1}}{\Delta} \left[-\alpha(1 - \beta(1 - \eta)) \int_0^\xi \left(\int_0^s \frac{(s-m)^{n-1}}{(n-1)!} f(m) dm \right) ds + \right. \\ &+ \beta(1 - \alpha\xi) \int_\eta^1 \left(\int_0^s \frac{(s-m)^{n-1}}{(n-1)!} y(m) dm \right) ds - \\ &\left. - (1 - \alpha\xi) \int_0^1 \frac{(1-s)^{n-1}}{(n-1)!} f(s) ds \right]. \end{aligned}$$

In view of (H_2) , for each $t \in [0, 1]$, we obtain

$$\begin{aligned}
 |x(t)| &\leq \left| \frac{\alpha}{n\Delta} [(n - \beta(1 - \eta^n)) - n(1 - \beta(1 - \eta))]t^{n-1} \right| \times \\
 &\quad \times \int_0^\xi \left(\int_0^s \frac{(s-m)^{n-1}}{(n-1)!} |f(m)| dm \right) ds + \\
 &\quad + \left| \frac{\beta}{n\Delta} [\alpha\xi^n + n(1 - \alpha\xi)t^{n-1}] \right| \int_\eta^1 \left(\int_0^s \frac{(s-m)^{n-1}}{(n-1)!} |f(m)| dm \right) ds + \\
 &\quad + \left| \frac{1}{n\Delta} [\alpha\xi^n + n(1 - \alpha\xi)t^{n-1}] \right| \int_0^1 \frac{(1-s)^{n-1}}{(n-1)!} |f(s)| ds + \\
 &\quad + \int_0^t \frac{(t-s)^{n-1}}{(n-1)!} |f(s)| ds \leq \\
 &\leq \left\{ \left| \frac{\alpha}{n\Delta} [(n - \beta(1 - \eta^n)) - n(1 - \beta(1 - \eta))]t^{n-1} \right| \frac{\xi^n}{(n-1)!} + \right. \\
 &\quad + \left| \frac{\beta}{n\Delta} [\alpha + n(1 - \alpha\xi)t^{n-1}] \right| \frac{1}{(n-1)!} + \\
 &\quad \left. + \left| \frac{1}{n\Delta} [\alpha\xi^n + n(1 - \alpha\xi)t^{n-1}] \right| \left[\frac{1}{(n-1)!} + \frac{1}{(n-1)!} \right] \int_0^1 |f(s)| ds \right\} \leq \\
 &\leq \frac{1}{(n-1)!} \left(1 + \frac{\delta_1 + \delta_2}{|\Delta|} \right) \psi(\|x\|_\infty) \int_0^1 p(s) ds.
 \end{aligned}$$

Consequently, we have

$$\frac{\|x\|_\infty}{\frac{1}{(n-1)!} \left(1 + \frac{\delta_1 + \delta_2}{|\Delta|} \right) \psi(\|x\|_\infty) \|p\|_{L^1}} \leq 1.$$

In view of (H_3) , there exists M such that $\|x\|_\infty \neq M$. Let us set

$$U = \{x \in C([0, 1], \mathbb{R}) : \|x\|_\infty < M + 1\}.$$

Note that the operator $\Omega : \bar{U} \rightarrow \mathcal{P}(C([0, 1], \mathbb{R}))$ is upper semicontinuous and completely continuous. From the choice of U , there is no $x \in \partial U$ such that $x \in \mu\Omega(x)$ for some $\mu \in (0, 1)$. Consequently, by Lemma 2.6, we deduce that Ω has a fixed point $x \in \bar{U}$ which is a solution of the problem (1.1). This completes the proof. \square

As a next result, we study the case when F is not necessarily convex valued. Our strategy to deal with this problems is based on the nonlinear alternative of Leray Schauder type together with the selection theorem of Bressan and Colombo [12] for lower semi-continuous maps with decomposable values.

Theorem 3.2. Assume that (H_2) , (H_3) and the following conditions hold:

(H_4) $F : [0, 1] \times \mathbb{R} \rightarrow \mathcal{P}(\mathbb{R})$ is a nonempty compact-valued multivalued map such that:

(a) $(t, x) \mapsto F(t, x)$ is $\mathcal{L} \otimes \mathcal{B}$ measurable,

(b) $x \mapsto F(t, x)$ is lower semicontinuous for each $t \in [0, 1]$,

(H_5) for each $\sigma > 0$, there exists $\varphi_\sigma \in L^1([0, 1], \mathbb{R}^+)$ such that

$$\|F(t, x)\| = \sup\{|y| : y \in F(t, x)\} \leq \varphi_\sigma(t) \text{ for all } \|x\|_\infty \leq \sigma \text{ and for a.e. } t \in [0, 1].$$

Then the boundary value problem (1.1) has at least one solution on $[0, 1]$.

Proof. It follows from (H_4) and (H_5) that F is of l.s.c. type. Then from Lemma 2.7, there exists a continuous function $f : C([0, 1], \mathbb{R}) \rightarrow L^1([0, 1], \mathbb{R})$ such that $f(x) \in \mathcal{F}(x)$ for all $x \in C([0, 1], \mathbb{R})$.

Consider the problem

$$\begin{cases} x^{(n)}(t) = f(x(t)), & 0 < t < 1, \\ x(0) = \alpha \int_0^\xi x(s) ds, \quad x'(0) = 0, \quad x''(0) = 0, \dots, \quad x^{(n-2)}(0) = 0, \\ x(1) = \beta \int_\eta^1 x(s) ds, \quad 0 < \xi < \eta < 1. \end{cases} \quad (3.4)$$

Observe that if $x \in C^2([0, 1], \mathbb{R})$ is a solution of (3.4), then x is a solution to the problem (1.1). In order to transform the problem (3.4) into a fixed point problem, we define the operator $\bar{\Omega}$ as

$$\begin{aligned} \bar{\Omega}x(t) = & \int_0^t \frac{(t-s)^{n-1}}{(n-1)!} f(x(s)) ds + \\ & + \frac{1}{n\Delta} \left[\alpha(n - \beta(1 - \eta^n)) \int_0^\xi \left(\int_0^s \frac{(s-m)^{n-1}}{(n-1)!} f(x(m)) dm \right) ds + \right. \\ & + \alpha\beta\xi^n \int_\eta^1 \left(\int_0^s \frac{(s-m)^{n-1}}{(n-1)!} f(x(m)) dm \right) ds - \alpha\xi^n \int_0^1 \frac{(1-s)^{n-1}}{(n-1)!} f(x(s)) ds \Big] + \\ & + \frac{t^{n-1}}{\Delta} \left[-\alpha(1 - \beta(1 - \eta)) \int_0^\xi \left(\int_0^s \frac{(s-m)^{n-1}}{(n-1)!} f(x(m)) dm \right) ds + \right. \\ & + \beta(1 - \alpha\xi) \int_\eta^1 \left(\int_0^s \frac{(s-m)^{n-1}}{(n-1)!} f(x(m)) dm \right) ds - \\ & \left. - (1 - \alpha\xi) \int_0^1 \frac{(1-s)^{n-1}}{(n-1)!} f(x(s)) ds \right]. \end{aligned}$$

It can easily be shown that $\bar{\Omega}$ is continuous and completely continuous. The remaining part of the proof is similar to that of Theorem 3.1, so we omit it. This completes the proof. \square

Now we prove the existence of solutions for the problem (1.1) with a nonconvex valued right hand side by applying a fixed point theorem for a multivalued map due to Covitz and Nadler [15].

Theorem 3.3. *Assume that the following conditions hold:*

- (H₆) $F : [0, 1] \times \mathbb{R} \rightarrow P_{cp}(\mathbb{R})$ is such that $F(\cdot, x) : [0, 1] \rightarrow P_{cp}(\mathbb{R})$ is measurable for each $x \in \mathbb{R}$,
- (H₇) $H_d(F(t, x), F(t, \bar{x})) \leq m(t)|x - \bar{x}|$ for almost all $t \in [0, 1]$ and $x, \bar{x} \in \mathbb{R}$ with $m \in L^1([0, 1], \mathbb{R}^+)$ and $d(0, F(t, 0)) \leq m(t)$ for almost all $t \in [0, 1]$.

Then the boundary value problem (1.1) has at least one solution on $[0, 1]$ if

$$\frac{1}{(n-1)!} \left(1 + \frac{\delta_1 + \delta_2}{|\Delta|} \right) \|m\|_{L^1} < 1.$$

Proof. Observe that the set $S_{F,x}$ is nonempty for each $x \in C([0, 1], \mathbb{R})$ by the assumption (H₆), so F has a measurable selection (see [13, Theorem III.6]). Now we show that the operator Ω satisfies the assumptions of Lemma 2.8. To show that $\Omega(x) \in P_{cl}((C[0, 1], \mathbb{R}))$ for each $x \in C([0, 1], \mathbb{R})$, let $\{u_n\}_{n \geq 0} \in \Omega(x)$ be such that $u_n \rightarrow u$ ($n \rightarrow \infty$) in $C([0, 1], \mathbb{R})$. Then $u \in C([0, 1], \mathbb{R})$ and there exists $v_n \in S_{F,x}$ such that, for each $t \in [0, 1]$,

$$\begin{aligned} u_n(t) = & \int_0^t \frac{(t-s)^{n-1}}{(n-1)!} v_n(s) ds + \\ & + \frac{1}{n\Delta} \left[\alpha(n - \beta(1 - \eta^n)) \int_0^\xi \left(\int_0^s \frac{(s-m)^{n-1}}{(n-1)!} v_n(m) dm \right) ds + \right. \\ & + \alpha\beta\xi^n \int_\eta^1 \left(\int_0^s \frac{(s-m)^{n-1}}{(n-1)!} v_n(m) dm \right) ds - \alpha\xi^n \int_0^1 \frac{(1-s)^{n-1}}{(n-1)!} v_n(s) ds \left. \right] + \\ & + \frac{t^{n-1}}{\Delta} \left[-\alpha(1 - \beta(1 - \eta)) \int_0^\xi \left(\int_0^s \frac{(s-m)^{n-1}}{(n-1)!} v_n(m) dm \right) ds + \right. \\ & + \beta(1 - \alpha\xi) \int_\eta^1 \left(\int_0^s \frac{(s-m)^{n-1}}{(n-1)!} v_n(m) dm \right) ds - \\ & \left. - (1 - \alpha\xi) \int_0^1 \frac{(1-s)^{n-1}}{(n-1)!} v_n(s) ds \right]. \end{aligned}$$

As F has compact values, we pass onto a subsequence to obtain that v_n converges to v in $L^1([0, 1], \mathbb{R})$. Thus, $v \in S_{F,x}$ and for each $t \in [0, 1]$,

$$\begin{aligned} u_n(t) \rightarrow u(t) &= \int_0^t \frac{(t-s)^{n-1}}{(n-1)!} v(s) ds + \\ &+ \frac{1}{n\Delta} \left[\alpha(n - \beta(1 - \eta^n)) \int_0^\xi \left(\int_0^s \frac{(s-m)^{n-1}}{(n-1)!} v(m) dm \right) ds + \right. \\ &+ \alpha\beta\xi^n \int_\eta^1 \left(\int_0^s \frac{(s-m)^{n-1}}{(n-1)!} v(m) dm \right) ds - \alpha\xi^n \int_0^1 \frac{(1-s)^{n-1}}{(n-1)!} v(s) ds \left. \right] + \\ &+ \frac{t^{n-1}}{\Delta} \left[-\alpha(1 - \beta(1 - \eta)) \int_0^\xi \left(\int_0^s \frac{(s-m)^{n-1}}{(n-1)!} v(m) dm \right) ds + \right. \\ &+ \beta(1 - \alpha\xi) \int_\eta^1 \left(\int_0^s \frac{(s-m)^{n-1}}{(n-1)!} v(m) dm \right) ds - (1 - \alpha\xi) \int_0^1 \frac{(1-s)^{n-1}}{(n-1)!} v(s) ds \left. \right]. \end{aligned}$$

Hence, $u \in \Omega(x)$.

Next we show that there exists $\gamma < 1$ such that

$$H_d(\Omega(x), \Omega(\bar{x})) \leq \gamma \|x - \bar{x}\|_\infty \quad \text{for each } x, \bar{x} \in C([0, 1], \mathbb{R}).$$

Let $x, \bar{x} \in C([0, 1], \mathbb{R})$ and $h_1 \in \Omega(x)$. Then there exists $v_1(t) \in F(t, x(t))$ such that, for each $t \in [0, 1]$,

$$\begin{aligned} h_1(t) &= \int_0^t \frac{(t-s)^{n-1}}{(n-1)!} v_1(s) ds \\ &+ \frac{1}{n\Delta} \left[\alpha(n - \beta(1 - \eta^n)) \int_0^\xi \left(\int_0^s \frac{(s-m)^{n-1}}{(n-1)!} v_1(m) dm \right) ds + \right. \\ &+ \alpha\beta\xi^n \int_\eta^1 \left(\int_0^s \frac{(s-m)^{n-1}}{(n-1)!} v_1(m) dm \right) ds - \alpha\xi^n \int_0^1 \frac{(1-s)^{n-1}}{(n-1)!} v_1(s) ds \left. \right] + \\ &+ \frac{t^{n-1}}{\Delta} \left[-\alpha(1 - \beta(1 - \eta)) \int_0^\xi \left(\int_0^s \frac{(s-m)^{n-1}}{(n-1)!} v_1(m) dm \right) ds + \right. \\ &+ \beta(1 - \alpha\xi) \int_\eta^1 \left(\int_0^s \frac{(s-m)^{n-1}}{(n-1)!} v_1(m) dm \right) ds - (1 - \alpha\xi) \int_0^1 \frac{(1-s)^{n-1}}{(n-1)!} v_1(s) ds \left. \right]. \end{aligned}$$

By (H_7) , we have

$$H_d(F(t, x), F(t, \bar{x})) \leq m(t) |x(t) - \bar{x}(t)|.$$

So, there exists $w \in F(t, \bar{x}(t))$ such that

$$|v_1(t) - w| \leq m(t)|x(t) - \bar{x}(t)|, \quad t \in [0, 1].$$

Define $U : [0, 1] \rightarrow \mathcal{P}(\mathbb{R})$ by

$$U(t) = \{w \in \mathbb{R} : |v_1(t) - w| \leq m(t)|x(t) - \bar{x}(t)|\}.$$

Since the multivalued operator $U(t) \cap F(t, \bar{x}(t))$ is measurable ([13, Proposition III.4]), there exists a function $v_2(t)$ which is a measurable selection for V . So $v_2(t) \in F(t, \bar{x}(t))$ and for each $t \in [0, 1]$, we have $|v_1(t) - v_2(t)| \leq m(t)|x(t) - \bar{x}(t)|$.

For each $t \in [0, 1]$, let us define

$$\begin{aligned} h_2(t) &= \int_0^t \frac{(t-s)^{n-1}}{(n-1)!} v_2(s) ds + \\ &+ \frac{1}{n\Delta} \left[\alpha(n - \beta(1 - \eta^n)) \int_0^\xi \left(\int_0^s \frac{(s-m)^{n-1}}{(n-1)!} v_2(m) dm \right) ds + \right. \\ &+ \alpha\beta\xi^n \int_\eta^1 \left(\int_0^s \frac{(s-m)^{n-1}}{(n-1)!} v_2(m) dm \right) ds - \alpha\xi^n \int_0^1 \frac{(1-s)^{n-1}}{(n-1)!} v_2(s) ds \left. \right] + \\ &+ \frac{t^{n-1}}{\Delta} \left[-\alpha(1 - \beta(1 - \eta)) \int_0^\xi \left(\int_0^s \frac{(s-m)^{n-1}}{(n-1)!} v_2(m) dm \right) ds + \right. \\ &+ \beta(1 - \alpha\xi) \int_\eta^1 \left(\int_0^s \frac{(s-m)^{n-1}}{(n-1)!} v_2(m) dm \right) ds - \\ &\left. - (1 - \alpha\xi) \int_0^1 \frac{(1-s)^{n-1}}{(n-1)!} v_2(s) ds \right]. \end{aligned}$$

Thus,

$$\begin{aligned}
|h_1(t) - h_2(t)| &\leq \int_0^t \frac{(t-s)^{n-1}}{(n-1)!} |v_1(s) - v_2(s)| ds + \\
&+ \frac{1}{n\Delta} \left[\alpha(n - \beta(1 - \eta^n)) \int_0^\xi \left(\int_0^s \frac{(s-m)^{n-1}}{(n-1)!} |v_1(m) - v_2(m)| dm \right) ds + \right. \\
&+ \alpha\beta\xi^n \int_\eta^1 \left(\int_0^s \frac{(s-m)^{n-1}}{(n-1)!} |v_1(m) - v_2(m)| dm \right) ds - \\
&- \alpha\xi^n \int_0^1 \frac{(1-s)^{n-1}}{(n-1)!} |v_1(s) - v_2(s)| ds \left. \right] + \\
&+ \frac{t^{n-1}}{\Delta} \left[-\alpha(1 - \beta(1 - \eta)) \int_0^\xi \left(\int_0^s \frac{(s-m)^{n-1}}{(n-1)!} |v_1(m) - v_2(m)| dm \right) ds + \right. \\
&+ \beta(1 - \alpha\xi) \int_\eta^1 \left(\int_0^s \frac{(s-m)^{n-1}}{(n-1)!} |v_1(m) - v_2(m)| dm \right) ds - \\
&- (1 - \alpha\xi) \int_0^1 \frac{(1-s)^{n-1}}{(n-1)!} |v_1(s) - v_2(s)| ds \left. \right] \leq \\
&\leq \frac{1}{(n-1)!} \left(1 + \frac{\delta_1 + \delta_2}{|\Delta|} \right) \int_0^1 m(s) \|x - \bar{x}\| ds.
\end{aligned}$$

Hence,

$$\|h_1 - h_2\|_\infty \leq \frac{1}{(n-1)!} \left(1 + \frac{\delta_1 + \delta_2}{|\Delta|} \right) \|m\|_{L^1} \|x - \bar{x}\|_\infty.$$

Analogously, interchanging the roles of x and \bar{x} , we obtain

$$\begin{aligned}
H_d(\Omega(x), \Omega(\bar{x})) &\leq \gamma \|x - \bar{x}\|_\infty \leq \\
&\leq \frac{1}{(n-1)!} \left(1 + \frac{\delta_1 + \delta_2}{|\Delta|} \right) \|m\|_{L^1} \|x - \bar{x}\|_\infty.
\end{aligned}$$

Since Ω is a contraction, it follows by Lemma 2.8 that Ω has a fixed point x which is a solution of (1.1). This completes the proof. \square

REFERENCES

- [1] A.R. Aftabizadeh, *Existence and uniqueness theorems for fourth-order boundary value problems*, J. Math. Anal. Appl. **116** (1986), 415–426.
- [2] B. Ahmad, A. Alsaedi, B. Alghamdi, *Analytic approximation of solutions of the forced Duffing equation with integral boundary conditions*, Nonlinear Anal. Real World Appl. **9** (2008), 1727–1740.
- [3] B. Ahmad, S.K. Ntouyas, *Existence results for nonlocal boundary value problems for fractional differential inclusions*, International Electronic Journal of Pure and Applied Mathematics **1** (2010) 2, 101–114.
- [4] B. Ahmad, S.K. Ntouyas, *Some existence results for boundary value problems of fractional differential inclusions with non-separated boundary conditions*, Electron. J. Qual. Theory Differ. Equ. **71** (2010), 1–17.
- [5] B. Ahmad, S.K. Ntouyas, *Existence results for a second order boundary value problem with four-point nonlocal integral boundary conditions*, Commun. Appl. Nonlinear Anal. **17** (2010), 43–52.
- [6] J. Andres, *A four-point boundary value problem for the second-order ordinary differential equations*, Arch. Math. (Basel) **53** (1989), 384–389.
- [7] J. Andres, *Four-point and asymptotic boundary value problems via a possible modification of Poincaré’s mapping*, Math. Nachr. **149** (1990), 155–162.
- [8] E.O. Ayoola, *Quantum stochastic differential inclusions satisfying a general Lipschitz condition*, Dynam. Systems Appl. **17** (2008), 487–502
- [9] M. Benaïm, J. Hofbauer, S. Sorin, *Stochastic approximations and differential inclusions II. Applications*, Math. Oper. Res. **31** (2006), 673–695.
- [10] A.V. Bicaдзе, A.A. Samarskii, *Some elementary generalizations of linear elliptic boundary value problems*, Anal. Dokl. Akad. Nauk SSSR **185** (1969), 739–740 [in Russian].
- [11] A. Boucherif, *Second-order boundary value problems with integral boundary conditions*, Nonlinear Anal. **70** (2009), 364–371.
- [12] A. Bressan, G. Colombo, *Extensions and selections of maps with decomposable values*, Studia Math. **90** (1988), 69–86.
- [13] C. Castaing, M. Valadier, *Convex Analysis and Measurable Multifunctions*, Lecture Notes in Mathematics 580, Springer-Verlag, Berlin-Heidelberg-New York, 1977.
- [14] Y.K. Chang, W.T. Li, *Existence results for second order impulsive functional differential inclusions*, J. Math. Anal. Appl. **301** (2005), 477–490.
- [15] H. Covitz, S.B. Nadler (Jr.), *Multivalued contraction mappings in generalized metric spaces*, Israel J. Math. **8** (1970), 5–11.
- [16] K. Deimling, *Multivalued Differential Equations*, Walter De Gruyter, Berlin-New York, 1992.
- [17] P.W. Eloe, B. Ahmad, *Positive solutions of a nonlinear n th order boundary value problem with nonlocal conditions*, Appl. Math. Lett. **18** (2005), 521–527.

- [18] A. Granas, J. Dugundji, *Fixed Point Theory*, Springer-Verlag, New York, 2005.
- [19] J.R. Graef, J.R.L. Webb, *Third order boundary value problems with nonlocal boundary conditions*, *Nonlinear Anal.* **71** (2009), 1542–1551.
- [20] M. Greguš, F. Neumann, F.M. Arscott, *Three-point boundary value problems in differential equations*, *Proc. London Math. Soc.* **3** (1964), 459–470.
- [21] C.P. Gupta, *A second order m -point boundary value problem at resonance*, *Nonlinear Anal.* **24** (1995), 1483–1489.
- [22] Sh. Hu, N. Papageorgiou, *Handbook of Multivalued Analysis, Volume I: Theory*, Kluwer, Dordrecht, 1997.
- [23] V.A. Il'in, E.I. Moiseev, *Nonlocal boundary-value problem of the first kind for a Sturm-Liouville operator in its differential and finite difference aspects*, *Differential Equations* **23** (1987), 803–810.
- [24] M. Kisielewicz, *Differential Inclusions and Optimal Control*, Kluwer, Dordrecht, 1991.
- [25] A. Lasota, Z. Opial, *An application of the Kakutani-Ky Fan theorem in the theory of ordinary differential equations*, *Bull. Acad. Polon. Sci. Ser. Sci. Math. Astronom. Phys.* **13** (1965), 781–786.
- [26] R. Ma, *Multiple positive solutions for nonlinear m -point boundary value problems*, *Appl. Math. Comput.* **148** (2004), 249–262.
- [27] S.K. Ntouyas, *Nonlocal Initial and Boundary Value Problems: A Survey*, *Handbook on Differential Equations: Ordinary Differential Equations*, A. Canada, P. Drabek and A. Fonda (Eds.), Elsevier Science B.V., 2005, 459–555.
- [28] S.K. Ntouyas, *Neumann boundary value problems for impulsive differential inclusions*, *Electron. J. Qual. Theory Differ. Equ.* 2009, Special Edition I, No. 22, 13 pp.
- [29] G.V. Smirnov, *Introduction to the Theory of Differential Inclusions*, American Mathematical Society, Providence, RI, 2002.
- [30] L. Wang, M. Pei, W. Ge, *Existence and approximation of solutions for nonlinear second-order four-point boundary value problems*, *Math. Comput. Modelling* **50** (2009), 1348–1359.
- [31] J.R.L. Webb, G. Infante, *Positive solutions of nonlocal boundary value problems: A unified approach*, *J. London Math. Soc.* **74** (2006), 673–693.

Bashir Ahmad
bashir_qau@yahoo.com

King Abdulaziz University,
Faculty of Science
Department of Mathematics
P.O. Box 80203, Jeddah 21589, Saudi Arabia

Sotiris K. Ntouyas
sntouyas@uoi.gr

University of Ioannina
Department of Mathematics
451 10 Ioannina, Greece

Received: January 11, 2011.

Revised: April 6, 2011.

Accepted: April 7, 2011.