

ON THE UNIQUENESS OF MINIMAL PROJECTIONS IN BANACH SPACES

Ewa Szlachtowska and Dominik Mielczarek

Abstract. Let X be a uniformly convex Banach space with a continuous semi-inner product. We investigate the relation of orthogonality in X and generalized projections acting on X . We prove uniqueness of orthogonal and co-orthogonal projections.

Keywords: minimal projection, orthogonal projection, co-orthogonal projection, uniqueness of norm-one projection.

Mathematics Subject Classification: 41A65, 46B20, 46B25.

1. INTRODUCTION

In the theory of operators on a Hilbert space most of the terminology and techniques are developed by use of the inner-product. It is known that a Banach space can be represented as a semi-inner product space with a more general axiom system than that of a Hilbert space (see [10]). Hence, in a Banach space we can define orthogonality and transversality relations. A natural consequence of these relations are an orthogonal set M^\perp and a transversal set M^\top for a set M . In a Hilbert space X we have $M^\perp = M^\top$ and $X = M \oplus M^\perp$ for a closed subspace M of X . It turns out that there holds the decomposition theorem on a uniformly convex Banach space with a continuous semi-inner product. This result is presented in detail in Theorem 2.5. However, M^\perp is not always a linear subspace of X . If it were, the space X would have to be isomorphic to some Hilbert space by the Lindenstrauss-Tzafriri theorem [9], but this is not always true. In Theorem 3.10 we give conditions for M^\perp to be a subspace of X . If the set M^\perp is a subspace of the space X , then M is one-co-complemented and the converse of this statement is also true. In this paper a new definition of a generalized projection is given. The inspiration for this was the metric projection. The main result in this article is Theorem 3.5. We show that for a closed subspace M in a uniformly convex Banach space with a continuous semi-inner product there exists at most one homogenous generalized projection $P : X \rightarrow M$ satisfying the Lipschitz condition with the constant equal to one.

2. AUXILIARY RESULTS

To apply Hilbert space type methods to the theory of Banach spaces, G. Lumer [10] constructed a semi-inner product (s.i.p.) on a complex linear space X as a complex function $[\cdot, \cdot]$ on $X \times X$ with the following properties:

$$[\alpha x + \beta y, z] = \alpha[x, z] + \beta[y, z], \quad x, y, z \in X, \quad \alpha, \beta \in \mathbb{C}, \quad (2.1)$$

$$[x, \lambda y] = \bar{\lambda}[x, y], \quad x, y \in X, \quad \lambda \in \mathbb{C}. \quad (2.2)$$

$$[x, x] > 0 \quad \text{for } x \neq 0, \quad (2.3)$$

$$|[x, y]|^2 \leq [x, x][y, y], \quad x, y \in X. \quad (2.4)$$

$(X, [\cdot, \cdot])$ is called a complex space with semi-inner product.

The importance of a semi-inner product space (s.i.p.s.) is that every normed space can be represented as a semi-inner product space so that the theory of operators on a Banach space can be represented by Hilbert space type arguments.

Theorem 2.1 ([4], [10]). *A semi-inner product space $(X, [\cdot, \cdot])$ is a normed linear space with the norm*

$$\|x\| = [x, x]^{1/2}, \quad x \in X.$$

Every normed linear space can be made into a semi-inner product space (in general, in infinitely many different ways).

In a normed space X we set

$$S = \{x \in X : \|x\| = 1\}.$$

We introduce additional properties of the semi-inner product that will help us to carry over Hilbert space type arguments to the case of a Banach space. Note that a semi-inner product is continuous with respect to the first component. A very convenient property of a s.i.p. is continuity with respect to the second variable.

A s.i.p.s. X is called a *continuous s.i.p. space* when a semi-inner product satisfies the following additional condition:

for every $x, y \in S$,

$$\operatorname{Re}[y, x + \lambda y] \rightarrow \operatorname{Re}[y, x] \quad \text{for all real } \lambda \rightarrow 0. \quad (2.5)$$

The space X is a *uniformly continuous s.i.p.s.* if the above limit (2.5) is approached uniformly for all $(x, y) \in S \times S$.

Define a relation on a s.i.p. space which may be called an orthogonality relation. Let $x, y \in X$. We say that x is *normal* to y and y is *transversal* to x if $[y, x] = 0$. A vector $x \in X$ is normal to a subspace N and N is transversal to x if x is normal to all vectors from N .

For a normed space, R.C. James [6] studied the orthogonality relation (in the sense of Birkhoff) defined as follows:

A vector x is *orthogonal* to y in the sense of Birkhoff if

$$\|x + \lambda y\| \geq \|x\| \text{ for all } \lambda \in \mathbb{C}.$$

It is worth noting that orthogonality in the sense of Birkhoff is very close to the concept of an element of best approximation. It was shown that in a continuous s.i.p.s. an orthogonality relation is equivalent to a Birkhoff orthogonality relation (see [4]).

Theorem 2.2 ([4]). *In a continuous s.i.p.s. x is normal to y if and only if x is orthogonal to y in the sense of Birkhoff.*

Since a s.i.p. is not commutative, this orthogonality relation is not symmetric, i.e. if x is normal to y , then y is not necessarily normal to x . So, for a subset M of X we define an *orthogonal set* by

$$M^\perp = \{x \in X : \forall y \in M [y, x] = 0\}$$

and a *transversal set* by

$$M^\top = \{x \in X : \forall y \in M [x, y] = 0\}.$$

It is easy to see that

$$X^\perp = X^\top = \{0\}, \tag{2.6}$$

$$M \cap M^\perp = \{0\}, \tag{2.7}$$

$$M \cap M^\top = \{0\}. \tag{2.8}$$

2.1. THE DECOMPOSITION THEOREM

To extend Hilbert space type arguments to the theory of decomposition we need to impose an additional structure on a s.i.p. chiefly to guarantee the existence of normal vectors to closed subspaces.

A normed space is *uniformly convex* if given $\varepsilon \in (0, 2]$, there exists $\delta(\varepsilon) > 0$ such that for $x, y \in S$, $\|x - y\| > \varepsilon$ implies $\|x + y\|/2 \leq 1 - \delta(\varepsilon)$.

Recall the notion of strict convexity. A normed space is *strictly convex* if whenever $\|x\| + \|y\| = \|x + y\|$, where $x, y \neq 0$, then $y = \lambda x$ for some real $\lambda > 0$.

It is well known that uniform convexity implies strict convexity. The following two lemmas will help us to characterize a strictly convex space by the structure of the semi-inner product. We will also need them for further considerations. Note also that for linearly dependent elements we have equality in the Schwarz inequality.

Lemma 2.3 ([4]). *A s.i.p.s. is strictly convex if and only if whenever $[x, y] = \|x\|\|y\|$, where $x, y \neq 0$, then $y = \lambda x$ for some real $\lambda > 0$.*

Lemma 2.4 ([4]). *Let X be a strictly convex space with a semi-inner product. Let $y, z \in X$. If $[x, y] = [x, z]$ for all $x \in X$, then $y = z$.*

Let M and M' be subsets of a linear space X . We say that $X = M \oplus M'$ if and only if for $x \in X$ there exist unique elements $x_M \in M$, $x_{M'} \in M'$ such that $x = x_M + x_{M'}$ and $M \cap M' = \{0\}$.

We will prove that in a uniformly convex Banach space with a continuous semi-inner product we have

$$X = M \oplus M^\perp$$

for a closed subspace M of X .

Theorem 2.5. *Let X be a uniformly convex Banach space with a continuous semi-inner product. Let M be a closed subspace of X . Then each $x \in X$ can be uniquely decomposed in the form $x = y + z$ with $y \in M$ and $z \in M^\perp$.*

Proof. It is well known that, in a uniformly convex Banach space, for a closed subspace M and a vector $x \notin M$, there exists a unique nonzero vector $y \in M$ such that

$$\|x - y\| = d(x, M) = \inf\{\|x - y'\| : y' \in M\}.$$

Let us set $z = x - y$. Then z is normal to M .

In order to prove the uniqueness of the representation $x = y + z$ we assume that $x = y_1 + z_1 = y_2 + z_2$, where $y_1, y_2 \in M$ and $z_1, z_2 \in M^\perp$. It follows that $z_1 - z_2 = y_1 - y_2 \in M$. If $z_1 - z_2 \in M \cap M^\perp$, then $z_1 - z_2 = 0$ and $y_1 = y_2$. If $z_1 - z_2 \notin M^\perp$, then

$$\begin{aligned} 0 &= [z_1 - z_2, z_1] = [z_1, z_1] - [z_2, z_1] \geq \|z_1\|^2 - \|z_1\| \|z_2\|, \\ 0 &= [z_2 - z_1, z_2] = [z_2, z_2] - [z_1, z_2] \geq \|z_2\|^2 - \|z_1\| \|z_2\|. \end{aligned}$$

Therefore,

$$\|z_1\| = \|z_2\| \text{ and } \|z_1\| \|z_2\| = [z_1, z_2].$$

By the strict convexity of X , we obtain $z_1 = z_2$. This implies that $y_1 = y_2$. \square

In all that follows, we assume that X is a uniformly convex Banach space with a continuous semi-inner product and M is a proper closed subspace of X .

In this case, we can define a metric projection $P_m : X \rightarrow M$ such that $P_m(x)$ is an element that best approximates $x \in X$ with respect to M , i.e.

$$\|x - P_m(x)\| = \text{dist}(x, M).$$

In a Hilbert space we have $M^\perp = M^\top$. The following theorem shows the relationship between an orthogonal set and a transversal set.

Theorem 2.6. *Let X be a uniformly convex Banach space with a continuous semi-inner product. Let M be a closed subspace of X . Then*

$$\begin{aligned} M &\subset (M^\top)^\perp, \\ (M^\perp)^\top &= M. \end{aligned}$$

Proof. If $x \in M$, then $[y, x] = 0$ for $y \in M^\top$, hence

$$M \subset (M^\top)^\perp.$$

If $x \in M$, then we have $[x, y] = 0$ for $y \in M^\perp$, i.e. $x \in (M^\perp)^\top$.

Conversely, suppose that $x \in (M^\perp)^\top$, i.e. $[x, y] = 0$ for $y \in M^\perp$. By Theorem 2.5, there exist $x_1 \in M$ and $x_2 \in M^\perp$ such that $x = x_1 + x_2$. Then $[x_1 + x_2, y] = 0$. Hence $[x_2, y] = 0$ for all $y \in M^\perp$. Setting $y = x_2$, we deduce that $x_2 = 0$. Therefore, $x = x_1 \in M$. \square

It should be noted that the set M^\top is a closed subspace of X . According to Theorem 2.2 M^\perp is a closed subset (but not necessarily a subspace) of X .

Example 2.7. Let $X = l_p$, $1 < p < \infty$. Let us equip l_p with a semi-inner product given by

$$[y, x] = \begin{cases} \|x\|_p^{2-p} \sum_{k=1}^{\infty} y_k \overline{x_k} |x_k|^{p-2}, & x \neq 0, \\ 0, & x = 0, \end{cases}$$

where $\|x\|_p = \left(\sum_{k=1}^{\infty} |x_k|^p \right)^{1/p}$.

(i) Let $M = \text{span}\{e_1, e_2, \dots, e_n\}$, where $n \in \mathbb{N}$ and $(e_i)_{i=1}^n$ are elements from the standard basis in l_p . Then v is orthogonal to M if and only if $v(i) = 0$ for $i = 1, 2, \dots, n$. Note that in this case M^\perp is a linear subspace of l_p .

(ii) Take $n \in \mathbb{N}$, $n > 1$. Let $M = \text{span}\{e\}$, where $e(i) = 1$ for $i = 1, 2, \dots, n$ and $e(i) = 0$ otherwise. Then v is orthogonal to M if and only if $\sum_{i=1}^n |v(i)|^{p-2} \overline{v(i)} = 0$. In this case M^\perp is not a linear subspace of l_p .

3. ORTHOGONAL PROJECTIONS IN BANACH SPACES

3.1. GENERALIZED PROJECTIONS

Let X be a Banach space and M be a subspace of X . An operator $P : X \rightarrow M$ is called a *generalized projection* if it satisfies the following conditions:

- (P1) P is continuous;
- (P2) $\ker P = \{x - Px : x \in X\}$;
- (P3) $X = \ker P \oplus M$;
- (P4) For every $x \in X$, we set $P(x) = x_M$, where $x = x_{\ker P} + x_M$, $x_{\ker P} \in \ker P$, $x_M \in M$.

An inspiration to define a generalized projection was the metric projection P_m that satisfies the conditions (P1)-(P4). The continuous property of the metric projection is a consequence of the assumptions on the space X (see [5]). It is easy to show that every continuous linear projection is a generalized projection.

Note that if there exists a projection $P : X \rightarrow M$, then M is closed. Moreover, a projection P is linear if and only if $\ker P$ is a subspace of X . Furthermore, every linear and continuous projection has properties from (P1) to (P4).

3.2. ORTHOGONAL PROJECTIONS

Let $P : X \rightarrow M$ be a generalized projection. We say that P is *orthogonal* if $(\ker P)^\perp = M$.

The following theorem holds.

Theorem 3.1. *Let M be a closed subspace of a uniformly convex Banach space X with a continuous semi-inner product. Let $P : X \rightarrow M$ be a projection (not necessarily linear) satisfying conditions (P2)-(P4). If P is homogeneous and*

$$\|P(x) - P(y)\| \leq \|x - y\| \text{ for all } x, y \in X,$$

then P is orthogonal.

Proof. Note that $P(0) = 0$, hence for $x \in X$ we have

$$\|P(x)\| \leq \|x\|. \quad (3.1)$$

Moreover, if $y - P(y) \in \ker P$, then $\lambda(y - P(y)) \in \ker P$. Indeed, using the homogeneity of P we obtain

$$P(\lambda(y - P(y))) = \lambda P(y - P(y)) = 0.$$

We shall show that $(\ker P)^\perp = M$. Setting x equal to $P(x) + \lambda(y - P(y))$ in (3.1) we obtain

$$\|P(P(x) + \lambda(y - P(y)))\| \leq \|P(x) + \lambda(y - P(y))\|,$$

hence

$$\|P(x)\| \leq \|P(x) + \lambda(y - P(y))\|$$

by virtue of Theorem 2.2, which is equivalent to the fact that $P(x)$ is orthogonal to every $z \in \ker P$.

Conversely, suppose that $x \in (\ker P)^\perp$. Then $[z, x] = 0$ for $z \in \ker P$. Hence $[x - P(x), x] = 0$ and

$$\|x\|^2 = [x - P(x) + P(x), x] = [x - P(x), x] + [P(x), x] \leq \|x\| \|P(x)\| \leq \|x\|^2.$$

By assumptions, it follows that $\|x\| = \|P(x)\|$ and $\|P(x)\| \|x\| = [P(x), x]$. By the strict convexity of X , we obtain $P(x) = x$, and so $x \in M$. \square

Now we can conclude that in a uniformly convex Banach space with a continuous semi-inner product every orthogonal projection is linear and satisfies the Lipschitz condition.

Theorem 3.2. *Assume that X is a uniformly convex Banach space with a continuous semi-inner product and M is a closed subspace of X . Let $P : X \rightarrow M$ be a generalized projection. If P is orthogonal, then P is linear and Lipschitz continuous with the constant equal to one.*

Proof. Note that $P(x_1 + x_2) - P(x_1) - P(x_2) \in M$ for $x_1, x_2 \in X$. Let

$$y = P(x_1 + x_2) - P(x_1) - P(x_2).$$

Then

$$\begin{aligned} \|P(x_1 + x_2) - P(x_1) - P(x_2)\|^2 &= [P(x_1 + x_2) - P(x_1) - P(x_2), y] = \\ &= -[(x_1 + x_2) - P(x_1 + x_2), y] + [x_1 - P(x_1), y] + [x_2 - P(x_2), y] = 0. \end{aligned}$$

Therefore, $P(x_1 + x_2) = P(x_1) + P(x_2)$. Now, let $y = P(\alpha x) - \alpha P(x)$. Then

$$\begin{aligned} \|P(\alpha x) - \alpha P(x)\|^2 &= [P(\alpha x) - \alpha P(x), y] = \\ &= -[\alpha x - P(\alpha x), y] + \alpha[x - P(x), y] = 0, \end{aligned}$$

hence $P(\alpha x) = \alpha P(x)$ for $x \in X$ and a scalar α .

We next show that $\|P\| = 1$. Let $x \in X$. Then $Px - x \in \ker P$ and

$$\|Px\|^2 = [Px, Px] = [Px - x + x, Px] = [Px - x, Px] + [x, Px] = [x, Px].$$

Using (2.4) we get

$$\|Px\| \leq \|x\|,$$

hence $\|P\| = 1$. □

Lemma 3.3. *Let $P: X \rightarrow M$ be an orthogonal projection. Then P is a unique orthogonal projection.*

Proof. Let P_i be an orthogonal projection ($i = 1, 2$). Hence $(\ker P_i)^\perp = M$ ($i = 1, 2$). Then $P_1x - P_2x \in M$ and

$$\begin{aligned} \|P_1x - P_2x\|^2 &= [P_1x - P_2x, P_1x - P_2x] = \\ &= [P_1x - x + x - P_2x, P_1x - P_2x] = \\ &= [P_1x - x, P_1x - P_2x] + [x - P_2x, P_1x - P_2x] = 0. \end{aligned}$$

Consequently, we conclude that $P_1x = P_2x$, which completes the proof. □

Lewicki and Skrzypek proved that the minimal projection onto a symmetric subspace of a smooth Banach space is unique (see [8, Theorem 2.9]). Now, we show an analogous theorem in a uniformly convex Banach space X with a continuous s.i.p. In its proof we use the structure of a semi-inner product.

Theorem 3.4. *Let X be a uniformly convex Banach space with continuous semi-inner product. Let M be a closed subspace of X . If there exists a linear projection $P: X \rightarrow M$ such that $\|P\| = 1$, then P is unique.*

Proof. Suppose that there exist linear projections P_1, P_2 such that $\|P_1\| = \|P_2\| = 1$. Then according to Theorem 3.1 they are orthogonal and hence $P_1 = P_2$ by Lemma 3.3. □

A stronger result is given below. Its proof is similar to those of Theorem 3.4, so we omit it.

Theorem 3.5. *Let M be a closed subspace of a uniformly convex Banach space X with continuous semi-inner product. If there exists a homogeneous projection $P : X \rightarrow M$ satisfying (P1)-(P4) such that*

$$\|P(x) - P(y)\| \leq \|x - y\| \text{ for all } x, y \in X,$$

then P is unique.

A linear subspace M is *one-complemented* if there exists a linear projection $P : X \rightarrow M$ such that $\|P\| = 1$.

Remark 3.6. Let M be a subspace of X such that $\dim M = 1$. Then from the Hahn-Banach theorem there exists a linear projection such that $\|P\| = 1$. Therefore, P is an orthogonal projection and M is one-complemented.

In this paper we give a necessary and sufficient condition for the set M^\perp to be a subspace of X . We also show when the equality

$$M = (M^\top)^\perp \tag{3.2}$$

holds.

Theorem 3.7. *Let X be a uniformly convex Banach space with continuous semi-inner product and M be a closed subspace of X . Then M is one-complemented if and only if there exists a closed subspace V of X such that $V^\perp = M$.*

Proof. Let M be one-complemented, hence there exists a linear, continuous projection $P : X \rightarrow M$ such that $\|P\| = 1$. By virtue of Theorem 3.1, P is an orthogonal projection, thus $(\ker P)^\perp = M$. Setting $V = \ker P$ we complete the first part of the proof.

Conversely, suppose that there exists a closed subspace V such that $V^\perp = M$. Then $X = V \oplus V^\perp = V \oplus M$. We define an orthogonal projection $P_V : X \rightarrow M$ such that

$$P_V x = P_V(x_V + x_M) = x_M,$$

where $x_V \in V$, $x_M \in M$. This finishes the proof. □

The following theorem gives a characterization of one-complemented spaces.

Theorem 3.8. *A subspace M of a uniformly convex Banach space X with continuous semi-inner product is one-complemented if and only if*

$$M = (M^\top)^\perp. \tag{3.3}$$

Moreover, if (3.3) holds, then a projection $P : X \rightarrow M$ given by

$$P(x_M + x_{M^\top}) = x_M, \quad x_M \in M, \quad x_{M^\top} \in M^\top, \tag{3.4}$$

is the only projection with the norm equal to one.

Proof. From Theorem 3.7 we deduce that exists a closed subspace V of X such that

$$V^\perp = M. \tag{3.5}$$

Hence

$$V = (V^\perp)^\top = M^\top. \tag{3.6}$$

From (3.5) and (3.6) we get $M = V^\perp = (M^\top)^\perp$. By Theorem 2.5, we deduce that

$$X = V \oplus V^\perp = M \oplus M^\top.$$

Conversely, let $M = (M^\top)^\perp$. Hence

$$X = (M^\top)^\perp \oplus M^\top = M \oplus M^\top. \tag{3.7}$$

From (3.7) it easy to see that a linear projection $P: X \rightarrow M$ given by the formula (3.4) is orthogonal. \square

3.3. CO-ORTHOGONAL PROJECTIONS

A projection P is called *co-orthogonal* if $M^\perp = \ker P$. Note that not every co-orthogonal projection is linear, for example a metric projection.

We start with the following theorem.

Theorem 3.9. *Let M be a closed proper subspace of a uniformly convex Banach space X with a continuous semi-inner product. Let $P: X \rightarrow M$ be a linear projection. Then the following conditions are equivalent:*

- (i) P is co-orthogonal,
- (ii) $\|Id - P\| = 1$.

Proof. Suppose that the linear projection $P: X \rightarrow M$ is co-orthogonal.

Let $x \in X$. Then $x - Px \in \ker P$ and

$$\begin{aligned} \|x - Px\|^2 &= [x - Px, x - Px] = [x, x - Px] - [Px, x - Px] = \\ &= [x, x - Px] \leq \|x\| \|x - Px\|. \end{aligned}$$

Therefore, we have

$$\|x - Px\| \leq \|x\|,$$

hence $\|Id - P\| = 1$.

Conversely, suppose that for each $x \in X$ we get

$$\|x - Px\| \leq \|x\|. \tag{3.8}$$

We now show that $\ker P = M^\perp$. Setting x equal to $x - Px + \lambda Py$ in (3.8) we obtain

$$\|x - Px + \lambda Py - P(x - Px + \lambda Py)\| \leq \|x - Px + \lambda Py\|,$$

hence

$$\|x - Px\| \leq \|x - Px + \lambda Py\|$$

by virtue of Theorem 2.2, which is equivalent to $x - Px$ is orthogonal to every $z \in M$.

On the other hand, suppose that $x \in M^\perp$. Then $[z, x] = 0$ for $z \in M$. Hence $[Px, x] = 0$ and

$$\|x\|^2 = [x - Px, x].$$

Therefore,

$$\|x\|^2 = [x - Px, x] \leq \|x - Px\| \|x\| \leq \|x\|^2.$$

By assumption it follows $\|x - Px\| = \|x\|$. By Lemma 2.4, we obtain that $x - Px = x$, therefore $x \in \ker P$. \square

Let us now characterize the linearity of the set of M^\perp . We present the following theorem.

Theorem 3.10. *Let M be a closed proper subspace of a uniformly convex Banach space X with a continuous semi-inner product. Then the following conditions are equivalent:*

- (i) *the set M^\perp is a linear space,*
- (ii) *there exists a linear projection $P: X \rightarrow M$ such that $\|Id - P\| = 1$.*

Proof. If M^\perp is a linear subspace, we get $X = M \oplus M^\perp$. Then it easy to see that linear projection $P: X \rightarrow M$ given by the formula

$$Px = P(x_M + x_{M^\perp}) = x_M, \quad x \in X \tag{3.9}$$

is co-orthogonal.

Conversely, if a linear projection $P: X \rightarrow M$ is co-orthogonal, then $\ker P = M^\perp$. \square

Finally, we will prove the following lemma.

Lemma 3.11 ([8]). *Let $P: X \rightarrow M$ be a co-orthogonal linear projection. Then P is a unique co-orthogonal linear projection.*

Proof. Let P_i be a co-orthogonal projection, hence $\ker P_i = M^\perp$ ($i = 1, 2$). Then $P_1x - P_2x \in M$ and $x - P_1x \in M^\perp$, $x - P_2x \in M^\perp$. Since M^\perp is a subspace of X , then $P_1x - P_2x \in M^\perp$. According to (2.7) we conclude $P_1x = P_2x$, which completes the proof. \square

Let M be a closed proper subspace of a uniformly convex Banach space X with a continuous semi-inner product.

We say that M is *one-co-complemented* if there exists a linear projection $P: X \rightarrow M$ such that $\|Id - P\| = 1$.

From the above discussion we obtain the following result.

Theorem 3.12. *Let M be a closed proper subspace of a uniformly convex Banach space X with a continuous semi-inner product. Then M is one-co-complemented if and only if M^\perp is a vector space. Moreover, if M^\perp is a linear space, then a projection $P: X \rightarrow M$ given by*

$$P(x_M + x_{M^\perp}) = x_M,$$

is the only projection which satisfies the equality $\|Id - P\| = 1$.

Acknowledgments

The authors would like to express their thanks to prof. Petru Cojuhari for useful comments and discussions on the paper.

The authors also would like to thank prof. Grzegorz Lewicki for his valuable comments and discussions regarding the subject of this paper.

The authors were supported by the Polish Ministry of Sciences and Higher Education.

REFERENCES

- [1] J.A. Clarkson, *Uniformly convex spaces*, Trans. Amer. Math. Soc. **40** (1936), 396–414.
- [2] P.A. Cojuhari, *Generalized Hardy type inequalities and some applications to spectral theory*, Operator Theory, Operator Algebras and Related Topics, Theta Found., Bucharest, (1997), 79–99.
- [3] P.A. Cojuhari, M.A Nowak, *Projection-iterative methods for a class of difference equations*, Integral Equations Operator Theory **64** (2009), 155–175.
- [4] J.R. Giles, *Classes of semi-inner-product spaces*, Trans. Amer. Math. Soc., **129** (1961), 436–446.
- [5] H. Hudzik, W. Kowalewski, G. Lewicki, *Approximative compactness and full rotundity in Musielak-Orlicz spaces and Lorentz-Orlicz spaces*, Z. Anal. Anwendungen (Journal for Analysis and Its Applications) **25** (2006) 2, 163–192.
- [6] R.C. James, *Orthogonality and linear functionals in normed linear spaces*, Trans. Amer. Math. Soc. **61** (1947), 265–292.
- [7] A. Kufner, *Weighted Sobolev Spaces*, Teubner-Texte zur Mathematik, Band 31, 1980.
- [8] G. Lewicki, L. Skrzypek, *Chalmers-Metcalf operator and uniqueness of minimal projections*, J. Approx. Theory **148** (2007), 71–91.
- [9] J. Lindenstrauss, L. Tzafriri, *Classical Banach spaces I*, EMG 92 Springer Verlag, 1977.
- [10] G. Lumer, *Semi-inner-product spaces*, Trans. Amer. Math. Soc. **100** (1961), 29–43.
- [11] V. Smulian, *Sur la dérivabilité de la norme dans l'espace de Banach*, Dokl. Akad. Nauk SSSR **27** (1940), 643–648.

Ewa Szlachtowska
szlachto@agh.edu.pl

AGH University of Science and Technology
Faculty of Applied Mathematics
al. Mickiewicza 30, 30-059 Krakow, Poland

Dominik Mielczarek
dmielcza@wms.mat.agh.edu.pl

AGH University of Science and Technology
Faculty of Applied Mathematics
al. Mickiewicza 30, 30-059 Krakow, Poland

Received: February 19, 2012.

Revised: March 18, 2012.

Accepted: March 20, 2012.